

Chapter 1

Equations and Inequalities

Section 1.1

1. Distributive

2. Zero-Product

3. $\{x | x \neq 4\}$

4. False. Multiplying both sides of an equation by zero will not result in an equivalent equation.

5. identity

6. linear; first-degree

7. False. The solution is $\frac{8}{3}$.

8. True

9. b

10. d

11. $7x = 21$

$$\frac{7x}{7} = \frac{21}{7}$$

$$x = 3$$

The solution set is $\{3\}$.

12. $6x = -24$

$$\frac{6x}{6} = \frac{-24}{6}$$

$$x = -4$$

The solution set is $\{-4\}$.

13. $3x + 15 = 0$

$$3x + 15 - 15 = 0 - 15$$

$$3x = -15$$

$$\frac{3x}{3} = \frac{-15}{3}$$

$$x = -5$$

The solution set is $\{-5\}$.

14. $6x + 18 = 0$

$$6x + 18 - 18 = 0 - 18$$

$$6x = -18$$

$$\frac{6x}{6} = \frac{-18}{6}$$

$$x = -3$$

The solution set is $\{-3\}$.

15. $2x - 3 = 0$

$$2x - 3 + 3 = 0 + 3$$

$$2x = 3$$

$$\frac{2x}{2} = \frac{3}{2}$$

$$x = \frac{3}{2}$$

The solution set is $\left\{\frac{3}{2}\right\}$.

16. $3x + 4 = 0$

$$3x + 4 - 4 = 0 - 4$$

$$3x = -4$$

$$\frac{3x}{3} = \frac{-4}{3}$$

$$x = -\frac{4}{3}$$

The solution set is $\left\{-\frac{4}{3}\right\}$.

17. $\frac{1}{3}x = \frac{5}{12}$

$$3\left(\frac{1}{3}x\right) = 3\left(\frac{5}{12}\right)$$

$$x = \frac{5}{4}$$

The solution set is $\left\{\frac{5}{4}\right\}$.

18. $\frac{2}{3}x = \frac{9}{2}$
 $6\left(\frac{2}{3}x\right) = 6\left(\frac{9}{2}\right)$
 $4x = 27$
 $\frac{4x}{4} = \frac{27}{4}$
 $x = \frac{27}{4}$

The solution set is $\left\{\frac{27}{4}\right\}$.

19. $3x + 4 = x$
 $3x + 4 - 4 = x - 4$
 $3x = x - 4$
 $3x - x = x - 4 - x$
 $2x = -4$
 $\frac{2x}{2} = \frac{-4}{2}$
 $x = -2$

The solution set is $\{-2\}$.

20. $2x + 9 = 5x$
 $2x + 9 - 9 = 5x - 9$
 $2x = 5x - 9$
 $2x - 5x = 5x - 9 - 5x$
 $-3x = -9$
 $\frac{-3x}{-3} = \frac{-9}{-3}$
 $x = 3$

The solution set is $\{3\}$.

21. $2t - 6 = 3 - t$
 $2t - 6 + 6 = 3 - t + 6$
 $2t = 9 - t$
 $2t + t = 9 - t + t$
 $3t = 9$
 $\frac{3t}{3} = \frac{9}{3}$
 $t = 3$

The solution set is $\{3\}$.

22. $5y + 6 = -18 - y$
 $5y + 6 - 6 = -18 - y - 6$
 $5y = -y - 24$
 $5y + y = -y - 24 + y$
 $6y = -24$
 $\frac{6y}{6} = \frac{-24}{6}$
 $y = -4$

The solution set is $\{-4\}$.

23. $6 - x = 2x + 9$
 $6 - x - 6 = 2x + 9 - 6$
 $-x = 2x + 3$
 $-x - 2x = 2x + 3 - 2x$
 $-3x = 3$
 $\frac{-3x}{-3} = \frac{3}{-3}$
 $x = -1$

The solution set is $\{-1\}$.

24. $3 - 2x = 2 - x$
 $3 - 2x - 3 = 2 - x - 3$
 $-2x = -x - 1$
 $-2x + x = -x - 1 + x$
 $-x = -1$
 $\frac{-x}{-1} = \frac{-1}{-1}$
 $x = 1$

The solution set is $\{1\}$.

25. $3 + 2n = 4n + 7$
 $3 + 2n - 3 = 4n + 7 - 3$
 $2n = 4n + 4$
 $2n - 4n = 4n + 4 - 4n$
 $-2n = 4$
 $\frac{-2n}{-2} = \frac{4}{-2}$
 $n = -2$

The solution set is $\{-2\}$.

Chapter 1: Equations and Inequalities

26. $6 - 2m = 3m + 1$

$$6 - 2m - 6 = 3m + 1 - 6$$

$$-2m = 3m - 5$$

$$-2m - 3m = 3m - 5 - 3m$$

$$-5m = -5$$

$$\frac{-5m}{-5} = \frac{-5}{-5}$$

$$m = 1$$

The solution set is $\{1\}$.

27. $2(3 + 2x) = 3(x - 4)$

$$6 + 4x = 3x - 12$$

$$6 + 4x - 6 = 3x - 12 - 6$$

$$4x = 3x - 18$$

$$4x - 3x = 3x - 18 - 3x$$

$$x = -18$$

The solution set is $\{-18\}$.

28. $3(2 - x) = 2x - 1$

$$6 - 3x = 2x - 1$$

$$6 - 3x - 6 = 2x - 1 - 6$$

$$-3x = 2x - 7$$

$$-3x - 2x = 2x - 7 - 2x$$

$$-5x = -7$$

$$\frac{-5x}{-5} = \frac{-7}{-5}$$

$$x = \frac{7}{5}$$

The solution set is $\left\{\frac{7}{5}\right\}$.

29. $8x - (3x + 2) = 3x - 10$

$$8x - 3x - 2 = 3x - 10$$

$$5x - 2 = 3x - 10$$

$$5x - 2 + 2 = 3x - 10 + 2$$

$$5x = 3x - 8$$

$$5x - 3x = 3x - 8 - 3x$$

$$2x = -8$$

$$\frac{2x}{2} = \frac{-8}{2}$$

$$x = -4$$

The solution set is $\{-4\}$.

30. $7 - (2x - 1) = 10$

$$7 - 2x + 1 = 10$$

$$8 - 2x = 10$$

$$8 - 2x - 8 = 10 - 8$$

$$-2x = 2$$

$$\frac{-2x}{-2} = \frac{2}{-2}$$

$$x = -1$$

The solution set is $\{-1\}$.

31. $\frac{3}{2}x + 2 = \frac{1}{2} - \frac{1}{2}x$

$$2\left(\frac{3}{2}x + 2\right) = 2\left(\frac{1}{2} - \frac{1}{2}x\right)$$

$$3x + 4 = 1 - x$$

$$3x + 4 - 4 = 1 - x - 4$$

$$3x = -3 - x$$

$$3x + x = -3 - x + x$$

$$4x = -3$$

$$\frac{4x}{4} = \frac{-3}{4}$$

$$x = -\frac{3}{4}$$

The solution set is $\left\{-\frac{3}{4}\right\}$.

32. $\frac{1}{3}x = 2 - \frac{2}{3}x$

$$3\left(\frac{1}{3}x\right) = 3\left(2 - \frac{2}{3}x\right)$$

$$x = 6 - 2x$$

$$x + 2x = 6 - 2x + 2x$$

$$3x = 6$$

$$\frac{3x}{3} = \frac{6}{3}$$

$$x = 2$$

The solution set is $\{2\}$.

33. $\frac{1}{2}x - 5 = \frac{3}{4}x$

$$4\left(\frac{1}{2}x - 5\right) = 4\left(\frac{3}{4}x\right)$$

$$2x - 20 = 3x$$

$$2x - 20 - 2x = 3x - 2x$$

$$-20 = x$$

$$x = -20$$

The solution set is $\{-20\}$.

34. $1 - \frac{1}{2}x = 6$

$$2\left(1 - \frac{1}{2}x\right) = 2(6)$$

$$2 - x = 12$$

$$2 - x - 2 = 12 - 2$$

$$-x = 10$$

$$\frac{-x}{-1} = \frac{10}{-1}$$

$$x = -10$$

The solution set is $\{-10\}$.

35. $\frac{2}{3}p = \frac{1}{2}p + \frac{1}{3}$

$$6\left(\frac{2}{3}p\right) = 6\left(\frac{1}{2}p + \frac{1}{3}\right)$$

$$4p = 3p + 2$$

$$4p - 3p = 3p + 2 - 3p$$

$$p = 2$$

The solution set is $\{2\}$.

36. $\frac{1}{2} - \frac{1}{3}p = \frac{4}{3}$

$$6\left(\frac{1}{2} - \frac{1}{3}p\right) = 6\left(\frac{4}{3}\right)$$

$$3 - 2p = 8$$

$$3 - 2p - 3 = 8 - 3$$

$$-2p = 5$$

$$\frac{-2p}{-2} = \frac{5}{-2}$$

$$p = -\frac{5}{2}$$

The solution set is $\left\{-\frac{5}{2}\right\}$.

37. $0.9t = 0.4 + 0.1t$

$$0.9t - 0.1t = 0.4 + 0.1t - 0.1t$$

$$0.8t = 0.4$$

$$\frac{0.8t}{0.8} = \frac{0.4}{0.8}$$

$$t = 0.5$$

The solution set is $\{0.5\}$.

38. $0.9t = 1 + t$

$$0.9t - t = 1 + t - t$$

$$-0.1t = 1$$

$$\frac{-0.1t}{-0.1} = \frac{1}{-0.1}$$

$$t = -10$$

The solution set is $\{-10\}$.

39. $\frac{x+1}{3} + \frac{x+2}{7} = 2$

$$21\left(\frac{x+1}{3} + \frac{x+2}{7}\right) = 21(2)$$

$$7(x+1) + (3)(x+2) = 42$$

$$7x + 7 + 3x + 6 = 42$$

$$10x + 13 = 42$$

$$10x + 13 - 13 = 42 - 13$$

$$10x = 29$$

$$\frac{10x}{10} = \frac{29}{10}$$

$$x = \frac{29}{10}$$

The solution set is $\left\{\frac{29}{10}\right\}$.

40. $\frac{2x+1}{3} + 16 = 3x$

$$3\left(\frac{2x+1}{3} + 16\right) = 3(3x)$$

$$2x + 1 + 48 = 9x$$

$$2x + 49 = 9x$$

$$2x + 49 - 2x = 9x - 2x$$

$$49 = 7x$$

$$\frac{49}{7} = \frac{7x}{7}$$

$$x = 7$$

The solution set is $\{7\}$.

Chapter 1: Equations and Inequalities

41. $\frac{2}{y} + \frac{4}{y} = 3$
 $y\left(\frac{2}{y} + \frac{4}{y}\right) = y(3)$
 $2+4=3y$
 $6=3y$
 $\frac{6}{3}=\frac{3y}{3}$
 $2=y$

Since $y=2$ does not cause a denominator to equal zero, the solution set is $\{2\}$.

42. $\frac{4}{y} - 5 = \frac{5}{2y}$
 $2y\left(\frac{4}{y} - 5\right) = 2y\left(\frac{5}{2y}\right)$
 $8-10y=5$
 $8-10y-8=5-8$
 $-10y=-3$
 $\frac{-10y}{-10}=\frac{-3}{-10}$
 $y=\frac{3}{10}$

Since $y=\frac{3}{10}$ does not cause a denominator to equal zero, the solution set is $\left\{\frac{3}{10}\right\}$.

43. $\frac{1}{2} + \frac{2}{x} = \frac{3}{4}$
 $4x\left(\frac{1}{2} + \frac{2}{x}\right) = 4x\left(\frac{3}{4}\right)$
 $2x+8=3x$
 $2x+8-2x=3x-2x$
 $8=x$

Since $x=8$ does not cause any denominator to equal zero, the solution set is $\{8\}$.

44. $\frac{3}{x} - \frac{1}{3} = \frac{1}{6}$
 $6x\left(\frac{3}{x} - \frac{1}{3}\right) = 6x\left(\frac{1}{6}\right)$
 $18-2x=x$
 $18-2x+2x=x+2x$
 $18=3x$
 $\frac{18}{3}=\frac{3x}{3}$
 $6=x$

Since $x=6$ does not cause a denominator to equal zero, the solution set is $\{6\}$.

45. $(x+7)(x-1) = (x+1)^2$
 $x^2 + 6x - 7 = x^2 + 2x + 1$
 $x^2 + 6x - 7 - x^2 = x^2 + 2x + 1 - x^2$
 $6x - 7 = 2x + 1$
 $6x - 7 + 7 = 2x + 1 + 7$
 $6x = 2x + 8$
 $6x - 2x = 2x + 8 - 2x$
 $4x = 8$
 $\frac{4x}{4}=\frac{8}{4}$
 $x=2$
The solution set is $\{2\}$.

46. $(x+2)(x-3) = (x+3)^2$
 $x^2 - x - 6 = x^2 + 6x + 9$
 $x^2 - x - 6 - x^2 = x^2 + 6x + 9 - x^2$
 $-x - 6 = 6x + 9$
 $-x - 6 + 6 = 6x + 9 + 6$
 $-x = 6x + 15$
 $-x - 6x = 6x + 15 - 6x$
 $-7x = 15$
 $\frac{-7x}{-7}=\frac{15}{-7}$
 $x=-\frac{15}{7}$
The solution set is $\left\{-\frac{15}{7}\right\}$.

47. $x(2x - 3) = (2x + 1)(x - 4)$

$$2x^2 - 3x = 2x^2 - 7x - 4$$

$$2x^2 - 3x - 2x^2 = 2x^2 - 7x - 4 - 2x^2$$

$$-3x = -7x - 4$$

$$-3x + 7x = -7x - 4 + 7x$$

$$4x = -4$$

$$\frac{4x}{4} = \frac{-4}{4}$$

$$x = -1$$

The solution set is $\{-1\}$.

48. $x(1+2x) = (2x-1)(x-2)$

$$x + 2x^2 = 2x^2 - 5x + 2$$

$$x + 2x^2 - 2x^2 = 2x^2 - 5x + 2 - 2x^2$$

$$x = -5x + 2$$

$$x + 5x = -5x + 2 + 5x$$

$$6x = 2$$

$$\frac{6x}{6} = \frac{2}{6}$$

$$x = \frac{1}{3}$$

The solution set is $\left\{\frac{1}{3}\right\}$.

49. $z(z^2 + 1) = 3 + z^3$

$$z^3 + z = 3 + z^3$$

$$z^3 + z - z^3 = 3 + z^3 - z^3$$

$$z = 3$$

The solution set is $\{3\}$.

50. $w(4 - w^2) = 8 - w^3$

$$4w - w^3 = 8 - w^3$$

$$4w - w^3 + w^3 = 8 - w^3 + w^3$$

$$4w = 8$$

$$\frac{4w}{4} = \frac{8}{4}$$

$$w = 2$$

The solution set is $\{2\}$.

51. $\frac{x}{x-2} + 3 = \frac{2}{x-2}$

$$\left(\frac{x}{x-2} + 3\right)(x-2) = \left(\frac{2}{x-2}\right)(x-2)$$

$$x + 3(x-2) = 2$$

$$x + 3x - 6 = 2$$

$$4x - 6 = 2$$

$$4x - 6 + 6 = 2 + 6$$

$$4x = 8$$

$$\frac{4x}{4} = \frac{8}{4}$$

$$x = 2$$

Since $x = 2$ causes a denominator to equal zero, we must discard it. Therefore the original equation has no solution.

52. $\frac{2x}{x+3} = \frac{-6}{x+3} - 2$

$$\left(\frac{2x}{x+3}\right)(x+3) = \left(\frac{-6}{x+3} - 2\right)(x+3)$$

$$2x = -6 - (2)(x+3)$$

$$2x = -6 - 2x - 6$$

$$2x = -12 - 2x$$

$$2x + 2x = -12 - 2x + 2x$$

$$4x = -12$$

$$\frac{4x}{4} = \frac{-12}{4}$$

$$x = -3$$

Since $x = -3$ causes a denominator to equal zero, we must discard it. Therefore the original equation has no solution.

Chapter 1: Equations and Inequalities

53.

$$\frac{2x}{x^2 - 4} = \frac{4}{x^2 - 4} - \frac{3}{x + 2}$$

$$\frac{2x}{(x+2)(x-2)} = \frac{4}{(x+2)(x-2)} - \frac{3}{x+2}$$

$$\left(\frac{2x}{(x+2)(x-2)}\right)(x+2)(x-2) = \left(\frac{4}{(x+2)(x-2)} - \frac{3}{x+2}\right)(x+2)(x-2)$$

$$2x = 4 - 3(x-2)$$

$$2x = 4 - 3x + 6$$

$$2x = 10 - 3x$$

$$2x + 3x = 10 - 3x + 3x$$

$$5x = 10$$

$$\frac{5x}{5} = \frac{10}{5}$$

$$x = 2$$

Since $x = 2$ causes a denominator to equal zero, we must discard it. Therefore the original equation has no solution.

54.

$$\frac{x}{x^2 - 9} + \frac{4}{x+3} = \frac{3}{x^2 - 9}$$

$$\frac{x}{(x+3)(x-3)} + \frac{4}{x+3} = \frac{3}{(x+3)(x-3)}$$

$$\left(\frac{x}{(x+3)(x-3)} + \frac{4}{x+3}\right)(x+3)(x-3) = \left(\frac{3}{(x+3)(x-3)}\right)(x+3)(x-3)$$

$$x + 4(x-3) = 3$$

$$x + 4x - 12 = 3$$

$$5x - 12 = 3$$

$$5x - 12 + 12 = 3 + 12$$

$$5x = 15$$

$$\frac{5x}{5} = \frac{15}{5}$$

$$x = 3$$

Since $x = 3$ causes a denominator to equal zero, we must discard it. Therefore the original equation has no solution.

55.

$$\frac{x}{x+2} = \frac{3}{2}$$

$$2(x+2)\left(\frac{x}{x+2}\right) = 2(x+2)\left(\frac{3}{2}\right)$$

$$2x = 3(x+2)$$

$$2x = 3x + 6$$

$$2x - 3x = 3x + 6 - 3x$$

$$-x = 6$$

$$\frac{-x}{-1} = \frac{6}{-1}$$

$$x = -6$$

Since $x = -6$ does not cause any denominator to equal zero, the solution set is $\{-6\}$.

56.

$$\frac{3x}{x-1} = 2$$

$$\left(\frac{3x}{x-1}\right)(x-1) = 2(x-1)$$

$$3x = 2x - 2$$

$$3x - 2x = 2x - 2 - 2x$$

$$x = -2$$

Since $x = -2$ does not cause any denominator to equal zero, the solution set is $\{-2\}$.

57. $\frac{5}{2x-3} = \frac{3}{x+5}$

$$\left(\frac{5}{2x-3}\right)(2x-3)(x+5) = \left(\frac{3}{x+5}\right)(2x-3)(x+5)$$

$$5(x+5) = 3(2x-3)$$

$$5x+25 = 6x-9$$

$$5x+25-6x = 6x-9-6x$$

$$25-x = -9$$

$$25-x-25 = -9-25$$

$$-x = -34$$

$$\frac{-x}{-1} = \frac{-34}{-1}$$

$$x = 34$$

Since $x = 34$ does not cause any denominator to equal zero, the solution is $\{34\}$.

59. $\frac{6t+7}{4t-1} = \frac{3t+8}{2t-4}$

$$\left(\frac{6t+7}{4t-1}\right)(4t-1)(2t-4) = \left(\frac{3t+8}{2t-4}\right)(4t-1)(2t-4)$$

$$(6t+7)(2t-4) = (3t+8)(4t-1)$$

$$12t^2 - 24t + 14t - 28 = 12t^2 - 3t + 32t - 8$$

$$12t^2 - 10t - 28 = 12t^2 + 29t - 8$$

$$12t^2 - 10t - 28 - 12t^2 = 12t^2 + 29t - 8 - 12t^2$$

$$-10t - 28 = 29t - 8$$

$$-10t - 28 - 29t = 29t - 8 - 29t$$

$$-28 - 39t = -8$$

$$-28 - 39t + 28 = -8 + 28$$

$$-39t = 20$$

$$\frac{-39t}{-39} = \frac{20}{-39}$$

$$t = -\frac{20}{39}$$

Since $t = -\frac{20}{39}$ does not cause any denominator to equal zero, the solution set is $\left\{-\frac{20}{39}\right\}$.

58. $\frac{-4}{x+4} = \frac{-3}{x+6}$

$$\left(\frac{-4}{x+4}\right)(x+6)(x+4) = \left(\frac{-3}{x+6}\right)(x+6)(x+4)$$

$$-4(x+6) = -3(x+4)$$

$$-4x - 24 = -3x - 12$$

$$-4x - 24 + 4x = -3x - 12 + 4x$$

$$-24 = -12 + x$$

$$-24 + 12 = -12 + x + 12$$

$$-12 = x$$

Since $x = -12$ does not cause any denominator to equal zero, the solution set is $\{-12\}$.

Chapter 1: Equations and Inequalities

60.

$$\begin{aligned} \frac{8w+5}{10w-7} &= \frac{4w-3}{5w+7} \\ \left(\frac{8w+5}{10w-7}\right)(10w-7)(5w+7) &= \left(\frac{4w-3}{5w+7}\right)(10w-7)(5w+7) \\ (8w+5)(5w+7) &= (4w-3)(10w-7) \\ 40w^2 + 56w + 25w + 35 &= 40w^2 - 28w - 30w + 21 \\ 40w^2 + 81w + 35 &= 40w^2 - 58w + 21 \\ 40w^2 + 81w + 35 - 40w^2 &= 40w^2 - 58w + 21 - 40w^2 \\ 81w + 35 &= -58w + 21 \\ 81w + 35 + 58w &= -58w + 21 + 58w \\ 139w + 35 &= 21 \\ 139w + 35 - 35 &= 21 - 35 \\ 139w &= -14 \\ \frac{139w}{139} &= \frac{-14}{139} \\ w &= -\frac{14}{139} \end{aligned}$$

Since $w = -\frac{14}{139}$ does not cause any denominator to equal zero, the solution set is $\left\{-\frac{14}{139}\right\}$.

61.

$$\begin{aligned} \frac{4}{x-2} &= \frac{-3}{x+5} + \frac{7}{(x+5)(x-2)} \\ \left(\frac{4}{x-2}\right)(x+5)(x-2) &= \left(\frac{-3}{x+5} + \frac{7}{(x+5)(x-2)}\right)(x+5)(x-2) \\ 4(x+5) &= -3(x-2) + 7 \\ 4x + 20 &= -3x + 6 + 7 \\ 4x + 20 &= -3x + 13 \\ 4x + 20 + 3x &= -3x + 13 + 3x \\ 7x + 20 &= 13 \\ 7x + 20 - 20 &= 13 - 20 \\ 7x &= -7 \\ \frac{7x}{7} &= \frac{-7}{7} \\ x &= -1 \end{aligned}$$

Since $x = -1$ does not cause any denominator to equal zero, the solution set is $\{-1\}$.

62.
$$\frac{-4}{2x+3} + \frac{1}{x-1} = \frac{1}{(2x+3)(x-1)}$$

$$\left(\frac{-4}{2x+3} + \frac{1}{x-1}\right)(2x+3)(x-1) = \left(\frac{1}{(2x+3)(x-1)}\right)(2x+3)(x-1)$$

$$-4(x-1) + 1(2x+3) = 1$$

$$-4x + 4 + 2x + 3 = 1$$

$$-2x + 7 = 1$$

$$-2x + 7 - 7 = 1 - 7$$

$$-2x = -6$$

$$\frac{-2x}{-2} = \frac{-6}{-2}$$

$$x = 3$$

Since $x = 3$ does not cause any denominator to equal zero, the solution set is $\{3\}$.

63.
$$\frac{2}{y+3} + \frac{3}{y-4} = \frac{5}{y+6}$$

$$\left(\frac{2}{y+3} + \frac{3}{y-4}\right)(y+3)(y-4)(y+6) = \left(\frac{5}{y+6}\right)(y+3)(y-4)(y+6)$$

$$2(y-4)(y+6) + 3(y+3)(y+6) = 5(y+3)(y-4)$$

$$2(y^2 + 6y - 4y - 24) + 3(y^2 + 6y + 3y + 18) = 5(y^2 - 4y + 3y - 12)$$

$$2(y^2 + 2y - 24) + 3(y^2 + 9y + 18) = 5(y^2 - y - 12)$$

$$2y^2 + 4y - 48 + 3y^2 + 27y + 54 = 5y^2 - 5y - 60$$

$$5y^2 + 31y + 6 = 5y^2 - 5y - 60$$

$$5y^2 + 31y + 6 - 5y^2 = 5y^2 - 5y - 60 - 5y^2$$

$$31y + 6 = -5y - 60$$

$$31y + 6 + 5y = -5y - 60 + 5y$$

$$36y + 6 = -60$$

$$36y + 6 - 6 = -60 - 6$$

$$36y = -66$$

$$\frac{36y}{36} = \frac{-66}{36}$$

$$y = -\frac{11}{6}$$

Since $y = -\frac{11}{6}$ does not cause any denominator to equal zero, the solution set is $\left\{-\frac{11}{6}\right\}$.

Chapter 1: Equations and Inequalities

64.

$$\begin{aligned} \frac{5}{5z-11} + \frac{4}{2z-3} &= \frac{-3}{5-z} \\ \left(\frac{5}{5z-11} + \frac{4}{2z-3} \right) (5z-11)(2z-3)(5-z) &= \left(\frac{-3}{5-z} \right) (5z-11)(2z-3)(5-z) \\ 5(2z-3)(5-z) + 4(5z-11)(5-z) &= -3(5z-11)(2z-3) \\ 5(10z-2z^2-15+3z) + 4(25z-5z^2-55+11z) &= -3(10z^2-15z-22z+33) \\ 5(-2z^2+13z-15) + 4(-5z^2+36z-55) &= -3(10z^2-37z+33) \\ -10z^2+65z-75-20z^2+144z-220 &= -30z^2+111z-99 \\ -30z^2+209z-295 &= -30z^2+111z-99 \\ -30z^2+209z-295+30z^2 &= -30z^2+111z-99+30z^2 \\ 209z-295 &= 111z-99 \\ 209z-295-209z &= 111z-99-209z \\ -295 &= -98z-99 \\ -295+99 &= -98z-99+99 \\ -196 &= -98z \\ \frac{-196}{-98} &= \frac{-118z}{-98} \\ 2 &= z \end{aligned}$$

Since $z = 2$ does not cause any denominator to equal zero, the solution set is $\{2\}$.

65.

$$\begin{aligned} \frac{x}{x^2-1} - \frac{x+3}{x^2-x} &= \frac{-3}{x^2+x} \\ \frac{x}{(x+1)(x-1)} - \frac{x+3}{x(x-1)} &= \frac{-3}{x(x+1)} \\ \left(\frac{x}{(x+1)(x-1)} - \frac{x+3}{x(x-1)} \right) x(x+1)(x-1) &= \left(\frac{-3}{x(x+1)} \right) x(x+1)(x-1) \\ (x)(x) - (x+3)(x+1) &= -3(x-1) \\ x^2 - (x^2 + x + 3x + 3) &= -3x + 3 \\ x^2 - (x^2 + 4x + 3) &= -3x + 3 \\ x^2 - x^2 - 4x - 3 &= -3x + 3 \\ -4x - 3 &= -3x + 3 \\ -4x - 3 + 4x &= -3x + 3 + 4x \\ -3 &= 3 + x \\ -3 - 3 &= 3 + x - 3 \\ -6 &= x \end{aligned}$$

Since $x = -6$ does not cause any denominator to equal zero, the solution set is $\{-6\}$.

66.

$$\frac{x+1}{x^2+2x} - \frac{x+4}{x^2+x} = \frac{-3}{x^2+3x+2}$$

$$\frac{x+1}{x(x+2)} - \frac{x+4}{x(x+1)} = \frac{-3}{(x+2)(x+1)}$$

$$\left(\frac{x+1}{x(x+2)} - \frac{x+4}{x(x+1)} \right) x(x+2)(x+1) = \left(\frac{-3}{(x+2)(x+1)} \right) x(x+2)(x+1)$$

$$(x+1)(x+1) - (x+4)(x+2) = -3x$$

$$(x^2 + x + x + 1) - (x^2 + 2x + 4x + 8) = -3x$$

$$x^2 + 2x + 1 - (x^2 + 6x + 8) = -3x$$

$$x^2 + 2x + 1 - x^2 - 6x - 8 = -3x$$

$$2x + 1 - 6x - 8 = -3x$$

$$-4x - 7 = -3x$$

$$-4x - 7 + 4x = -3x + 4x$$

$$-7 = x$$

Since $x = -7$ does not cause any denominator to equal zero, the solution set is $\{-7\}$.

67.

$$3.2x + \frac{21.3}{65.871} = 19.23$$

$$3.2x + \frac{21.3}{65.871} - \frac{21.3}{65.871} = 19.23 - \frac{21.3}{65.871}$$

$$3.2x = 19.23 - \frac{21.3}{65.871}$$

$$\left(\frac{1}{3.2} \right) (3.2x) = \left(19.23 - \frac{21.3}{65.871} \right) \left(\frac{1}{3.2} \right)$$

$$x = \left(19.23 - \frac{21.3}{65.871} \right) \left(\frac{1}{3.2} \right) \approx 5.91$$

The solution set is approximately $\{5.91\}$.

68.

$$6.2x - \frac{19.1}{83.72} = 0.195$$

$$6.2x - \frac{19.1}{83.72} + \frac{19.1}{83.72} = 0.195 + \frac{19.1}{83.72}$$

$$6.2x = 0.195 + \frac{19.1}{83.72}$$

$$\left(\frac{1}{6.2} \right) (6.2x) = \left(0.195 + \frac{19.1}{83.72} \right) \left(\frac{1}{6.2} \right)$$

$$x = \left(0.195 + \frac{19.1}{83.72} \right) \left(\frac{1}{6.2} \right) \approx 0.07$$

The solution set is approximately $\{0.07\}$.

Chapter 1: Equations and Inequalities

69. $14.72 - 21.58x = \frac{18}{2.11}x + 2.4$

$$14.72 - 21.58x - \frac{18}{2.11}x = \frac{18}{2.11}x + 2.4 - \frac{18}{2.11}x$$

$$14.72 - 21.58x - \frac{18}{2.11}x = 2.4$$

$$14.72 - 21.58x - \frac{18}{2.11}x - 14.72 = 2.4 - 14.72$$

$$-21.58x - \frac{18}{2.11}x = -12.32$$

$$\left(-21.58 - \frac{18}{2.11}\right)x = -12.32$$

$$\left(\frac{1}{-21.58 - \frac{18}{2.11}}\right) \left(-21.58 - \frac{18}{2.11}\right)x = -12.32 \left(\frac{1}{-21.58 - \frac{18}{2.11}}\right)$$

$$x = -12.32 \left(\frac{1}{-21.58 - \frac{18}{2.11}}\right) \approx 0.41$$

The solution set is approximately $\{0.41\}$.

70. $18.63x - \frac{21.2}{2.6} = \frac{14}{2.32}x - 20$

$$18.63x - \frac{21.2}{2.6} - \frac{14}{2.32}x = \frac{14}{2.32}x - 20 - \frac{14}{2.32}x$$

$$18.63x - \frac{21.2}{2.6} - \frac{14}{2.32}x = -20$$

$$18.63x - \frac{21.2}{2.6} - \frac{14}{2.32}x + \frac{21.2}{2.6} = -20 + \frac{21.2}{2.6}$$

$$18.63x - \frac{14}{2.32}x = -20 + \frac{21.2}{2.6}$$

$$\left(18.63 - \frac{14}{2.32}\right)x = -20 + \frac{21.2}{2.6}$$

$$\left(\frac{1}{18.63 - \frac{14}{2.32}}\right) \left(18.63 - \frac{14}{2.32}\right)x = \left(-20 + \frac{21.2}{2.6}\right) \left(\frac{1}{18.63 - \frac{14}{2.32}}\right)$$

$$x = \left(-20 + \frac{21.2}{2.6}\right) \left(\frac{1}{18.63 - \frac{14}{2.32}}\right) \approx -0.94$$

The solution set is approximately $\{-0.94\}$.

71. $ax - b = c, \quad a \neq 0$

$$ax - b + b = c + b$$

$$ax = b + c$$

$$\frac{ax}{a} = \frac{b+c}{a}$$

$$x = \frac{b+c}{a}$$

72. $1 - ax = b, \quad a \neq 0$

$$1 - ax - 1 = b - 1$$

$$-ax = b - 1$$

$$\frac{-ax}{-a} = \frac{b-1}{-a}$$

$$x = \frac{b-1}{-a} = \frac{1-b}{a}$$

73. $\frac{x}{a} + \frac{x}{b} = c, \quad a \neq 0, b \neq 0, a \neq -b$

$$ab\left(\frac{x}{a} + \frac{x}{b}\right) = ab \cdot c$$

$$bx + ax = abc$$

$$(a+b)x = abc$$

$$\frac{(a+b)x}{a+b} = \frac{abc}{a+b}$$

$$x = \frac{abc}{a+b}$$

74. $\frac{a}{x} + \frac{b}{x} = c, \quad c \neq 0$

$$x\left(\frac{a}{x} + \frac{b}{x}\right) = x \cdot c$$

$$a + b = cx$$

$$\frac{a+b}{c} = \frac{cx}{c}$$

$$x = \frac{a+b}{c}$$

75. $x + 2a = 16 + ax - 6a$, if $x = 4$

$$4 + 2a = 16 + a(4) - 6a$$

$$4 + 2a = 16 + 4a - 6a$$

$$4 + 2a = 16 - 2a$$

$$4a = 12$$

$$\frac{4a}{4} = \frac{12}{4}$$

$$a = 3$$

76. $x + 2b = x - 4 + 2bx$, for $x = 2$

$$2 + 2b = 2 - 4 + 2b(2)$$

$$2 + 2b = 2 - 4 + 4b$$

$$2 + 2b = -2 + 4b$$

$$4 = 2b$$

$$\frac{4}{2} = b$$

$$b = 2$$

77. $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2}$

$$RR_1R_2\left(\frac{1}{R}\right) = RR_1R_2\left(\frac{1}{R_1} + \frac{1}{R_2}\right)$$

$$R_1R_2 = RR_2 + RR_1$$

$$R_1R_2 = R(R_2 + R_1)$$

$$\frac{R_1R_2}{R_2 + R_1} = \frac{R(R_2 + R_1)}{R_2 + R_1}$$

$$\frac{R_1R_2}{R_2 + R_1} = R$$

78. $A = P(1 + rt)$

$$A = P + Prt$$

$$A - P = Prt$$

$$\frac{A - P}{Pt} = \frac{Prt}{Pt}$$

$$\frac{A - P}{Pt} = r$$

79. $F = \frac{mv^2}{R}$

$$RF = R\left(\frac{mv^2}{R}\right)$$

$$RF = mv^2$$

$$\frac{RF}{F} = \frac{mv^2}{F}$$

$$R = \frac{mv^2}{F}$$

80. $PV = nRT$

$$\frac{PV}{nR} = \frac{nRT}{nR}$$

$$\frac{PV}{nR} = T$$

Chapter 1: Equations and Inequalities

81. $S = \frac{a}{1-r}$

$$S(1-r) = \left(\frac{a}{1-r}\right)(1-r)$$

$$S - Sr = a$$

$$S - Sr - S = a - S$$

$$-Sr = a - S$$

$$\frac{-Sr}{-S} = \frac{a - S}{-S}$$

$$r = \frac{S - a}{S}$$

82. $v = -gt + v_0$

$$v - v_0 = -gt$$

$$\frac{v - v_0}{-g} = \frac{-gt}{-g}$$

$$t = \frac{v - v_0}{-g} = \frac{v_0 - v}{g}$$

83.

Amount in bonds	Amount in CDs	Total
x	$x - 3000$	20,000

$$x + (x - 3000) = 20,000$$

$$2x - 3000 = 20,000$$

$$2x = 23,000$$

$$x = 11,500$$

\$11,500 will be invested in bonds and \$8500 will be invested in CD's.

84.

Sean's Amount	George's Amount	Total
x	$x - 3000$	10,000

$$x + (x - 3000) = 10,000$$

$$2x - 3000 = 10,000$$

$$2x = 13,000$$

$$x = 6500$$

Sean will receive \$6500 and George will receive \$3500.

85.

	Dollars per hour	Hours worked	Money earned
Regular wage	x	40	$40x$
Overtime wage	$1.5x$	8	$8(1.5x)$

$$40x + 8(1.5x) = 546$$

$$40x + 12x = 546$$

$$52x = 546$$

$$x = \frac{546}{52} = 10.50$$

Sandra's regular hourly wage is \$10.50.

86.

	Dollars per hour	Hours worked	Money earned
Regular wage	x	40	$40x$
Overtime wage	$1.5x$	6	$6(1.5x)$
Sunday wage	$2x$	4	$4(2x)$

$$40x + 6(1.5x) + 4(2x) = 627$$

$$40x + 9x + 8x = 627$$

$$57x = 627$$

$$x = \frac{627}{57} = 11$$

Leigh's regular hourly wage is \$11.00.

87. Let x represent the score on the final exam.

$$\frac{80 + 83 + 71 + 61 + 95 + x + x}{7} = 80$$

$$\frac{390 + 2x}{7} = 80$$

$$390 + 2x = 560$$

$$2x = 170$$

$$x = 85$$

Brooke needs a score of 85 on the final exam.

88. Let x represent the score on the final exam.

Note: since the final exam counts for two-thirds of the overall grade, the average of the four test scores count for one-third of the overall grade. For a B, the average score must be 80.

$$\begin{aligned} \frac{1}{3}\left(\frac{86+80+84+90}{4}\right) + \frac{2}{3}x &= 80 \\ \frac{1}{3}\left(\frac{340}{4}\right) + \frac{2}{3}x &= 80 \\ \frac{85}{3} + \frac{2}{3}x &= 80 \\ 3\left(\frac{85}{3} + \frac{2}{3}x\right) &= 3(80) \\ 85 + 2x &= 240 \\ 2x &= 155 \\ x &= 77.5 \end{aligned}$$

Mike needs a score of 78 to earn a B.

For an A, the average score must be 90.

$$\begin{aligned} \frac{1}{3}\left(\frac{86+80+84+90}{4}\right) + \frac{2}{3}x &= 90 \\ \frac{1}{3}\left(\frac{340}{4}\right) + \frac{2}{3}x &= 90 \\ \frac{85}{3} + \frac{2}{3}x &= 90 \\ 3\left(\frac{85}{3} + \frac{2}{3}x\right) &= 3(90) \\ 85 + 2x &= 270 \\ 2x &= 185 \\ x &= 92.5 \end{aligned}$$

Mike needs a score of 93 to earn an A.

89. Let x represent the original price of the house.
 Then $0.15x$ represents the reduction in the price of the house.
 The new price of the home is \$425,000.
 original price – reduction = new price

$$\begin{aligned} x - 0.15x &= 425,000 \\ 0.85x &= 425,000 \\ x &= 500,000 \end{aligned}$$

The original price of the house was \$500,000.
 The amount of the reduction (i.e., the savings) is $0.15(\$500,000) = \$75,000$.

90. Let x represent the original price of the car.
 Then $0.15x$ represents the reduction in the price of the car.
 The new price of the car is \$8000.
 list price – reduction = new price

$$\begin{aligned} x - 0.15x &= 8000 \\ 0.85x &= 8000 \\ x &\approx 9411.76 \end{aligned}$$

The list price of the car was \$9411.76.
 The amount of the reduction (i.e., the savings) is $0.15(\$9411.76) \approx \1411.76 .

91. Let x represent the price the theater pays for the candy.
 Then $2.75x$ represents the markup on the candy.
 The selling price of the candy is \$3.00.
 supplier price + markup = selling price

$$x + 2.75x = 3.00$$

$$3.75x = 3.00$$

$$x = 0.80$$

The theater paid \$0.80 for the candy.

92. Let x represent selling price for the new car.
 The dealer's cost is $0.85(\$18,000) = \$15,300$.
 The markup is \$100.
 selling price = dealer's cost + markup
 $x = 15,300 + 100 = \$15,400$
 At \$100 over the dealer's cost, the price of the care is \$15,400.

	Tickets sold	Price per ticket	Money earned
Adults	x	7.50	7.50x
Children	$5200 - x$	4.50	4.50(5200 - x)

$$7.50x + 4.50(5200 - x) = 29,961$$

$$7.50x + 23,400 - 4.50x = 29,961$$

$$3.00x + 23,400 = 29,961$$

$$3.00x = 6561$$

$$x = 2187$$

There were 2187 adult patrons.

94. Let p represent the original price for the suit.
 Then, $0.30p$ represents the discounted amount.
 original price – discount = clearance price

$$p - 0.30p = 399$$

$$0.70p = 399$$

$$p = 570$$

The suit originally cost \$570.

95. Let w represent the width of the rectangle.
 Then $w + 8$ is the length.
 Perimeter is given by the formula $P = 2l + 2w$.

Chapter 1: Equations and Inequalities

$$2(w+8) + 2w = 60$$

$$2w+16+2w=60$$

$$4w+16=60$$

$$4w=44$$

$$w=11$$

Now, $11 + 8 = 19$.

The width of the rectangle is 11 feet and the length is 19 feet.

- 96.** Let w represent the width of the rectangle.
Then $2w$ is the length.
Perimeter is given by the formula $P = 2l + 2w$.

$$2(2w) + 2w = 42$$

$$4w + 2w = 42$$

$$6w = 42$$

$$w = 7$$

Now, $2(7) = 14$.

The width of the rectangle is 7 meters and the length is 14 meters.

- 97.** We will let B be the calories from breakfast, L the calories from lunch and D the calories from dinner. So we have the following equations:

$$B = L + 125$$

$$D = 2L - 300$$

$$2025 = B + L + D$$

Now we substitute the first two into the last one and solve for L .

$$2025 = (L + 125) + L + (2L - 300)$$

$$2025 = 4L - 175$$

$$2200 = 4L$$

$$L = 550$$

Now we substitute L into the first two equations to get B and D .

$$B = 550 + 125 = 675$$

$$D = 2(550) - 300 = 800$$

So Herschel took in 675 calories from breakfast, 550 calories from lunch and 800 calories from dinner.

- 98.** We will let B be the calories from breakfast, L the calories from lunch, D the calories from dinner and S the calories from snacks. So we have the following equations:

$$L = 0.5B \quad D = B + 200$$

$$S = B - 120 \quad E = 700$$

$$1480 = B + L + D - E$$

Now we substitute the first four into the last one and solve for B .

$$1480 = B + 0.5B + (B + 200) + (B - 200) - 700$$

$$1480 = 3.5B - 620$$

$$2100 = 3.5B$$

$$B = 600$$

Now we substitute B to get S .

$$S = B - 120 = 600 - 120 = 480$$

So Tyshira took in 480 calories from snacks.

	Judy's Amount	Tom's Amount	Total
	x	$\frac{2}{3}x$	18

$$x + \frac{2}{3}x = 18$$

$$\frac{5}{3}x = 18$$

$$x = \frac{3}{5}(18)$$

$$x = 10.80$$

Judy pays \$10.80 and Tom pays \$7.20.

- 100.** To move from step (6) to step (7), we divided both sides of the equation by the expression $x - 2$. From step (1), however, we know $x = 2$, so this means we divided both sides of the equation by zero.

101–102. Answers will vary.

Section 1.2

1. $x^2 - 5x - 6 = (x - 6)(x + 1)$

2. $2x^2 - x - 3 = (2x - 3)(x + 1)$

3. $\left\{-\frac{5}{3}, 3\right\}$

4. True

5. $\frac{1}{2} \cdot 5 = \frac{5}{2}; \left(\frac{5}{2}\right)^2 = \frac{25}{4}; x^2 + 5x + \frac{25}{4}$

$$x^2 + 5x + \frac{25}{4} = \left(x + \frac{5}{2}\right)^2$$

6. discriminant; negative

7. False; a quadratic equation may have no real solutions.

8. False; if the discriminant is positive, the equation has two distinct real solutions.

9. b

10. d

11. $x^2 - 9x = 0$

$$x(x - 9) = 0$$

$$x = 0 \text{ or } x - 9 = 0$$

$$x = 0 \text{ or } x = 9$$

The solution set is $\{0, 9\}$.

12. $x^2 + 4x = 0$

$$x(x + 4) = 0$$

$$x = 0 \text{ or } x + 4 = 0$$

$$x = 0 \text{ or } x = -4$$

The solution set is $\{-4, 0\}$.

13. $x^2 - 25 = 0$

$$(x + 5)(x - 5) = 0$$

$$x + 5 = 0 \text{ or } x - 5 = 0$$

$$x = -5 \text{ or } x = 5$$

The solution set is $\{-5, 5\}$.

14. $x^2 - 9 = 0$

$$(x + 3)(x - 3) = 0$$

$$x + 3 = 0 \text{ or } x - 3 = 0$$

$$x = -3 \text{ or } x = 3$$

The solution set is $\{-3, 3\}$.

15. $z^2 + z - 6 = 0$

$$(z + 3)(z - 2) = 0$$

$$z + 3 = 0 \text{ or } z - 2 = 0$$

$$z = -3 \text{ or } z = 2$$

The solution set is $\{-3, 2\}$.

16. $v^2 + 7v + 6 = 0$

$$(v + 6)(v + 1) = 0$$

$$v + 6 = 0 \text{ or } v + 1 = 0$$

$$v = -6 \text{ or } v = -1$$

The solution set is $\{-6, -1\}$

17. $2x^2 - 5x - 3 = 0$

$$(2x + 1)(x - 3) = 0$$

$$2x + 1 = 0 \text{ or } x - 3 = 0$$

$$x = -\frac{1}{2} \text{ or } x = 3$$

The solution set is $\left\{-\frac{1}{2}, 3\right\}$

18. $3x^2 + 5x + 2 = 0$

$$(3x + 2)(x + 1) = 0$$

$$3x + 2 = 0 \text{ or } x + 1 = 0$$

$$x = -\frac{2}{3} \text{ or } x = -1$$

The solution set is $\left\{-1, -\frac{2}{3}\right\}$.

19. $3t^2 - 48 = 0$

$$3(t^2 - 16) = 0$$

$$3(t + 4)(t - 4) = 0$$

$$t + 4 = 0 \text{ or } t - 4 = 0$$

$$t = -4 \text{ or } t = 4$$

The solution set is $\{-4, 4\}$.

20. $2y^2 - 50 = 0$

$$2(y^2 - 25) = 0$$

$$2(y + 5)(y - 5) = 0$$

$$y + 5 = 0 \text{ or } y - 5 = 0$$

$$y = -5 \text{ or } y = 5$$

The solution set is $\{-5, 5\}$.

21. $x(x - 8) + 12 = 0$

$$x^2 - 8x + 12 = 0$$

$$(x - 6)(x - 2) = 0$$

$$x - 6 = 0 \text{ or } x - 2 = 0$$

$$x = 6 \text{ or } x = 2$$

The solution set is $\{2, 6\}$.

Chapter 1: Equations and Inequalities

22. $x(x+4)=12$

$$x^2 + 4x - 12 = 0$$

$$(x+6)(x-2) = 0$$

$$x+6=0 \quad \text{or} \quad x-2=0$$

$$x=-6 \quad \text{or} \quad x=2$$

The solution set is $\{-6, 2\}$.

23. $4x^2 + 9 = 12x$

$$4x^2 - 12x + 9 = 0$$

$$(2x-3)^2 = 0$$

$$2x-3=0$$

$$x=\frac{3}{2}$$

The solution set is $\left\{\frac{3}{2}\right\}$.

24. $25x^2 + 16 = 40x$

$$25x^2 - 40x + 16 = 0$$

$$(5x-4)^2 = 0$$

$$5x-4=0$$

$$x=\frac{4}{5}$$

The solution set is $\left\{\frac{4}{5}\right\}$.

25. $6(p^2 - 1) = 5p$

$$6p^2 - 6 = 5p$$

$$6p^2 - 5p - 6 = 0$$

$$(3p+2)(2p-3) = 0$$

$$3p+2=0 \quad \text{or} \quad 2p-3=0$$

$$p=-\frac{2}{3} \quad \text{or} \quad p=\frac{3}{2}$$

The solution set is $\left\{-\frac{2}{3}, \frac{3}{2}\right\}$.

26. $2(2u^2 - 4u) + 3 = 0$

$$4u^2 - 8u + 3 = 0$$

$$(2u-1)(2u-3) = 0$$

$$2u-1=0 \quad \text{or} \quad 2u-3=0$$

$$u=\frac{1}{2} \quad \text{or} \quad u=\frac{3}{2}$$

The solution set is $\left\{\frac{1}{2}, \frac{3}{2}\right\}$.

27. $6x - 5 = \frac{6}{x}$

$$(6x-5)x = \left(\frac{6}{x}\right)x$$

$$6x^2 - 5x = 6$$

$$6x^2 - 5x - 6 = 0$$

$$(3x+2)(2x-3) = 0$$

$$3x+2=0 \quad \text{or} \quad 2x-3=0$$

$$x=-\frac{2}{3} \quad \text{or} \quad x=\frac{3}{2}$$

Neither of these values causes a denominator to equal zero, so the solution set is $\left\{-\frac{2}{3}, \frac{3}{2}\right\}$.

28. $x + \frac{12}{x} = 7$

$$\left(x + \frac{12}{x}\right)x = 7x$$

$$x^2 + 12 = 7x$$

$$x^2 - 7x + 12 = 0$$

$$(x-3)(x-4) = 0$$

$$x-3=0 \quad \text{or} \quad x-4=0$$

$$x=3 \quad \text{or} \quad x=4$$

Neither of these values causes a denominator to equal zero, so the solution set is $\{3, 4\}$.

29. $\frac{4(x-2)}{x-3} + \frac{3}{x} = \frac{-3}{x(x-3)}$

$$\left(\frac{4(x-2)}{x-3} + \frac{3}{x}\right)x(x-3) = \left(\frac{-3}{x(x-3)}\right)x(x-3)$$

$$4x(x-2) + 3(x-3) = -3$$

$$4x^2 - 8x + 3x - 9 = -3$$

$$4x^2 - 5x - 6 = 0$$

$$(4x+3)(x-2) = 0$$

$$4x+3=0 \quad \text{or} \quad x-2=0$$

$$x=-\frac{3}{4} \quad \text{or} \quad x=2$$

Neither of these values causes a denominator to equal zero, so the solution set is $\left\{-\frac{3}{4}, 2\right\}$.

30.

$$\frac{5}{x+4} = 4 + \frac{3}{x-2}$$

$$\left(\frac{5}{x+4}\right)(x+4)(x-2) = \left(4 + \frac{3}{x-2}\right)(x+4)(x-2)$$

$$5(x-2) = 4(x+4)(x-2) + 3(x+4)$$

$$5x-10 = 4(x^2 + 2x - 8) + 3x + 12$$

$$5x-10 = 4x^2 + 8x - 32 + 3x + 12$$

$$0 = 4x^2 + 6x - 10$$

$$0 = 2(2x^2 + 3x - 5)$$

$$0 = 2(2x+5)(x-1)$$

$$2x+5=0 \quad \text{or} \quad x-1=0$$

$$x=-\frac{5}{2} \quad \text{or} \quad x=1$$

Neither of these values causes a denominator to equal zero, so the solution set is $\left\{-\frac{5}{2}, 1\right\}$.

31. $x^2 = 25$

$$x = \pm\sqrt{25}$$

$$x = \pm 5$$

The solution set is $\{-5, 5\}$.

32. $x^2 = 36$

$$x = \pm\sqrt{36}$$

$$x = \pm 6$$

The solution set is $\{-6, 6\}$.

33. $(x-1)^2 = 4$

$$x-1 = \pm\sqrt{4}$$

$$x-1 = \pm 2$$

$$x-1 = 2 \quad \text{or} \quad x-1 = -2$$

$$x = 3 \quad \text{or} \quad x = -1$$

The solution set is $\{-1, 3\}$.

34. $(x+2)^2 = 1$

$$x+2 = \pm\sqrt{1}$$

$$x+2 = \pm 1$$

$$x+2 = 1 \quad \text{or} \quad x+2 = -1$$

$$x = -1 \quad \text{or} \quad x = -3$$

The solution set is $\{-3, -1\}$.

35. $(2x+3)^2 = 9$

$$2x+3 = \pm\sqrt{9}$$

$$2x+3 = \pm 3$$

$$2x+3 = 3 \quad \text{or} \quad 2x+3 = -3$$

$$2x = 0 \quad \text{or} \quad 2x = -6$$

$$x = 0 \quad \text{or} \quad x = -3$$

The solution set is $\{-3, 0\}$.

36. $(3x-2)^2 = 4$

$$3x-2 = \pm\sqrt{4}$$

$$3x-2 = \pm 2$$

$$3x-2 = 2 \quad \text{or} \quad 3x-2 = -2$$

$$3x = 4 \quad \text{or} \quad 3x = 0$$

$$x = \frac{4}{3} \quad \text{or} \quad x = 0$$

The solution set is $\left\{0, \frac{4}{3}\right\}$.

37. $x^2 + 4x = 21$

$$x^2 + 4x + 4 = 21 + 4$$

$$(x+2)^2 = 25$$

$$x+2 = \pm\sqrt{25}$$

$$x+2 = \pm 5$$

$$x = -2 \pm 5$$

$$x = 3 \quad \text{or} \quad x = -7$$

The solution set is $\{-7, 3\}$.

38. $x^2 - 6x = 13$

$$x^2 - 6x + 9 = 13 + 9$$

$$(x-3)^2 = 22$$

$$x-3 = \pm\sqrt{22}$$

$$x = 3 \pm \sqrt{22}$$

The solution set is $\{3 - \sqrt{22}, 3 + \sqrt{22}\}$.

Chapter 1: Equations and Inequalities

39. $x^2 - \frac{1}{2}x - \frac{3}{16} = 0$

$$x^2 - \frac{1}{2}x = \frac{3}{16}$$

$$x^2 - \frac{1}{2}x + \frac{1}{16} = \frac{3}{16} + \frac{1}{16}$$

$$\left(x - \frac{1}{4}\right)^2 = \frac{1}{4}$$

$$x - \frac{1}{4} = \pm\sqrt{\frac{1}{4}} = \pm\frac{1}{2}$$

$$x = \frac{1}{4} \pm \frac{1}{2}$$

$$x = \frac{3}{4} \text{ or } x = -\frac{1}{4}$$

The solution set is $\left\{-\frac{1}{4}, \frac{3}{4}\right\}$.

40. $x^2 + \frac{2}{3}x - \frac{1}{3} = 0$

$$x^2 + \frac{2}{3}x = \frac{1}{3}$$

$$x^2 + \frac{2}{3}x + \frac{1}{9} = \frac{1}{3} + \frac{1}{9}$$

$$\left(x + \frac{1}{3}\right)^2 = \frac{4}{9}$$

$$x + \frac{1}{3} = \pm\sqrt{\frac{4}{9}} = \pm\frac{2}{3}$$

$$x = -\frac{1}{3} \pm \frac{2}{3}$$

$$x = \frac{1}{3} \text{ or } x = -1$$

The solution set is $\left\{-1, \frac{1}{3}\right\}$.

41. $3x^2 + x - \frac{1}{2} = 0$

$$x^2 + \frac{1}{3}x - \frac{1}{6} = 0$$

$$x^2 + \frac{1}{3}x = \frac{1}{6}$$

$$x^2 + \frac{1}{3}x + \frac{1}{36} = \frac{1}{6} + \frac{1}{36}$$

$$\left(x + \frac{1}{6}\right)^2 = \frac{7}{36}$$

$$x + \frac{1}{6} = \pm\sqrt{\frac{7}{36}}$$

$$x + \frac{1}{6} = \pm\frac{\sqrt{7}}{6}$$

$$x = \frac{-1 \pm \sqrt{7}}{6}$$

The solution set is $\left\{\frac{-1-\sqrt{7}}{6}, \frac{-1+\sqrt{7}}{6}\right\}$.

42. $2x^2 - 3x - 1 = 0$

$$x^2 - \frac{3}{2}x - \frac{1}{2} = 0$$

$$x^2 - \frac{3}{2}x = \frac{1}{2}$$

$$x^2 - \frac{3}{2}x + \frac{9}{16} = \frac{1}{2} + \frac{9}{16}$$

$$\left(x - \frac{3}{4}\right)^2 = \frac{17}{16}$$

$$x - \frac{3}{4} = \pm\sqrt{\frac{17}{16}}$$

$$x - \frac{3}{4} = \pm\frac{\sqrt{17}}{4}$$

$$x = \frac{3 \pm \sqrt{17}}{4}$$

The solution set is $\left\{\frac{3-\sqrt{17}}{4}, \frac{3+\sqrt{17}}{4}\right\}$.

43. $x^2 - 4x + 2 = 0$

$$a = 1, \quad b = -4, \quad c = 2$$

$$x = \frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(2)}}{2(1)} = \frac{4 \pm \sqrt{16-8}}{2}$$

$$= \frac{4 \pm \sqrt{8}}{2} = \frac{4 \pm 2\sqrt{2}}{2} = 2 \pm \sqrt{2}$$

The solution set is $\{2 - \sqrt{2}, 2 + \sqrt{2}\}$.

44. $x^2 + 4x + 2 = 0$

$$a = 1, \quad b = 4, \quad c = 2$$

$$x = \frac{-4 \pm \sqrt{4^2 - 4(1)(2)}}{2(1)} = \frac{-4 \pm \sqrt{16 - 8}}{2} \\ = \frac{-4 \pm \sqrt{8}}{2} = \frac{-4 \pm 2\sqrt{2}}{2} = -2 \pm \sqrt{2}$$

The solution set is $\{-2 - \sqrt{2}, -2 + \sqrt{2}\}$.

45. $x^2 - 4x - 1 = 0$

$$a = 1, \quad b = -4, \quad c = -1$$

$$x = \frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(-1)}}{2(1)} = \frac{4 \pm \sqrt{16 + 4}}{2} \\ = \frac{4 \pm \sqrt{20}}{2} = \frac{4 \pm 2\sqrt{5}}{2} = 2 \pm \sqrt{5}$$

The solution set is $\{2 - \sqrt{5}, 2 + \sqrt{5}\}$.

46. $x^2 + 6x + 1 = 0$

$$a = 1, \quad b = 6, \quad c = 1$$

$$x = \frac{-6 \pm \sqrt{6^2 - 4(1)(1)}}{2(1)} = \frac{-6 \pm \sqrt{36 - 4}}{2} \\ = \frac{-6 \pm \sqrt{32}}{2} = \frac{-6 \pm 4\sqrt{2}}{2} = -3 \pm 2\sqrt{2}$$

The solution set is $\{-3 - 2\sqrt{2}, -3 + 2\sqrt{2}\}$.

47. $2x^2 - 5x + 3 = 0$

$$a = 2, \quad b = -5, \quad c = 3$$

$$x = \frac{-(-5) \pm \sqrt{(-5)^2 - 4(2)(3)}}{2(2)} \\ = \frac{5 \pm \sqrt{25 - 24}}{4} = \frac{5 \pm \sqrt{1}}{4} = \frac{5 \pm 1}{4}$$

$$x = \frac{5+1}{4} \text{ or } x = \frac{5-1}{4}$$

$$x = \frac{6}{4} \text{ or } x = \frac{4}{4}$$

$$x = \frac{3}{2} \text{ or } x = 1$$

The solution set is $\left\{1, \frac{3}{2}\right\}$.

48. $2x^2 + 5x + 3 = 0$

$$a = 2, \quad b = 5, \quad c = 3$$

$$x = \frac{-5 \pm \sqrt{5^2 - 4(2)(3)}}{2(2)} \\ = \frac{-5 \pm \sqrt{25 - 24}}{4} = \frac{-5 \pm \sqrt{1}}{4} = \frac{-5 \pm 1}{4}$$

$$x = \frac{-5+1}{4} \text{ or } x = \frac{-5-1}{4}$$

$$x = \frac{-4}{4} \text{ or } x = \frac{-6}{4}$$

$$x = -1 \text{ or } x = -\frac{3}{2}$$

The solution set is $\left\{-\frac{3}{2}, -1\right\}$.

49. $4y^2 - y + 2 = 0$

$$a = 4, \quad b = -1, \quad c = 2$$

$$y = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(4)(2)}}{2(4)} \\ = \frac{1 \pm \sqrt{1 - 32}}{8} = \frac{1 \pm \sqrt{-31}}{8}$$

No real solution.

50. $4t^2 + t + 1 = 0$

$$a = 4, \quad b = 1, \quad c = 1$$

$$t = \frac{-1 \pm \sqrt{1^2 - 4(4)(1)}}{2(4)} \\ = \frac{-1 \pm \sqrt{1 - 16}}{8} = \frac{-1 \pm \sqrt{-15}}{8}$$

No real solution.

51. $4x^2 = 1 - 2x$

$$4x^2 + 2x - 1 = 0$$

$$a = 4, \quad b = 2, \quad c = -1$$

$$x = \frac{-2 \pm \sqrt{2^2 - 4(4)(-1)}}{2(4)}$$

$$= \frac{-2 \pm \sqrt{4 + 16}}{8} = \frac{-2 \pm \sqrt{20}}{8}$$

$$= \frac{-2 \pm 2\sqrt{5}}{8} = \frac{-1 \pm \sqrt{5}}{4}$$

The solution set is $\left\{\frac{-1 - \sqrt{5}}{4}, \frac{-1 + \sqrt{5}}{4}\right\}$.

Chapter 1: Equations and Inequalities

52. $2x^2 = 1 - 2x$

$$2x^2 + 2x - 1 = 0$$

$$a = 2, \quad b = 2, \quad c = -1$$

$$x = \frac{-2 \pm \sqrt{2^2 - 4(2)(-1)}}{2(2)} = \frac{-2 \pm \sqrt{4+8}}{4} = \frac{-2 \pm \sqrt{12}}{4} = \frac{-2 \pm 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2}$$

The solution set is $\left\{ \frac{-1-\sqrt{3}}{2}, \frac{-1+\sqrt{3}}{2} \right\}$.

53. $4x^2 = 9x$

$$4x^2 - 9x = 0$$

$$x(4x - 9) = 0$$

$$x = 0 \quad \text{or} \quad 4x - 9 = 0$$

$$x = 0 \quad \text{or} \quad x = \frac{9}{4}$$

The solution set is $\left\{ 0, \frac{9}{4} \right\}$.

54. $5x = 4x^2$

$$0 = 4x^2 - 5x$$

$$0 = x(4x - 5)$$

$$x = 0 \quad \text{or} \quad 4x - 5 = 0$$

$$x = 0 \quad \text{or} \quad x = \frac{5}{4}$$

The solution set is $\left\{ 0, \frac{5}{4} \right\}$.

55. $9t^2 - 6t + 1 = 0$

$$a = 9, \quad b = -6, \quad c = 1$$

$$t = \frac{-(-6) \pm \sqrt{(-6)^2 - 4(9)(1)}}{2(9)} = \frac{6 \pm \sqrt{36-36}}{18} = \frac{6 \pm 0}{18} = \frac{1}{3}$$

The solution set is $\left\{ \frac{1}{3} \right\}$.

56. $4u^2 - 6u + 9 = 0$

$$a = 4, \quad b = -6, \quad c = 9$$

$$u = \frac{-(-6) \pm \sqrt{(-6)^2 - 4(4)(9)}}{2(4)} = \frac{6 \pm \sqrt{36-144}}{8} = \frac{6 \pm \sqrt{-108}}{8}$$

No real solution.

57. $\frac{3}{4}x^2 - \frac{1}{4}x - \frac{1}{2} = 0$

$$4\left(\frac{3}{4}x^2 - \frac{1}{4}x - \frac{1}{2}\right) = 4(0)$$

$$3x^2 - x - 2 = 0$$

$$a = 3, \quad b = -1, \quad c = -2$$

$$x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(3)(-2)}}{2(3)} = \frac{1 \pm \sqrt{1+24}}{6} = \frac{1 \pm \sqrt{25}}{6} = \frac{1 \pm 5}{6}$$

$$x = \frac{1+5}{6} \quad \text{or} \quad x = \frac{1-5}{6}$$

$$x = \frac{6}{6} \quad \text{or} \quad x = \frac{-4}{6}$$

$$x = 1 \quad \text{or} \quad x = -\frac{2}{3}$$

The solution set is $\left\{ -\frac{2}{3}, 1 \right\}$.

58. $\frac{2}{3}x^2 - x - 3 = 0$

$$3\left(\frac{2}{3}x^2 - x - 3\right) = 3(0)$$

$$2x^2 - 3x - 9 = 0$$

$$a = 2, \quad b = -3, \quad c = -9$$

$$x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(2)(-9)}}{2(2)} = \frac{3 \pm \sqrt{9+72}}{4} = \frac{3 \pm \sqrt{81}}{4} = \frac{3 \pm 9}{4}$$

$$x = \frac{3+9}{4} \quad \text{or} \quad x = \frac{3-9}{4}$$

$$x = \frac{12}{4} \quad \text{or} \quad x = \frac{-6}{4}$$

$$x = 3 \quad \text{or} \quad x = -\frac{3}{2}$$

The solution set is $\left\{ -\frac{3}{2}, 3 \right\}$.

59. $\frac{5}{3}x^2 - x = \frac{1}{3}$

$$3\left(\frac{5}{3}x^2 - x\right) = 3\left(\frac{1}{3}\right)$$

$$5x^2 - 3x = 1$$

$$5x^2 - 3x - 1 = 0$$

$$a = 5, \quad b = -3, \quad c = -1$$

$$x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(5)(-1)}}{2(5)}$$

$$= \frac{3 \pm \sqrt{9 + 20}}{10} = \frac{3 \pm \sqrt{29}}{10}$$

The solution set is $\left\{\frac{3 - \sqrt{29}}{10}, \frac{3 + \sqrt{29}}{10}\right\}$.

60. $\frac{3}{5}x^2 - x = \frac{1}{5}$

$$5\left(\frac{3}{5}x^2 - x\right) = 5\left(\frac{1}{5}\right)$$

$$3x^2 - 5x = 1$$

$$3x^2 - 5x - 1 = 0$$

$$a = 3, \quad b = -5, \quad c = -1$$

$$x = \frac{-(-5) \pm \sqrt{(-5)^2 - 4(3)(-1)}}{2(3)}$$

$$= \frac{5 \pm \sqrt{25 + 12}}{6} = \frac{5 \pm \sqrt{37}}{6}$$

The solution set is $\left\{\frac{5 - \sqrt{37}}{6}, \frac{5 + \sqrt{37}}{6}\right\}$.

61. $2x(x + 2) = 3$

$$2x^2 + 4x - 3 = 0$$

$$a = 2, \quad b = 4, \quad c = -3$$

$$x = \frac{-4 \pm \sqrt{4^2 - 4(2)(-3)}}{2(2)} = \frac{-4 \pm \sqrt{16 + 24}}{4}$$

$$= \frac{-4 \pm \sqrt{40}}{4} = \frac{-4 \pm 2\sqrt{10}}{4} = \frac{-2 \pm \sqrt{10}}{2}$$

The solution set is $\left\{\frac{-2 - \sqrt{10}}{2}, \frac{-2 + \sqrt{10}}{2}\right\}$.

62. $3x(x + 2) = 1$

$$3x^2 + 6x - 1 = 0$$

$$a = 3, \quad b = 6, \quad c = -1$$

$$x = \frac{-6 \pm \sqrt{6^2 - 4(3)(-1)}}{2(3)} = \frac{-6 \pm \sqrt{36 + 12}}{6}$$

$$= \frac{-6 \pm \sqrt{48}}{6} = \frac{-6 \pm 4\sqrt{3}}{6} = \frac{-3 \pm 2\sqrt{3}}{3}$$

The solution set is $\left\{\frac{-3 - 2\sqrt{3}}{3}, \frac{-3 + 2\sqrt{3}}{3}\right\}$.

63. $4 - \frac{1}{x} - \frac{2}{x^2} = 0$

$$x^2 \left(4 - \frac{1}{x} - \frac{2}{x^2}\right) = x^2(0)$$

$$4x^2 - x - 2 = 0$$

$$a = 4, \quad b = -1, \quad c = -2$$

$$x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(4)(-2)}}{2(4)}$$

$$= \frac{1 \pm \sqrt{1 + 32}}{8} = \frac{1 \pm \sqrt{33}}{8}$$

Neither of these values causes a denominator to equal zero, so the solution set is

$$\left\{\frac{1 - \sqrt{33}}{8}, \frac{1 + \sqrt{33}}{8}\right\}.$$

64. $4 + \frac{1}{x} - \frac{1}{x^2} = 0$

$$x^2 \left(4 + \frac{1}{x} - \frac{1}{x^2}\right) = x^2(0)$$

$$4x^2 + x - 1 = 0$$

$$a = 4, \quad b = 1, \quad c = -1$$

$$x = \frac{-1 \pm \sqrt{1^2 - 4(4)(-1)}}{2(4)}$$

$$= \frac{-1 \pm \sqrt{1 + 16}}{8} = \frac{-1 \pm \sqrt{17}}{8}$$

Neither of these values causes a denominator to equal zero, so the solution set is

$$\left\{\frac{-1 - \sqrt{17}}{8}, \frac{-1 + \sqrt{17}}{8}\right\}.$$

Chapter 1: Equations and Inequalities

65. $\frac{3x}{x-2} + \frac{1}{x} = 4$

$$\left(\frac{3x}{x-2} + \frac{1}{x}\right)x(x-2) = 4x(x-2)$$

$$3x(x) + (x-2) = 4x^2 - 8x$$

$$3x^2 + x - 2 = 4x^2 - 8x$$

$$0 = x^2 - 9x + 2$$

$$a = 1, \quad b = -9, \quad c = 2$$

$$x = \frac{-(-9) \pm \sqrt{(-9)^2 - 4(1)(2)}}{2(1)}$$

$$= \frac{9 \pm \sqrt{81-8}}{2} = \frac{9 \pm \sqrt{73}}{2}$$

Neither of these values causes a denominator to equal zero, so the solution set is

$$\left\{ \frac{9-\sqrt{73}}{2}, \frac{9+\sqrt{73}}{2} \right\}.$$

66. $\frac{2x}{x-3} + \frac{1}{x} = 4$

$$\left(\frac{2x}{x-3} + \frac{1}{x}\right)x(x-3) = 4x(x-3)$$

$$2x(x) + (x-3) = 4x^2 - 12x$$

$$2x^2 + x - 3 = 4x^2 - 12x$$

$$0 = 2x^2 - 13x + 3$$

$$a = 2, \quad b = -13, \quad c = 3$$

$$x = \frac{-(-13) \pm \sqrt{(-13)^2 - 4(2)(3)}}{2(2)}$$

$$= \frac{13 \pm \sqrt{169-24}}{4} = \frac{13 \pm \sqrt{145}}{4}$$

Neither of these values causes a denominator to equal zero, so the solution set is

$$\left\{ \frac{13-\sqrt{145}}{4}, \frac{13+\sqrt{145}}{4} \right\}.$$

67. $x^2 - 4.1x + 2.2 = 0$

$$a = 1, \quad b = -4.1, \quad c = 2.2$$

$$x = \frac{-(-4.1) \pm \sqrt{(-4.1)^2 - 4(1)(2.2)}}{2(1)}$$

$$= \frac{4.1 \pm \sqrt{16.81-8.8}}{2} = \frac{4.1 \pm \sqrt{8.01}}{2}$$

$$x \approx 3.47 \text{ or } x \approx 0.63$$

The solution set is $\{0.63, 3.47\}$.

68. $x^2 + 3.9x + 1.8 = 0$

$$a = 1, \quad b = 3.9, \quad c = 1.8$$

$$x = \frac{-3.9 \pm \sqrt{(3.9)^2 - 4(1)(1.8)}}{2(1)}$$

$$= \frac{-3.9 \pm \sqrt{15.21-7.2}}{2} = \frac{-3.9 \pm \sqrt{8.01}}{2}$$

$$x \approx -0.53 \text{ or } x \approx -3.37$$

The solution set is $\{-3.37, -0.53\}$.

69. $x^2 + \sqrt{3}x - 3 = 0$

$$a = 1, \quad b = \sqrt{3}, \quad c = -3$$

$$x = \frac{-\sqrt{3} \pm \sqrt{(\sqrt{3})^2 - 4(1)(-3)}}{2(1)}$$

$$= \frac{-\sqrt{3} \pm \sqrt{3+12}}{2} = \frac{-\sqrt{3} \pm \sqrt{15}}{2}$$

$$x \approx 1.07 \text{ or } x \approx -2.80$$

The solution set is $\{-2.80, 1.07\}$.

70. $x^2 + \sqrt{2}x - 2 = 0$

$$a = 1, \quad b = \sqrt{2}, \quad c = -2$$

$$x = \frac{-\sqrt{2} \pm \sqrt{(\sqrt{2})^2 - 4(1)(-2)}}{2(1)}$$

$$= \frac{-\sqrt{2} \pm \sqrt{2+8}}{2} = \frac{-\sqrt{2} \pm \sqrt{10}}{2}$$

$$x \approx 0.87 \text{ or } x \approx -2.29$$

The solution set is $\{-2.29, 0.87\}$.

71. $\pi x^2 - x - \pi = 0$

$$a = \pi, \quad b = -1, \quad c = -\pi$$

$$x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(\pi)(-\pi)}}{2(\pi)}$$

$$= \frac{1 \pm \sqrt{1+4\pi^2}}{2\pi}$$

$$x \approx 1.17 \text{ or } x \approx -0.85$$

The solution set is $\{-0.85, 1.17\}$.

72. $\pi x^2 + \pi x - 2 = 0$

$$a = \pi, \quad b = \pi, \quad c = -2$$

$$x = \frac{-\pi \pm \sqrt{(\pi)^2 - 4(\pi)(-2)}}{2(\pi)}$$

$$= \frac{-\pi \pm \sqrt{\pi^2 + 8\pi}}{2\pi}$$

$$x \approx 0.44 \text{ or } x \approx -1.44$$

The solution set is $\{-1.44, 0.44\}$.

73. $2x^2 - 6x + 7 = 0$

$$a = 2, \quad b = -6, \quad c = 7$$

$$b^2 - 4ac = (-6)^2 - 4(2)(7) = 36 - 56 = -20$$

Since the $b^2 - 4ac < 0$, the equation has no real solution.

74. $x^2 + 4x + 7 = 0$

$$a = 1, \quad b = 4, \quad c = 7$$

$$b^2 - 4ac = (4)^2 - 4(1)(7) = 16 - 28 = -12$$

Since the $b^2 - 4ac < 0$, the equation has no real solution.

75. $9x^2 - 30x + 25 = 0$

$$a = 9, \quad b = -30, \quad c = 25$$

$$b^2 - 4ac = (-30)^2 - 4(9)(25) = 900 - 900 = 0$$

Since $b^2 - 4ac = 0$, the equation has one repeated real solution.

76. $25x^2 - 20x + 4 = 0$

$$a = 25, \quad b = -20, \quad c = 4$$

$$b^2 - 4ac = (-20)^2 - 4(25)(4) = 400 - 400 = 0$$

Since $b^2 - 4ac = 0$, the equation has one repeated real solution.

77. $3x^2 + 5x - 8 = 0$

$$a = 3, \quad b = 5, \quad c = -8$$

$$b^2 - 4ac = (5)^2 - 4(3)(-8) = 25 + 96 = 121$$

Since $b^2 - 4ac > 0$, the equation has two unequal real solutions.

78. $2x^2 - 3x - 7 = 0$

$$a = 2, \quad b = -3, \quad c = -7$$

$$b^2 - 4ac = (-3)^2 - 4(2)(-7) = 9 + 56 = 65$$

Since $b^2 - 4ac > 0$, the equation has two unequal real solutions.

79. $x^2 - 5 = 0$

$$x^2 = 5$$

$$x = \pm\sqrt{5}$$

The solution set is $\{-\sqrt{5}, \sqrt{5}\}$.

80. $x^2 - 6 = 0$

$$x^2 = 6$$

$$x = \pm\sqrt{6}$$

The solution set is $\{-\sqrt{6}, \sqrt{6}\}$.

81. $16x^2 - 8x + 1 = 0$

$$(4x-1)(4x-1) = 0$$

$$4x-1 = 0$$

$$x = \frac{1}{4}$$

The solution set is $\left\{\frac{1}{4}\right\}$.

82. $9x^2 - 12x + 4 = 0$

$$(3x-2)(3x-2) = 0$$

$$3x-2 = 0$$

$$x = \frac{2}{3}$$

The solution set is $\left\{\frac{2}{3}\right\}$.

83. $10x^2 - 19x - 15 = 0$

$$(5x+3)(2x-5) = 0$$

$$5x+3 = 0 \quad \text{or} \quad 2x-5 = 0$$

$$x = -\frac{3}{5} \quad \text{or} \quad x = \frac{5}{2}$$

The solution set is $\left\{-\frac{3}{5}, \frac{5}{2}\right\}$.

84. $6x^2 + 7x - 20 = 0$

$$(3x-4)(2x+5) = 0$$

$$3x-4 = 0 \quad \text{or} \quad 2x+5 = 0$$

$$x = \frac{4}{3} \quad \text{or} \quad x = -\frac{5}{2}$$

The solution set is $\left\{-\frac{5}{2}, \frac{4}{3}\right\}$.

85. $2+z=6z^2$

$$0=6z^2-z-2$$

$$0=(3z-2)(2z+1)$$

$$3z-2=0 \text{ or } 2z+1=0$$

$$z=\frac{2}{3} \text{ or } z=-\frac{1}{2}$$

The solution set is $\left\{-\frac{1}{2}, \frac{2}{3}\right\}$.

86. $2=y+6y^2$

$$0=6y^2+y-2$$

$$0=(3y+2)(2y-1)$$

$$3y+2=0 \text{ or } 2y-1=0$$

$$y=-\frac{2}{3} \text{ or } y=\frac{1}{2}$$

The solution set is $\left\{-\frac{2}{3}, \frac{1}{2}\right\}$.

87. $x^2+\sqrt{2}x=\frac{1}{2}$

$$x^2+\sqrt{2}x-\frac{1}{2}=0$$

$$2\left(x^2+\sqrt{2}x-\frac{1}{2}\right)=2(0)$$

$$2x^2+2\sqrt{2}x-1=0$$

$$a=2, \quad b=2\sqrt{2}, \quad c=-1$$

$$x=\frac{-(2\sqrt{2})\pm\sqrt{(2\sqrt{2})^2-4(2)(-1)}}{2(2)}$$

$$=\frac{-2\sqrt{2}\pm\sqrt{8+8}}{4}=\frac{-2\sqrt{2}\pm\sqrt{16}}{4}$$

$$=\frac{-2\sqrt{2}\pm 4}{4}=\frac{-\sqrt{2}\pm 2}{2}$$

The solution set is $\left\{\frac{-\sqrt{2}-2}{2}, \frac{-\sqrt{2}+2}{2}\right\}$

88. $\frac{1}{2}x^2=\sqrt{2}x+1$

$$\frac{1}{2}x^2-\sqrt{2}x-1=0$$

$$2\left(\frac{1}{2}x^2-\sqrt{2}x-1\right)=2(0)$$

$$x^2-2\sqrt{2}x-2=0$$

$$a=1, \quad b=-2\sqrt{2}, \quad c=-2$$

$$x=\frac{-(-2\sqrt{2})\pm\sqrt{(-2\sqrt{2})^2-4(1)(-2)}}{2(1)}$$

$$=\frac{2\sqrt{2}\pm\sqrt{8+8}}{2}=\frac{2\sqrt{2}\pm\sqrt{16}}{2}$$

$$=\frac{2\sqrt{2}\pm 4}{2}=\frac{\sqrt{2}\pm 2}{1}$$

The solution set is $\left\{\sqrt{2}-2, \sqrt{2}+2\right\}$.

89. $x^2+x=4$

$$x^2+x-4=0$$

$$a=1, \quad b=1, \quad c=-4$$

$$x=\frac{-(1)\pm\sqrt{(1)^2-4(1)(-4)}}{2(1)}$$

$$=\frac{-1\pm\sqrt{1+16}}{2}=\frac{-1\pm\sqrt{17}}{2}$$

The solution set is $\left\{\frac{-1-\sqrt{17}}{2}, \frac{-1+\sqrt{17}}{2}\right\}$.

90. $x^2+x=1$

$$x^2+x-1=0$$

$$a=1, \quad b=1, \quad c=-1$$

$$x=\frac{-(1)\pm\sqrt{(1)^2-4(1)(-1)}}{2(1)}$$

$$=\frac{-1\pm\sqrt{1+4}}{2}=\frac{-1\pm\sqrt{5}}{2}$$

The solution set is $\left\{\frac{-1-\sqrt{5}}{2}, \frac{-1+\sqrt{5}}{2}\right\}$.

91.

$$\frac{x}{x-2} + \frac{2}{x+1} = \frac{7x+1}{x^2 - x - 2}$$

$$\frac{x}{x-2} + \frac{2}{x+1} = \frac{7x+1}{(x-2)(x+1)}$$

$$\left(\frac{x}{x-2} + \frac{2}{x+1}\right)(x-2)(x+1) = \left(\frac{7x+1}{(x-2)(x+1)}\right)(x-2)(x+1)$$

$$x(x+1) + 2(x-2) = 7x+1$$

$$x^2 + x + 2x - 4 = 7x+1$$

$$x^2 + 3x - 4 = 7x+1$$

$$x^2 - 4x - 5 = 0$$

$$(x+1)(x-5) = 0$$

$$x+1=0 \quad \text{or} \quad x-5=0$$

$$x=-1 \quad \text{or} \quad x=5$$

The value $x = -1$ causes a denominator to equal zero, so we disregard it. Thus, the solution set is $\{5\}$.

92.

$$\frac{3x}{x+2} + \frac{1}{x-1} = \frac{4-7x}{x^2 + x - 2}$$

$$\frac{3x}{x+2} + \frac{1}{x-1} = \frac{4-7x}{(x+2)(x-1)}$$

$$\left(\frac{3x}{x+2} + \frac{1}{x-1}\right)(x+2)(x-1) = \left(\frac{4-7x}{(x+2)(x-1)}\right)(x+2)(x-1)$$

$$3x(x-1) + (x+2) = 4-7x$$

$$3x^2 - 3x + x + 2 = 4-7x$$

$$3x^2 - 2x + 2 = 4-7x$$

$$3x^2 + 5x - 2 = 0$$

$$(3x-1)(x+2) = 0$$

$$3x-1=0 \quad \text{or} \quad x+2=0$$

$$x=\frac{1}{3} \quad \text{or} \quad x=-2$$

The value $x = -2$ causes a denominator to equal zero, so we disregard it. Thus, the solution set is $\left\{\frac{1}{3}\right\}$.

93. Since this is a right triangle then we can use the Pythagorean Theorem. So

$$(2x+3)^2 = (2x-5)^2 + (x+7)^2$$

$$4x^2 + 12x + 9 = 4x^2 - 20x + 25 + x^2 + 14x + 49$$

$$12x + 9 = x^2 - 6x + 74$$

$$0 = x^2 - 18x + 65$$

$$0 = (x-5)(x-13)$$

$$x-5=0 \quad \text{or} \quad x-13=0$$

$$x=5 \quad \text{or} \quad x=13$$

This means there are 2 possible that meet these requirements. Substituting x into the given sides gives:

When $x = 5$: 5m, 12m, 13m

When $x = 13$: 20m, 21m, 29m

Thus there are 2 solutions.

Chapter 1: Equations and Inequalities

94. Since this is a right triangle then we can use the Pythagorean Theorem. So

$$\begin{aligned}(4x+5)^2 &= (3x+13)^2 + x^2 \\ 16x^2 + 40x + 25 &= 9x^2 + 78x + 169 + x^2 \\ 6x^2 - 38x - 144 &= 0 \\ 2(3x^2 - 19x - 72) &= 0 \\ 2(3x+8)(x-9) &= 0 \\ 3x+8 = 0 \text{ or } x-9 &= 0 \\ x = -\frac{8}{3} \text{ or } x &= 9\end{aligned}$$

This means there are 2 possible solutions that meet these requirements. Substituting x into the given sides gives:

When $x = 9$: 41m, 40m, 9m

When $x = -\frac{8}{3}$ at least one side of the triangle has a negative measurement which is impossible. Thus there is only 1 triangle possible

95. Let w represent the width of window.

Then $l = w+2$ represents the length of the window.

Since the area is 143 square feet, we have:

$$w(w+2) = 143$$

$$w^2 + 2w - 143 = 0$$

$$(w+13)(w-11) = 0$$

$$\cancel{w=-13} \text{ or } w=11$$

Discard the negative solution since width cannot be negative. The width of the rectangular window is 11 feet and the length is 13 feet.

96. Let w represent the width of window.

Then $l = w+1$ represents the length of the window.

Since the area is 306 square centimeters, we have: $w(w+1) = 306$

$$w^2 + w - 306 = 0$$

$$(w+18)(w-17) = 0$$

$$\cancel{w=-18} \text{ or } w=17$$

Discard the negative solution since width cannot be negative. The width of the rectangular window is 17 centimeters and the length is 18 centimeters.

97. Let l represent the length of the rectangle.

Let w represent the width of the rectangle.

The perimeter is 26 meters and the area is 40

square meters.

$$2l + 2w = 26$$

$$l + w = 13 \text{ so } w = 13 - l$$

$$lw = 40$$

$$l(13-l) = 40$$

$$13l - l^2 = 40$$

$$l^2 - 13l + 40 = 0$$

$$(l-8)(l-5) = 0$$

$$l = 8 \text{ or } l = 5$$

$$w = 5 \text{ or } w = 8$$

The dimensions are 5 meters by 8 meters.

98. Let r represent the radius of the circle.

Since the field is a square with area 1250 square feet, the length of a side of the square is

$$\sqrt{1250} = 25\sqrt{2} \text{ feet.}$$

The length of the diagonal is $2r$.

Use the Pythagorean Theorem to solve for r :

$$(2r)^2 = (25\sqrt{2})^2 + (25\sqrt{2})^2$$

$$4r^2 = 1250 + 1250$$

$$4r^2 = 2500$$

$$r^2 = 625$$

$$r = 25$$

The shortest radius setting for the sprinkler is 25 feet.

99. Let x = length of side of original sheet in feet.

Length of box: $x-2$ feet

Width of box: $x-2$ feet

Height of box: 1 foot

$$V = l \cdot w \cdot h$$

$$4 = (x-2)(x-2)(1)$$

$$4 = x^2 - 4x + 4$$

$$0 = x^2 - 4x$$

$$0 = x(x-4)$$

$$x = 0 \text{ or } x = 4$$

Discard $x = 0$ since that is not a feasible length for the original sheet. Therefore, the original sheet should measure 4 feet on each side.

- 100.** Let x = width of original sheet in feet.

Length of sheet: $2x$

Length of box: $2x - 2$ feet

Width of box: $x - 2$ feet

Height of box: 1 foot

$$V = l \cdot w \cdot h$$

$$4 = (2x - 2)(x - 2)(1)$$

$$4 = 2x^2 - 6x + 4$$

$$0 = 2x^2 - 6x$$

$$0 = x^2 - 3x$$

$$0 = x(x - 3)$$

$$x = 0 \quad \text{or} \quad x = 3$$

Discard $x = 0$ since that is not a feasible length for the original sheet. Therefore, the original sheet is 3 feet wide and 6 feet long.

- 101. a.** When the ball strikes the ground, the distance from the ground will be 0.

Therefore, we solve

$$96 + 80t - 16t^2 = 0$$

$$-16t^2 + 80t + 96 = 0$$

$$t^2 - 5t - 6 = 0$$

$$(t - 6)(t + 1) = 0$$

$$t = 6 \quad \text{or} \quad t = -1$$

Discard the negative solution since the time of flight must be positive. The ball will strike the ground after 6 seconds.

- b.** When the ball passes the top of the building, it will be 96 feet from the ground. Therefore, we solve

$$96 + 80t - 16t^2 = 96$$

$$-16t^2 + 80t = 0$$

$$t^2 - 5t = 0$$

$$t(t - 5) = 0$$

$$t = 0 \quad \text{or} \quad t = 5$$

The ball is at the top of the building at time $t = 0$ when it is thrown. It will pass the top of the building on the way down after 5 seconds.

- 102. a.** To find when the object will be 15 meters above the ground, we solve

$$-4.9t^2 + 20t = 15$$

$$-4.9t^2 + 20t - 15 = 0$$

$$a = -4.9, b = 20, c = -15$$

$$t = \frac{-20 \pm \sqrt{20^2 - 4(-4.9)(-15)}}{2(-4.9)}$$

$$= \frac{-20 \pm \sqrt{106}}{-9.8} = \frac{20 \pm \sqrt{106}}{9.8}$$

$$t \approx 0.99 \quad \text{or} \quad t \approx 3.09$$

The object will be 15 meters above the ground after about 0.99 seconds (on the way up) and about 3.09 seconds (on the way down).

- b.** The object will strike the ground when the distance from the ground is 0. Therefore, we solve

$$-4.9t^2 + 20t = 0$$

$$t(-4.9t + 20) = 0$$

$$t = 0 \quad \text{or} \quad -4.9t + 20 = 0$$

$$-4.9t = -20$$

$$t \approx 4.08$$

The object will strike the ground after about 4.08 seconds.

- c.** $-4.9t^2 + 20t = 100$

$$-4.9t^2 + 20t - 100 = 0$$

$$a = -4.9, b = 20, c = -100$$

$$t = \frac{-20 \pm \sqrt{20^2 - 4(-4.9)(-100)}}{2(-4.9)}$$

$$= \frac{-20 \pm \sqrt{-1560}}{-9.8}$$

There is no real solution. The object never reaches a height of 100 meters.

- 103.** Let x represent the number of centimeters the length and width should be reduced.

$12 - x$ = the new length, $7 - x$ = the new width. The new volume is 90% of the old volume.

$$(12 - x)(7 - x)(3) = 0.9(12)(7)(3)$$

$$3x^2 - 57x + 252 = 226.8$$

$$3x^2 - 57x + 25.2 = 0$$

$$x^2 - 19x + 8.4 = 0$$

Chapter 1: Equations and Inequalities

$$x = \frac{-(-19) \pm \sqrt{(-19)^2 - 4(1)(8.4)}}{2(1)} = \frac{19 \pm \sqrt{327.4}}{2}$$

$$x \approx 0.45 \text{ or } x \approx 18.55$$

Since 18.55 exceeds the dimensions, it is discarded. The dimensions of the new chocolate bar are: 11.55 cm by 6.55 cm by 3 cm.

- 104.** Let x represent the number of centimeters the length and width should be reduced.
 $12-x$ = the new length, $7-x$ = the new width.
 The new volume is 80% of the old volume.
 $(12-x)(7-x)(3) = 0.8(12)(7)(3)$

$$3x^2 - 57x + 252 = 201.6$$

$$3x^2 - 57x + 50.4 = 0$$

$$x^2 - 19x + 16.8 = 0$$

$$x = \frac{-(-19) \pm \sqrt{(-19)^2 - 4(1)(16.8)}}{2(1)} = \frac{19 \pm \sqrt{293.8}}{2}$$

$$x \approx 0.93 \text{ or } x \approx 18.07$$

Since 18.07 exceeds the dimensions, it is discarded. The dimensions of the new chocolate bar are: 11.07 cm by 6.07 cm by 3 cm.

- 105.** Let x represent the width of the border measured in feet. The radius of the pool is 5 feet. Then $x+5$ represents the radius of the circle, including both the pool and the border. The total area of the pool and border is

$$A_T = \pi(x+5)^2.$$

The area of the pool is $A_P = \pi(5)^2 = 25\pi$.

The area of the border is

$$A_B = A_T - A_P = \pi(x+5)^2 - 25\pi.$$

Since the concrete is 3 inches or 0.25 feet thick, the volume of the concrete in the border is

$$0.25A_B = 0.25(\pi(x+5)^2 - 25\pi)$$

Solving the volume equation:

$$0.25(\pi(x+5)^2 - 25\pi) = 27$$

$$\pi(x^2 + 10x + 25 - 25) = 108$$

$$\pi x^2 + 10\pi x - 108 = 0$$

$$x = \frac{-10\pi \pm \sqrt{(10\pi)^2 - 4(\pi)(-108)}}{2(\pi)}$$

$$= \frac{-31.42 \pm \sqrt{100\pi^2 + 432\pi}}{6.28}$$

$$x \approx 2.71 \text{ or } x \approx -12.71$$

Discard the negative solution. The width of the border is roughly 2.71 feet.

- 106.** Let x represent the width of the border measured in feet. The radius of the pool is 5 feet. Then $x+5$ represents the radius of the circle, including both the pool and the border. The total area of the pool and border is

$$A_T = \pi(x+5)^2.$$

The area of the pool is $A_P = \pi(5)^2 = 25\pi$.

The area of the border is

$$A_B = A_T - A_P = \pi(x+5)^2 - 25\pi.$$

Since the concrete is 4 inches = $\frac{1}{3}$ foot thick, the volume of the concrete in the border is

$$\frac{1}{3}A_B = \frac{1}{3}(\pi(x+5)^2 - 25\pi)$$

Solving the volume equation:

$$\frac{1}{3}(\pi(x+5)^2 - 25\pi) = 27$$

$$\pi(x^2 + 10x + 25 - 25) = 81$$

$$\begin{aligned} \pi x^2 + 10\pi x - 81 &= 0 \\ x &= \frac{-10\pi \pm \sqrt{(10\pi)^2 - 4(\pi)(-81)}}{2(\pi)} \\ &= \frac{-31.42 \pm \sqrt{100\pi^2 + 324\pi}}{6.28} \end{aligned}$$

$$x \approx 2.13 \text{ or } x \approx -12.13$$

Discard the negative solution. The width of the border is approximately 2.13 feet.

- 107.** Let x represent the width of the border measured in feet.

The total area is $A_T = (6+2x)(10+2x)$.

The area of the garden is $A_G = 6 \cdot 10 = 60$.

The area of the border is

$$A_B = A_T - A_G = (6+2x)(10+2x) - 60.$$

Since the concrete is 3 inches or 0.25 feet thick, the volume of the concrete in the border is

$$0.25A_B = 0.25((6+2x)(10+2x) - 60)$$

Solving the volume equation:

$$0.25((6+2x)(10+2x) - 60) = 27$$

$$60 + 32x + 4x^2 - 60 = 108$$

$$4x^2 + 32x - 108 = 0$$

$$x^2 + 8x - 27 = 0$$

$$x = \frac{-8 \pm \sqrt{8^2 - 4(1)(-27)}}{2(1)} = \frac{-8 \pm \sqrt{172}}{2}$$

$$x \approx 2.56 \text{ or } x \approx -10.56$$

Discard the negative solution. The width of the border is approximately 2.56 feet.

- 108.** Let x = the width and $2x$ = the length of the patio. The height is $\frac{1}{3}$ foot and the concrete available is $8(27) = 216$ cubic feet..

$$V = lwh = x(2x) \cdot \frac{1}{3} = 216$$

$$\frac{2}{3}x^2 = 216$$

$$x^2 = 324$$

$$x = \pm 18$$

The dimensions of the patio are 18 feet by 36 feet.

- 109.** Let x = the length of a 9.7-inch iPad Air™ in a 4:3 format.

Then $\frac{3}{4}x$ = the width of the iPad. The diagonal of the 9.7-inch iPad is 9.7 inches, so by the Pythagorean theorem we have:

$$x^2 + \left(\frac{3}{4}x\right)^2 = 9.7^2$$

$$x^2 + \frac{9}{16}x^2 = 94.09$$

$$16x^2 + \frac{9}{16}x^2 = 16(94.09)$$

$$16x^2 + 9x^2 = 1505.44$$

$$25x^2 = 1505.44$$

$$x^2 = 60.2176$$

$$x = \pm\sqrt{60.2176} = \pm 7.76$$

Since the length cannot be negative, the length of the iPad is 7.76 inches and the width is

$$\frac{3}{4}(7.76) = 5.82 \text{ inches. Thus, the area of the}$$

iPad is $(7.76)(5.85) = 45.1632$ square inches.

Let y = the length of a 10-inch 16:10 format Google Nexus 10™.

Then $\frac{10}{16}y$ = the width of the Nexus.

The diagonal of a 10-inch Nexus is 10 inches, so by the Pythagorean theorem we have:

$$y^2 + \left(\frac{10}{16}y\right)^2 = 10^2$$

$$y^2 + \frac{100}{256}y^2 = 100$$

$$256y^2 + \frac{100}{256}y^2 = 256(100)$$

$$256y^2 + 100y^2 = 25600$$

$$356y^2 = 25600$$

$$y^2 = \frac{25600}{356}$$

$$y = \pm\sqrt{\frac{25600}{356}} \approx \pm 8.48$$

Since the length cannot be negative, the length of the Nexus is $\sqrt{\frac{25600}{356}} \approx 8.48$ inches and the width is $\frac{10}{16}\sqrt{\frac{25600}{356}} \approx 5.30$ inches. Thus, the area of the 10-inch 16:10 format Nexus is

$$\sqrt{\frac{25600}{356}} \quad \frac{10}{16}\sqrt{\frac{25600}{356}}$$

$$\approx 44.94 \text{ square inches.}$$

The iPad Air™ 4:3 format has the larger screen since its area is larger.

- 110.** Let x = the length of a 7.9-inch iPad Mini with Retina™ in a 4:3 format.

Then $\frac{3}{4}x$ = the width of the iPad. The diagonal of the 7.9-inch iPad is 7.9 inches, so by the Pythagorean theorem we have:

$$x^2 + \left(\frac{3}{4}x\right)^2 = 7.9^2$$

$$x^2 + \frac{9}{16}x^2 = 62.41$$

$$16x^2 + \frac{9}{16}x^2 = 16(62.41)$$

$$16x^2 + 9x^2 = 998.56$$

$$25x^2 = 998.56$$

$$x^2 = 39.9424$$

$$x = \pm\sqrt{39.9424} = \pm 6.32$$

Since the length cannot be negative, the length of the iPad is 6.32 inches and the width is

$$\frac{3}{4}(6.32) = 4.74 \text{ inches. Thus, the area of the}$$

iPad is $(6.32)(4.74) = 29.9568$ square inches.

Let y = the length of a 8-inch 16:9 format Dell Venue Pro™. Then $\frac{9}{16}y$ = the width of the

Dell. The diagonal of a 8-inch Dell is 8 inches, so by the Pythagorean theorem we have:

Chapter 1: Equations and Inequalities

$$\begin{aligned}
 y^2 + \frac{9}{16}y^2 &= 8^2 \\
 y^2 + \frac{81}{256}y^2 &= 64 \\
 256y^2 + \frac{81}{256}y^2 &= 256(64) \\
 256y^2 + 81y^2 &= 16384 \\
 337y^2 &= 16384 \\
 y^2 &= \frac{16384}{337} \\
 y &= \pm\sqrt{\frac{16384}{337}} \approx \pm 6.97
 \end{aligned}$$

Since the length cannot be negative, the length of the Nexus is $\sqrt{\frac{16384}{337}} \approx 6.97$ inches and the width is $\frac{9}{16}\sqrt{\frac{16384}{337}} \approx 3.922$ inches. Thus, the area of the 8-inch 16:9 format Dell is $(6.97)(3.922) \approx 27.34$ square inches.

The iPad Mini™ 4:3 format has the larger screen since its area is larger.

- 111.** Let x be 35. Then

$$\begin{aligned}
 h &= -0.00025x^2 + 0.04x \\
 &= -0.00025(35)^2 + 0.04(35) \\
 &\approx 1.1 \text{ ft}
 \end{aligned}$$

- 112.** Since d is expressed in 1000's we will set $d = 15$ and solve for x using the Quadratic Formula.

$$\begin{aligned}
 d &= -0.002x^2 + 0.319x + 7.512 \\
 15 &= -0.002x^2 + 0.319x + 7.512 \\
 0 &= -0.002x^2 + 0.319x - 7.488 \\
 x &= \frac{-0.319 \pm \sqrt{(0.319)^2 - 4(-0.002)(-7.448)}}{2(-0.002)} \\
 &= \frac{-0.319 \pm \sqrt{0.042177}}{-0.004} \\
 x &\approx 28.4 \text{ or } x \approx 131.1
 \end{aligned}$$

So the nearest year when the difference was \$15,000 occurred about 29 years after 1965 or 1994. The value 131.1 has no meaning since it is in the future.

- 113.** We will set $g = 2.97$ and solve for h using the Quadratic Formula.

$$\begin{aligned}
 g &= -0.0006x^2 + 0.015x + 3.04 \\
 2.97 &= -0.0006x^2 + 0.015x + 3.04 \\
 0 &= -0.0006x^2 + 0.015x + 0.07 \\
 x &= \frac{-0.015 \pm \sqrt{(0.015)^2 - 4(-0.0006)(0.07)}}{2(-0.0006)} \\
 &= \frac{-0.015 \pm \sqrt{0.000393}}{-0.0012} \\
 x &\approx 29 \text{ or } x \approx -4.02
 \end{aligned}$$

So the estimated numbers of hours worked by a student with a GPA of 2.97 is 29 hours. The value -4.02 has no meaning since it is negative.

- 114.** Let x be the numbers of members in the fraternity and s be the share paid by each member. Then $s = \frac{1470}{x}$. If there are 7 members who cannot contribute then the share goes up by \$5. So we have the following equation:

$$s + 5 = \frac{1470}{x-7} \text{ or } (s+5)(x-7) = 1470$$

Solving these two equations together:

$$\begin{aligned}
 (s+5)(x-7) &= 1470 \text{ and } s = \frac{1470}{x} \\
 \frac{1470}{x} + 5(x-7) &= 1470 \\
 1470 - \frac{10290}{x} + 5x - 35 &= 1470 \\
 5x - \frac{10290}{x} - 35 &= 0 \\
 5x^2 - 35x - 10290 &= 0 \\
 5x^2 - 35x - 10290 &= 0 \\
 x^2 - 7x - 2058 &= 0 \\
 (x+42)(x-49) &= 0 \\
 x &= -42 \text{ or } x = 49
 \end{aligned}$$

Since x is the number of members, it must be positive so the number of members is 49.

115. $\frac{1}{2}n(n+1) = 703$
 $n(n+1) = 1406$

$$\begin{aligned} n^2 + n - 1406 &= 0 \\ (n-37)(n+38) &= 0 \\ n = 37 \quad \text{or} \quad n = -38 \end{aligned}$$

Since the number of consecutive integers cannot be negative, we discard the negative value. We must add 37 consecutive integers, beginning at 1, in order to get a sum of 703.

116. $\frac{1}{2}n(n-3) = 65$
 $n(n-3) = 130$

$$\begin{aligned} n^2 - 3n - 130 &= 0 \\ (n-13)(n+10) &= 0 \\ n = 13 \quad \text{or} \quad n = -10 \end{aligned}$$

Since the number of sides cannot be negative, we discard the negative value. A polygon with 65 diagonals will have 13 sides.

$$\begin{aligned} \frac{1}{2}n(n-3) &= 80 \\ n(n-3) &= 160 \\ n^2 - 3n - 160 &= 0 \\ a = 1, b = -3, c = -160 \\ n = \frac{3 \pm \sqrt{(-3)^2 - 4(1)(-160)}}{2(1)} &= \frac{3 \pm \sqrt{646}}{2} \end{aligned}$$

Neither solution is an integer, so there is no polygon that has 80 diagonals.

117. The roots of a quadratic equation are
 $x_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$ and $x_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$

$$\begin{aligned} x_1 + x_2 &= \frac{-b - \sqrt{b^2 - 4ac}}{2a} + \frac{-b + \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-b - \sqrt{b^2 - 4ac} - b + \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-2b}{2a} \\ &= -\frac{b}{a} \end{aligned}$$

118. The roots of a quadratic equation are
 $x_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$ and $x_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$

$$\begin{aligned} x_1 \cdot x_2 &= \left(\frac{-b - \sqrt{b^2 - 4ac}}{2a} \right) \left(\frac{-b + \sqrt{b^2 - 4ac}}{2a} \right) \\ &= \frac{(-b)^2 - (\sqrt{b^2 - 4ac})^2}{(2a)^2} = \frac{b^2 - b^2 + 4ac}{4a^2} \\ &= \frac{4ac}{4a^2} \\ &= \frac{c}{a} \end{aligned}$$

119. In order to have one repeated solution, we need the discriminant to be 0.

$$\begin{aligned} b^2 - 4ac &= 0 \\ 1^2 - 4(k)(k) &= 0 \\ 1 - 4k^2 &= 0 \\ 4k^2 &= 1 \\ k^2 &= \frac{1}{4} \\ k &= \pm \sqrt{\frac{1}{4}} \\ k &= \frac{1}{2} \quad \text{or} \quad k = -\frac{1}{2} \end{aligned}$$

120. In order to have one repeated solution, we need the discriminant to be 0.

$$\begin{aligned} b^2 - 4ac &= 0 \\ (-k)^2 - 4(1)(4) &= 0 \\ k^2 - 16 &= 0 \\ (k-4)(k+4) &= 0 \\ k = 4 \quad \text{or} \quad k = -4 \end{aligned}$$

121. For $ax^2 + bx + c = 0$:

$$x_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a} \quad \text{and} \quad x_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$

For $ax^2 - bx + c = 0$:

$$\begin{aligned}x_1^* &= \frac{-(-b) - \sqrt{(-b)^2 - 4ac}}{2a} \\&= \frac{b - \sqrt{b^2 - 4ac}}{2a} \\&= -\left(\frac{-b + \sqrt{b^2 - 4ac}}{2a}\right) \\&= -x_2\end{aligned}$$

and

$$\begin{aligned}x_2^* &= \frac{-(-b) + \sqrt{(-b)^2 - 4ac}}{2a} \\&= \frac{b + \sqrt{b^2 - 4ac}}{2a} \\&= -\left(\frac{-b - \sqrt{b^2 - 4ac}}{2a}\right) \\&= -x_1\end{aligned}$$

122. For $ax^2 + bx + c = 0$:

$$x_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a} \text{ and } x_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$

For $cx^2 + bx + a = 0$:

$$\begin{aligned}x_1^* &= \frac{-b - \sqrt{b^2 - 4(c)(a)}}{2c} = \frac{-b - \sqrt{b^2 - 4ac}}{2c} \\&= \frac{-b - \sqrt{b^2 - 4ac}}{2c} \cdot \frac{-b + \sqrt{b^2 - 4ac}}{-b + \sqrt{b^2 - 4ac}} \\&= \frac{b^2 - (b^2 - 4ac)}{2c(-b + \sqrt{b^2 - 4ac})} = \frac{4ac}{2c(-b + \sqrt{b^2 - 4ac})} \\&= \frac{2a}{-b + \sqrt{b^2 - 4ac}} \\&= \frac{1}{x_2}\end{aligned}$$

and

$$\begin{aligned}x_2^* &= \frac{-b + \sqrt{b^2 - 4(c)(a)}}{2c} = \frac{-b + \sqrt{b^2 - 4ac}}{2c} \\&= \frac{-b + \sqrt{b^2 - 4ac}}{2c} \cdot \frac{-b - \sqrt{b^2 - 4ac}}{-b - \sqrt{b^2 - 4ac}} \\&= \frac{b^2 - (b^2 - 4ac)}{2c(-b - \sqrt{b^2 - 4ac})} = \frac{4ac}{2c(-b - \sqrt{b^2 - 4ac})} \\&= \frac{2a}{-b - \sqrt{b^2 - 4ac}} \\&= \frac{1}{x_1}\end{aligned}$$

123. a. $x^2 = 9$ and $x = 3$ are not equivalent because they do not have the same solution set. In the first equation we can also have $x = -3$.

b. $x = \sqrt{9}$ and $x = 3$ are equivalent because $\sqrt{9} = 3$.

c. $(x-1)(x-2) = (x-1)^2$ and $x-2 = x-1$ are not equivalent because they do not have the same solution set.

The first equation has the solution set $\{1\}$ while the second equation has no solutions.

124. Answers will vary. Methods may include the quadratic formula, completing the square, graphing, etc.

125. Answers will vary. Knowing the discriminant allows us to know how many real solutions the equation will have.

126. Answers will vary. One possibility:
Two distinct: $x^2 - 3x - 18 = 0$
One repeated: $x^2 - 14x + 49 = 0$
No real: $x^2 + x + 4 = 0$

127. Answers will vary.

Section 1.3

1. Integers: $\{-3, 0\}$

Rationals: $\left\{-3, 0, \frac{6}{5}\right\}$

Section 1.3: Complex Numbers; Quadratic Equations in the Complex Number System

2. True; the set of real numbers consists of all rational and irrational numbers.

$$\begin{aligned} 3. \quad \frac{3}{2+\sqrt{3}} &= \frac{3}{2+\sqrt{3}} \cdot \frac{2-\sqrt{3}}{2-\sqrt{3}} \\ &= \frac{3(2-\sqrt{3})}{2^2 - (\sqrt{3})^2} \\ &= \frac{3(2-\sqrt{3})}{4-3} \\ &= 3(2-\sqrt{3}) \end{aligned}$$

4. real; imaginary; imaginary unit

5. False; the conjugate of $2+5i$ is $2-5i$.

6. True; the set of real numbers is a subset of the set of complex numbers.

7. False; if $2-3i$ is a solution of a quadratic equation with real coefficients, then its conjugate, $2+3i$, is also a solution.

8. b

9. a

10. c

$$11. \quad (2-3i)+(6+8i) = (2+6)+(-3+8)i = 8+5i$$

$$12. \quad (4+5i)+(-8+2i) = (4+(-8))+(5+2)i \\ = -4+7i$$

$$13. \quad (-3+2i)-(4-4i) = (-3-4)+(2-(-4))i \\ = -7+6i$$

$$14. \quad (3-4i)-(-3-4i) = (3-(-3)) + (-4-(-4))i \\ = 6+0i = 6$$

$$15. \quad (2-5i)-(8+6i) = (2-8)+(-5-6)i \\ = -6-11i$$

$$16. \quad (-8+4i)-(2-2i) = (-8-2)+(4-(-2))i \\ = -10+6i$$

$$17. \quad 3(2-6i) = 6-18i$$

$$18. \quad -4(2+8i) = -8-32i$$

$$19. \quad 2i(2-3i) = 4i-6i^2 = 4i-6(-1) = 6+4i$$

$$20. \quad 3i(-3+4i) = -9i+12i^2 = -9i+12(-1) = -12-9i$$

$$21. \quad (3-4i)(2+i) = 6+3i-8i-4i^2 \\ = 6-5i-4(-1) \\ = 10-5i$$

$$22. \quad (5+3i)(2-i) = 10-5i+6i-3i^2 \\ = 10+i-3(-1) \\ = 13+i$$

$$23. \quad (-6+i)(-6-i) = 36+6i-6i-i^2 \\ = 36-(-1) \\ = 37$$

$$24. \quad (-3+i)(3+i) = -9-3i+3i+i^2 \\ = -9+(-1) \\ = -10$$

$$25. \quad \frac{10}{3-4i} = \frac{10}{3-4i} \cdot \frac{3+4i}{3+4i} = \frac{30+40i}{9+12i-12i-16i^2} \\ = \frac{30+40i}{9-16(-1)} = \frac{30+40i}{25} \\ = \frac{30}{25} + \frac{40}{25}i \\ = \frac{6}{5} + \frac{8}{5}i$$

$$26. \quad \frac{13}{5-12i} = \frac{13}{5-12i} \cdot \frac{5+12i}{5+12i} \\ = \frac{65+156i}{25+60i-60i-144i^2} \\ = \frac{65+156i}{25-144(-1)} = \frac{65+156i}{169} \\ = \frac{65}{169} + \frac{156}{169}i \\ = \frac{5}{13} + \frac{12}{13}i$$

$$27. \quad \frac{2+i}{i} = \frac{2+i}{i} \cdot \frac{-i}{-i} = \frac{-2i-i^2}{-i^2} \\ = \frac{-2i-(-1)}{-(-1)} = \frac{1-2i}{1} = 1-2i$$

Chapter 1: Equations and Inequalities

$$28. \frac{2-i}{-2i} = \frac{2-i}{-2i} \cdot \frac{i}{i} = \frac{2i-i^2}{-2i^2}$$

$$= \frac{2i-(-1)}{-2(-1)} = \frac{1+2i}{2} = \frac{1}{2} + i$$

$$29. \frac{6-i}{1+i} = \frac{6-i}{1+i} \cdot \frac{1-i}{1-i} = \frac{6-6i-i+i^2}{1-i+i-i^2}$$

$$= \frac{6-7i+(-1)}{1-(-1)} = \frac{5-7i}{2} = \frac{5}{2} - \frac{7}{2}i$$

$$30. \frac{2+3i}{1-i} = \frac{2+3i}{1-i} \cdot \frac{1+i}{1+i} = \frac{2+2i+3i+3i^2}{1+i-i-i^2}$$

$$= \frac{2+5i+3(-1)}{1-(-1)} = \frac{-1+5i}{2} = -\frac{1}{2} + \frac{5}{2}i$$

$$31. \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)^2 = \frac{1}{4} + 2\left(\frac{1}{2}\right)\left(\frac{\sqrt{3}}{2}i\right) + \frac{3}{4}i^2$$

$$= \frac{1}{4} + \frac{\sqrt{3}}{2}i + \frac{3}{4}(-1) = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$$

$$32. \left(\frac{\sqrt{3}}{2} - \frac{1}{2}i\right)^2 = \frac{3}{4} - 2\left(\frac{\sqrt{3}}{2}\right)\left(\frac{1}{2}i\right) + \frac{1}{4}i^2$$

$$= \frac{3}{4} - \frac{\sqrt{3}}{2}i + \frac{1}{4}(-1) = \frac{1}{2} - \frac{\sqrt{3}}{2}i$$

$$33. (1+i)^2 = 1+2i+i^2 = 1+2i+(-1) = 2i$$

$$34. (1-i)^2 = 1-2i+i^2 = 1-2i+(-1) = -2i$$

$$35. i^{23} = i^{22+1} = i^{22} \cdot i = (i^2)^{11} \cdot i = (-1)^{11}i = -i$$

$$36. i^{14} = (i^2)^7 = (-1)^7 = -1$$

$$37. i^{-15} = \frac{1}{i^{15}} = \frac{1}{i^{14+1}} = \frac{1}{i^{14} \cdot i} = \frac{1}{(i^2)^7 \cdot i}$$

$$= \frac{1}{(-1)^7 i} = \frac{1}{-i} = \frac{1}{-i} \cdot \frac{i}{i} = \frac{i}{-i^2} = \frac{i}{-(1)} = i$$

$$38. i^{-23} = \frac{1}{i^{23}} = \frac{1}{i^{22+1}} = \frac{1}{i^{22} \cdot i} = \frac{1}{(i^2)^{11} \cdot i}$$

$$= \frac{1}{(-1)^{11} i} = \frac{1}{-i} = \frac{1}{-i} \cdot \frac{i}{i} = \frac{i}{-i^2} = \frac{i}{-(1)} = i$$

$$39. i^6 - 5 = (i^2)^3 - 5 = (-1)^3 - 5 = -1 - 5 = -6$$

$$40. 4+i^3 = 4+i^2 \cdot i = 4+(-1)i = 4-i$$

$$41. 6i^3 - 4i^5 = i^3(6-4i^2)$$

$$= i^2 \cdot i(6-4(-1)) = -1 \cdot i(10) = -10i$$

$$42. 4i^3 - 2i^2 + 1 = 4i^2 \cdot i - 2i^2 + 1$$

$$= 4(-1)i - 2(-1) + 1$$

$$= -4i + 2 + 1$$

$$= 3 - 4i$$

$$43. (1+i)^3 = (1+i)(1+i)(1+i) = (1+2i+i^2)(1+i)$$

$$= (1+2i-1)(1+i) = 2i(1+i)$$

$$= 2i+2i^2 = 2i+2(-1)$$

$$= -2+2i$$

$$44. (3i)^4 + 1 = 81i^4 + 1 = 81(1) + 1 = 82$$

$$45. i^7(1+i^2) = i^7(1+(-1)) = i^7(0) = 0$$

$$46. 2i^4(1+i^2) = 2(1)(1+(-1)) = 2(0) = 0$$

$$47. i^6 + i^4 + i^2 + 1 = (i^2)^3 + (i^2)^2 + i^2 + 1$$

$$= (-1)^3 + (-1)^2 + (-1) + 1$$

$$= -1 + 1 - 1 + 1$$

$$= 0$$

$$48. i^7 + i^5 + i^3 + i = (i^2)^3 \cdot i + (i^2)^2 \cdot i + i^2 \cdot i + i$$

$$= (-1)^3 \cdot i + (-1)^2 \cdot i + (-1) \cdot i + i$$

$$= -i + i - i + i$$

$$= 0$$

$$49. \sqrt{-4} = 2i$$

$$50. \sqrt{-9} = 3i$$

$$51. \sqrt{-25} = 5i$$

$$52. \sqrt{-64} = 8i$$

53. $\sqrt{(3+4i)(4i-3)} = \sqrt{12i-9+16i^2-12i}$
 $= \sqrt{-9+16(-1)}$
 $= \sqrt{-25}$
 $= 5i$

54. $\sqrt{(4+3i)(3i-4)} = \sqrt{12i-16+9i^2-12i}$
 $= \sqrt{-16+9(-1)}$
 $= \sqrt{-25}$
 $= 5i$

55. $x^2 + 4 = 0$
 $x^2 = -4$
 $x = \pm\sqrt{-4}$
 $x = \pm 2i$

The solution set is $\{-2i, 2i\}$.

56. $x^2 - 4 = 0$
 $(x+2)(x-2) = 0$
 $x = -2$ or $x = 2$
The solution set is $\{-2, 2\}$.

57. $x^2 - 16 = 0$
 $(x+4)(x-4) = 0$
 $x = -4$ or $x = 4$
The solution set is $\{-4, 4\}$.

58. $x^2 + 25 = 0$
 $x^2 = -25$
 $x = \pm\sqrt{-25} = \pm 5i$
The solution set is $\{-5i, 5i\}$.

59. $x^2 - 6x + 13 = 0$
 $a = 1, b = -6, c = 13,$
 $b^2 - 4ac = (-6)^2 - 4(1)(13) = 36 - 52 = -16$
 $x = \frac{-(-6) \pm \sqrt{-16}}{2(1)} = \frac{6 \pm 4i}{2} = 3 \pm 2i$

The solution set is $\{3-2i, 3+2i\}$.

60. $x^2 + 4x + 8 = 0$
 $a = 1, b = 4, c = 8$
 $b^2 - 4ac = 4^2 - 4(1)(8) = 16 - 32 = -16$
 $x = \frac{-4 \pm \sqrt{-16}}{2(1)} = \frac{-4 \pm 4i}{2} = -2 \pm 2i$

The solution set is $\{-2-2i, -2+2i\}$.

61. $x^2 - 6x + 10 = 0$
 $a = 1, b = -6, c = 10$
 $b^2 - 4ac = (-6)^2 - 4(1)(10) = 36 - 40 = -4$
 $x = \frac{-(-6) \pm \sqrt{-4}}{2(1)} = \frac{6 \pm 2i}{2} = 3 \pm i$

The solution set is $\{3-i, 3+i\}$.

62. $x^2 - 2x + 5 = 0$
 $a = 1, b = -2, c = 5$
 $b^2 - 4ac = (-2)^2 - 4(1)(5) = 4 - 20 = -16$
 $x = \frac{-(-2) \pm \sqrt{-16}}{2(1)} = \frac{2 \pm 4i}{2} = 1 \pm 2i$

The solution set is $\{1-2i, 1+2i\}$.

63. $8x^2 - 4x + 1 = 0$
 $a = 8, b = -4, c = 1$
 $b^2 - 4ac = (-4)^2 - 4(8)(1) = 16 - 32 = -16$
 $x = \frac{-(-4) \pm \sqrt{-16}}{2(8)} = \frac{4 \pm 4i}{16} = \frac{1}{4} \pm \frac{1}{4}i$

The solution set is $\left\{ \frac{1}{4} - \frac{1}{4}i, \frac{1}{4} + \frac{1}{4}i \right\}$.

64. $10x^2 + 6x + 1 = 0$
 $a = 10, b = 6, c = 1$
 $b^2 - 4ac = 6^2 - 4(10)(1) = 36 - 40 = -4$
 $x = \frac{-6 \pm \sqrt{-4}}{2(10)} = \frac{-6 \pm 2i}{20} = -\frac{3}{10} \pm \frac{1}{10}i$

The solution set is $\left\{ -\frac{3}{10} - \frac{1}{10}i, -\frac{3}{10} + \frac{1}{10}i \right\}$.

Chapter 1: Equations and Inequalities

65. $5x^2 + 1 = 2x$

$$5x^2 - 2x + 1 = 0$$

$$a = 5, b = -2, c = 1$$

$$b^2 - 4ac = (-2)^2 - 4(5)(1) = 4 - 20 = -16$$

$$x = \frac{-(2) \pm \sqrt{-16}}{2(5)} = \frac{2 \pm 4i}{10} = \frac{1}{5} \pm \frac{2}{5}i$$

The solution set is $\left\{\frac{1}{5} - \frac{2}{5}i, \frac{1}{5} + \frac{2}{5}i\right\}$.

66. $13x^2 + 1 = 6x$

$$13x^2 - 6x + 1 = 0$$

$$a = 13, b = -6, c = 1$$

$$b^2 - 4ac = (-6)^2 - 4(13)(1) = 36 - 52 = -16$$

$$x = \frac{-(-6) \pm \sqrt{-16}}{2(13)} = \frac{6 \pm 4i}{26} = \frac{3}{13} \pm \frac{2}{13}i$$

The solution set is $\left\{\frac{3}{13} - \frac{2}{13}i, \frac{3}{13} + \frac{2}{13}i\right\}$.

67. $x^2 + x + 1 = 0$

$$a = 1, b = 1, c = 1$$

$$b^2 - 4ac = 1^2 - 4(1)(1) = 1 - 4 = -3$$

$$x = \frac{-1 \pm \sqrt{-3}}{2(1)} = \frac{-1 \pm \sqrt{3}i}{2} = -\frac{1}{2} \pm \frac{\sqrt{3}}{2}i$$

The solution set is $\left\{-\frac{1}{2} - \frac{\sqrt{3}}{2}i, -\frac{1}{2} + \frac{\sqrt{3}}{2}i\right\}$.

68. $x^2 - x + 1 = 0$

$$a = 1, b = -1, c = 1$$

$$b^2 - 4ac = (-1)^2 - 4(1)(1) = 1 - 4 = -3$$

$$x = \frac{-(-1) \pm \sqrt{-3}}{2(1)} = \frac{1 \pm \sqrt{3}i}{2} = \frac{1}{2} \pm \frac{\sqrt{3}}{2}i$$

The solution set is $\left\{\frac{1}{2} - \frac{\sqrt{3}}{2}i, \frac{1}{2} + \frac{\sqrt{3}}{2}i\right\}$.

69. $x^3 - 8 = 0$

$$(x-2)(x^2 + 2x + 4) = 0$$

$$x - 2 = 0 \Rightarrow x = 2$$

$$\text{or } x^2 + 2x + 4 = 0$$

$$a = 1, b = 2, c = 4$$

$$b^2 - 4ac = 2^2 - 4(1)(4) = 4 - 16 = -12$$

$$x = \frac{-2 \pm \sqrt{-12}}{2(1)} = \frac{-2 \pm 2\sqrt{3}i}{2} = -1 \pm \sqrt{3}i$$

The solution set is $\{2, -1 - \sqrt{3}i, -1 + \sqrt{3}i\}$.

70. $x^3 + 27 = 0$

$$(x+3)(x^2 - 3x + 9) = 0$$

$$x + 3 = 0 \Rightarrow x = -3$$

$$\text{or } x^2 - 3x + 9 = 0$$

$$a = 1, b = -3, c = 9$$

$$b^2 - 4ac = (-3)^2 - 4(1)(9) = 9 - 36 = -27$$

$$x = \frac{-(-3) \pm \sqrt{-27}}{2(1)} = \frac{3 \pm 3\sqrt{3}i}{2} = \frac{3}{2} \pm \frac{3\sqrt{3}}{2}i$$

The solution set is $\left\{-3, \frac{3}{2} - \frac{3\sqrt{3}}{2}i, \frac{3}{2} + \frac{3\sqrt{3}}{2}i\right\}$.

71. $x^4 = 16$

$$x^4 - 16 = 0$$

$$(x^2 - 4)(x^2 + 4) = 0$$

$$(x-2)(x+2)(x^2 + 4) = 0$$

$$x - 2 = 0 \text{ or } x + 2 = 0 \text{ or } x^2 + 4 = 0$$

$$x = 2 \text{ or } x = -2 \text{ or } x^2 = -4$$

$$x = 2 \text{ or } x = -2 \text{ or } x = \pm\sqrt{-4} = \pm 2i$$

The solution set is $\{-2, 2, -2i, 2i\}$.

72. $x^4 = 1$

$$x^4 - 1 = 0$$

$$(x^2 - 1)(x^2 + 1) = 0$$

$$(x-1)(x+1)(x^2 + 1) = 0$$

$$x - 1 = 0 \text{ or } x + 1 = 0 \text{ or } x^2 + 1 = 0$$

$$x = 1 \text{ or } x = -1 \text{ or } x^2 = -1$$

$$x = 1 \text{ or } x = -1 \text{ or } x = \pm\sqrt{-1} = \pm i$$

The solution set is $\{-1, 1, -i, i\}$.

Section 1.3: Complex Numbers; Quadratic Equations in the Complex Number System

73. $x^4 + 13x^2 + 36 = 0$

$$(x^2 + 9)(x^2 + 4) = 0$$

$$x^2 + 9 = 0 \quad \text{or} \quad x^2 + 4 = 0$$

$$x^2 = -9 \quad \text{or} \quad x^2 = -4$$

$$x = \pm\sqrt{-9} \quad \text{or} \quad x = \pm\sqrt{-4}$$

$$x = \pm 3i \quad \text{or} \quad x = \pm 2i$$

The solution set is $\{-3i, 3i, -2i, 2i\}$.

74. $x^4 + 3x^2 - 4 = 0$

$$(x^2 - 1)(x^2 + 4) = 0$$

$$(x-1)(x+1)(x^2 + 4) = 0$$

$$x-1=0 \quad \text{or} \quad x+1=0 \quad \text{or} \quad x^2 + 4 = 0$$

$$x=1 \quad \text{or} \quad x=-1 \quad \text{or} \quad x^2 = -4$$

$$x=1 \quad \text{or} \quad x=-1 \quad \text{or} \quad x = \pm\sqrt{-4} = \pm 2i$$

The solution set is $\{-1, 1, -2i, 2i\}$.

75. $3x^2 - 3x + 4 = 0$

$$a = 3, b = -3, c = 4$$

$$b^2 - 4ac = (-3)^2 - 4(3)(4) = 9 - 48 = -39$$

The equation has two complex solutions that are conjugates of each other.

76. $2x^2 - 4x + 1 = 0$

$$a = 2, b = -4, c = 1$$

$$b^2 - 4ac = (-4)^2 - 4(2)(1) = 16 - 8 = 8$$

The equation has two unequal real number solutions.

77. $2x^2 + 3x = 4$

$$2x^2 + 3x - 4 = 0$$

$$a = 2, b = 3, c = -4$$

$$b^2 - 4ac = 3^2 - 4(2)(-4) = 9 + 32 = 41$$

The equation has two unequal real solutions.

78. $x^2 + 6 = 2x$

$$x^2 - 2x + 6 = 0$$

$$a = 1, b = -2, c = 6$$

$$b^2 - 4ac = (-2)^2 - 4(1)(6) = 4 - 24 = -20$$

The equation has two complex solutions that are conjugates of each other.

79. $9x^2 - 12x + 4 = 0$

$$a = 9, b = -12, c = 4$$

$$b^2 - 4ac = (-12)^2 - 4(9)(4) = 144 - 144 = 0$$

The equation has a repeated real solution.

80. $4x^2 + 12x + 9 = 0$

$$a = 4, b = 12, c = 9$$

$$b^2 - 4ac = 12^2 - 4(4)(9) = 144 - 144 = 0$$

The equation has a repeated real solution.

81. The other solution is $\overline{2+3i} = 2-3i$.

82. The other solution is $\overline{4-i} = 4+i$.

83. $z + \overline{z} = 3 - 4i + \overline{3-4i} = 3 - 4i + 3 + 4i = 6$

$$\begin{aligned} 84. w - \overline{w} &= 8 + 3i - (\overline{8+3i}) \\ &= 8 + 3i - (8-3i) \\ &= 8 + 3i - 8 + 3i \\ &= 0 + 6i = 6i \end{aligned}$$

85. $z \cdot \overline{z} = (3-4i)(\overline{3-4i})$

$$= (3-4i)(3+4i)$$

$$= 9 + 12i - 12i - 16i^2$$

$$= 9 - 16(-1) = 25$$

86. $\overline{z-w} = \overline{3-4i-(8+3i)}$

$$= \overline{3-4i-8-3i}$$

$$= \overline{-5-7i}$$

$$= -5+7i$$

$$87. Z = \frac{V}{I} = \frac{18+i}{3-4i} = \frac{18+i}{3-4i} \cdot \frac{3+4i}{3+4i}$$

$$= \frac{54+72i+3i+4i^2}{9+12i-12i-16i^2} = \frac{54+75i-4}{9+16}$$

$$= \frac{50+75i}{25} = 2+3i$$

The impedance is $2+3i$ ohms.

Chapter 1: Equations and Inequalities

88.
$$\begin{aligned} \frac{1}{Z} &= \frac{1}{Z_1} + \frac{1}{Z_2} = \frac{1}{2+i} + \frac{1}{4-3i} = \frac{(4-3i)+(2+i)}{(2+i)(4-3i)} \\ &= \frac{6-2i}{8-6i+4i-3i^2} = \frac{6-2i}{8-2i+3} = \frac{6-2i}{11-2i} \end{aligned}$$

$$\begin{aligned} \text{So, } Z &= \frac{11-2i}{6-2i} = \frac{11-2i}{6-2i} \cdot \frac{6+2i}{6+2i} \\ &= \frac{66+22i-12i-4i^2}{36+12i-12i-4i^2} = \frac{66+10i+4}{36+4} \\ &= \frac{70+10i}{40} = \frac{7}{4} + \frac{1}{4}i \end{aligned}$$

The total impedance is $\frac{7}{4} + \frac{1}{4}i$ ohms.

89.
$$z + \bar{z} = (a+bi) + (\overline{a+bi})$$

$$\begin{aligned} &= a+bi+a-bi \\ &= 2a \end{aligned}$$

$$\begin{aligned} z - \bar{z} &= a+bi - (\overline{a+bi}) \\ &= a+bi - (a-bi) \\ &= a+bi-a+bi \\ &= 2bi \end{aligned}$$

90.
$$z = \overline{\overline{a+bi}} = \overline{a-bi} = a+bi = z$$

91.
$$\begin{aligned} \overline{z+w} &= \overline{(a+bi)+(c+di)} \\ &= \overline{(a+c)+(b+d)i} \\ &= (a+c)-(b+d)i \\ &= (a-bi)+(c-d)i \\ &= \overline{a+bi} + \overline{c+di} \\ &= \overline{z} + \overline{w} \end{aligned}$$

92.
$$\begin{aligned} \overline{z \cdot w} &= \overline{(a+bi) \cdot (c+di)} \\ &= \overline{ac+ad i+bc i+bd i^2} \\ &= \overline{(ac-bd)+(ad+bc)i} \\ &= (ac-bd)-(ad+bc)i \\ \overline{z} \cdot \overline{w} &= \overline{a+bi} \cdot \overline{c+di} \\ &= (a-bi)(c-di) \\ &= ac-ad i-bc i+bd i^2 \\ &= (ac-bd)-(ad+bc)i \end{aligned}$$

93 – 95. Answers will vary.

96. Answers will vary. A complex number is the sum or difference of two numbers (real and imaginary parts of the complex number) just as a binomial is the sum or difference of two monomial terms. We multiply two binomials by using the FOIL method, an approach we can also use to multiply two complex numbers.

97. Although the set of real numbers is a subset of the set of complex numbers, not all rules that work in the real number system can be used in the larger complex number system. The rule that allows us to write the product of two square roots as the square root of the product only works in the real number system. That is, $\sqrt{a} \cdot \sqrt{b} = \sqrt{ab}$ only when \sqrt{a} and \sqrt{b} are real numbers. In the complex number system we must first convert the radicals to complex form. In this case this means we need to write $\sqrt{-9}$ as $\sqrt{-1 \cdot 9} = \sqrt{9} \cdot \sqrt{-1} = 3i$. Then we can multiply to get $\sqrt{-9} \cdot \sqrt{-9} = 3i \cdot 3i = 9i^2 = 9(-1) = -9$.

Section 1.4

1. True
2. $\sqrt[3]{-8} = -2$
3. $6x^3 - 2x^2 = 2x^2(3x-1)$
4. False; you can also use the Quadratic Formula or completing the square.
5. quadratic in form
6. True
7. a
8. c
9. $\begin{aligned} \sqrt{2t-1} &= 1 \\ (\sqrt{2t-1})^2 &= 1^2 \\ 2t-1 &= 1 \\ 2t &= 2 \\ t &= 1 \end{aligned}$
Check: $\sqrt{2(1)-1} = \sqrt{1} = 1$
The solution set is {1}.

Section 1.4: Radical Equations; Equations Quadratic in Form; Factorable Equations

10. $\sqrt{3t+4} = 2$

$$(\sqrt{3t+4})^2 = 2^2$$

$$3t+4 = 4$$

$$3t = 0$$

$$t = 0$$

Check: $\sqrt{3(0)+4} = \sqrt{4} = 2$

The solution set is $\{0\}$.

11. $\sqrt{3t+4} = -6$

Since the principal square root is never negative, the equation has no real solution.

12. $\sqrt{5t+3} = -2$

Since the principal square root is never negative, the equation has no real solution.

13. $\sqrt[3]{1-2x} - 3 = 0$

$$\sqrt[3]{1-2x} = 3$$

$$(\sqrt[3]{1-2x})^3 = 3^3$$

$$1-2x = 27$$

$$-2x = 26$$

$$x = -13$$

Check: $\sqrt[3]{1-2(-13)} - 3 = \sqrt[3]{27} - 3 = 0$

The solution set is $\{-13\}$.

14. $\sqrt[3]{1-2x} - 1 = 0$

$$\sqrt[3]{1-2x} = 1$$

$$(\sqrt[3]{1-2x})^3 = 1^3$$

$$1-2x = 1$$

$$-2x = 0$$

$$x = 0$$

Check: $\sqrt[3]{1-2(0)} - 1 = \sqrt[3]{1} - 1 = 0$

The solution set is $\{0\}$.

15. $\sqrt[5]{x^2 + 2x} = -1$

$$(\sqrt[5]{x^2 + 2x})^5 = (-1)^5$$

$$x^2 + 2x = -1$$

$$x^2 + 2x + 1 = 0$$

$$(x+1)^2 = 0$$

$$x+1 = 0$$

$$x = -1$$

Check: $\sqrt[5]{(-1)^2 + 2(-1)} = \sqrt[5]{1-2} = \sqrt[5]{-1} = -1$

The solution set is $\{-1\}$.

16. $\sqrt[4]{x^2 + 16} = \sqrt{5}$

$$(\sqrt[4]{x^2 + 16})^4 = (\sqrt{5})^4$$

$$x^2 + 16 = 25$$

$$x^2 = 9$$

$$x = \pm 3$$

Check $-3: \sqrt[4]{(-3)^2 + 16} = \sqrt[4]{9+16} = \sqrt[4]{25} = \sqrt{5}$

Check $3: \sqrt[4]{(3)^2 + 16} = \sqrt[4]{9+16} = \sqrt[4]{25} = \sqrt{5}$

The solution set is $\{-3, 3\}$.

17. $x = 8\sqrt{x}$

$$(x)^2 = (8\sqrt{x})^2$$

$$x^2 = 64x$$

$$x^2 - 64x = 0$$

$$x(x-64) = 0$$

$$x = 0 \quad \text{or} \quad x = 64$$

Check 0: $0 = 8\sqrt{0}$ Check 64: $64 = 8\sqrt{64}$

$$0 = 0$$

$$64 = 64$$

The solution set is $\{0, 64\}$.

18. $x = 3\sqrt{x}$

$$(x)^2 = (3\sqrt{x})^2$$

$$x^2 = 9x$$

$$x^2 - 9x = 0$$

$$x(x-9) = 0$$

$$x = 0 \quad \text{or} \quad x = 9$$

Chapter 1: Equations and Inequalities

Check 0: $0 = 3\sqrt{0}$ Check 9: $9 = 3\sqrt{9}$
 $0 = 0$ $9 = 9$

The solution set is $\{0, 9\}$.

19. $\sqrt{15-2x} = x$

$$(\sqrt{15-2x})^2 = x^2$$

$$15-2x = x^2$$

$$x^2 + 2x - 15 = 0$$

$$(x+5)(x-3) = 0$$

$$x = -5 \text{ or } x = 3$$

Check -5: $\sqrt{15-2(-5)} = \sqrt{25} = 5 \neq -5$

Check 3: $\sqrt{15-2(3)} = \sqrt{9} = 3 = 3$

Disregard $x = -5$ as extraneous.

The solution set is $\{3\}$.

20. $\sqrt{12-x} = x$

$$(\sqrt{12-x})^2 = x^2$$

$$12-x = x^2$$

$$x^2 + x - 12 = 0$$

$$(x+4)(x-3) = 0$$

$$x = -4 \text{ or } x = 3$$

Check -4: $\sqrt{12-(-4)} = \sqrt{16} = 4 \neq -4$

Check 3: $\sqrt{12-3} = \sqrt{9} = 3 = 3$

Disregard $x = -4$ as extraneous.

The solution set is $\{3\}$.

21. $x = 2\sqrt{x-1}$

$$x^2 = (2\sqrt{x-1})^2$$

$$x^2 = 4(x-1)$$

$$x^2 = 4x - 4$$

$$x^2 - 4x + 4 = 0$$

$$(x-2)^2 = 0$$

$$x = 2$$

Check: $2 = 2\sqrt{2-1}$

$$2 = 2$$

The solution set is $\{2\}$.

22. $x = 2\sqrt{-x-1}$
 $x^2 = (2\sqrt{-x-1})^2$
 $x^2 = 4(-x-1)$
 $x^2 = -4x - 4$

$$x^2 + 4x + 4 = 0$$

$$(x+2)^2 = 0$$

$$x = -2$$

Check: $-2 = 2\sqrt{-(-2)-1}$
 $-2 \neq 2$

The equation has no real solution.

23. $\sqrt{x^2 - x - 4} = x + 2$

$$(\sqrt{x^2 - x - 4})^2 = (x+2)^2$$

$$x^2 - x - 4 = x^2 + 4x + 4$$

$$-8 = 5x$$

$$-\frac{8}{5} = x$$

Check: $\sqrt{\left(-\frac{8}{5}\right)^2 - \left(-\frac{8}{5}\right) - 4} = \left(-\frac{8}{5}\right) + 2$
 $\sqrt{\frac{64}{25} + \frac{8}{5} - 4} = \frac{2}{5}$
 $\sqrt{\frac{4}{25}} = \frac{2}{5}$
 $\frac{2}{5} = \frac{2}{5}$

The solution set is $\left\{-\frac{8}{5}\right\}$.

24. $\sqrt{3-x+x^2} = x-2$

$$(\sqrt{3-x+x^2})^2 = (x-2)^2$$

$$3-x+x^2 = x^2 - 4x + 4$$

$$3x = 1$$

$$x = \frac{1}{3}$$

Check: $\sqrt{3 - \left(\frac{1}{3}\right) + \left(\frac{1}{3}\right)^2} = \left(\frac{1}{3}\right) - 2$
 $\sqrt{3 - \frac{1}{3} + \frac{1}{9}} = -\frac{5}{3}$

Since the principal square root is always a non-negative number; $x = \frac{1}{3}$ does not check.

Therefore this equation has no real solution.

Section 1.4: Radical Equations; Equations Quadratic in Form; Factorable Equations

25. $3 + \sqrt{3x+1} = x$

$$\sqrt{3x+1} = x - 3$$

$$(\sqrt{3x+1})^2 = (x-3)^2$$

$$3x+1 = x^2 - 6x + 9$$

$$0 = x^2 - 9x + 8$$

$$0 = (x-1)(x-8)$$

$$x = 1 \text{ or } x = 8$$

Check 1: $3 + \sqrt{3(1)+1} = 3 + \sqrt{4} = 5 \neq 1$

Check 8: $3 + \sqrt{3(8)+1} = 3 + \sqrt{25} = 8 = 8$

Discard $x = 1$ as extraneous.

The solution set is $\{8\}$.

26. $2 + \sqrt{12-2x} = x$

$$\sqrt{12-2x} = x - 2$$

$$(\sqrt{12-2x})^2 = (x-2)^2$$

$$12-2x = x^2 - 4x + 4$$

$$0 = x^2 - 2x - 8$$

$$(x+2)(x-4) = 0$$

$$x = -2 \text{ or } x = 4$$

Check -2: $2 + \sqrt{12-2(-2)} = 2 + \sqrt{16} = 6 \neq -2$

Check 4: $2 + \sqrt{12-2(4)} = 2 + \sqrt{4} = 4 = 4$

Discard $x = -2$ as extraneous.

The solution set is $\{4\}$.

27. $\sqrt{3(x+10)} - 4 = x$

$$\sqrt{3(x+10)} = x + 4$$

$$(\sqrt{3(x+10)})^2 = (x+4)^2$$

$$3x+30 = x^2 + 8x + 16$$

$$0 = x^2 + 5x - 14$$

$$0 = (x+7)(x-2)$$

$$x = -7 \text{ or } x = 2$$

Check -7: $\sqrt{3(-7+10)} - 4 = \sqrt{9} - 4 = -1 \neq -7$

Check 2: $\sqrt{3(2+10)} - 4 = \sqrt{36} - 4 = 2 = 2$

Discard $x = -7$ as extraneous.

The solution set is $\{2\}$.

28. $\sqrt{1-x} - 3 = x + 2$

$$\sqrt{1-x} = x + 5$$

$$(\sqrt{1-x})^2 = (x+5)^2$$

$$1-x = x^2 + 10x + 25$$

$$0 = x^2 + 11x + 24$$

$$0 = (x+3)(x+8)$$

$$x = -3 \text{ or } x = -8$$

Check -3: $\sqrt{1-(-3)} - 3 = -3 + 2 \rightarrow -1 = -1$

Check -8: $\sqrt{1-(-8)} - 3 = -8 + 2 \rightarrow 0 = -6$

Discard $x = -8$ as extraneous.

The solution set is $\{-3\}$.

29. $\sqrt{2x+3} - \sqrt{x+1} = 1$

$$\sqrt{2x+3} = 1 + \sqrt{x+1}$$

$$(\sqrt{2x+3})^2 = (1 + \sqrt{x+1})^2$$

$$2x+3 = 1 + 2\sqrt{x+1} + x+1$$

$$x+1 = 2\sqrt{x+1}$$

$$(x+1)^2 = (2\sqrt{x+1})^2$$

$$x^2 + 2x + 1 = 4(x+1)$$

$$x^2 + 2x + 1 = 4x + 4$$

$$x^2 - 2x - 3 = 0$$

$$(x+1)(x-3) = 0$$

$$x = -1 \text{ or } x = 3$$

Check -1: $\sqrt{2(-1)+3} - \sqrt{-1+1}$

$$= \sqrt{1} - \sqrt{0} = 1 - 0 = 1 = 1$$

Check 3: $\sqrt{2(3)+3} - \sqrt{3+1}$

$$= \sqrt{9} - \sqrt{4} = 3 - 2 = 1 = 1$$

The solution set is $\{-1, 3\}$.

Chapter 1: Equations and Inequalities

30. $\sqrt{3x+7} + \sqrt{x+2} = 1$
 $\sqrt{3x+7} = 1 - \sqrt{x+2}$
 $(\sqrt{3x+7})^2 = (1 - \sqrt{x+2})^2$
 $3x+7 = 1 - 2\sqrt{x+2} + x+2$
 $2x+4 = -2\sqrt{x+2}$
 $-x-2 = \sqrt{x+2}$
 $(-x-2)^2 = (\sqrt{x+2})^2$
 $x^2 + 4x + 4 = x + 2$
 $x^2 + 3x + 2 = 0$
 $(x+1)(x+2) = 0$
 $x = -1 \text{ or } x = -2$
Check -1: $\sqrt{3(-1)+7} + \sqrt{-1+2} = \sqrt{4} + \sqrt{1} = 2 + 1 = 3 \neq 1$
Check -2: $\sqrt{3(-2)+7} + \sqrt{-2+2} = \sqrt{1} + \sqrt{0} = 1 + 0 = 1 = 1$
Discard $x = -1$ as extraneous.
The solution set is $\{-2\}$.

31. $\sqrt{3x+1} - \sqrt{x-1} = 2$
 $\sqrt{3x+1} = 2 + \sqrt{x-1}$
 $(\sqrt{3x+1})^2 = (2 + \sqrt{x-1})^2$
 $3x+1 = 4 + 4\sqrt{x-1} + x-1$
 $2x-2 = 4\sqrt{x-1}$
 $(2x-2)^2 = (4\sqrt{x-1})^2$
 $4x^2 - 8x + 4 = 16(x-1)$
 $x^2 - 2x + 1 = 4x - 4$
 $x^2 - 6x + 5 = 0$
 $(x-1)(x-5) = 0$
 $x = 1 \text{ or } x = 5$
Check 1: $\sqrt{3(1)+1} - \sqrt{1-1} = \sqrt{4} - \sqrt{0} = 2 - 0 = 2 = 2$
Check 5: $\sqrt{3(5)+1} - \sqrt{5-1} = \sqrt{16} - \sqrt{4} = 4 - 2 = 2 = 2$
The solution set is $\{1, 5\}$.

32. $\sqrt{3x-5} - \sqrt{x+7} = 2$
 $\sqrt{3x-5} = 2 + \sqrt{x+7}$
 $(\sqrt{3x-5})^2 = (2 + \sqrt{x+7})^2$
 $3x-5 = 4 + 4\sqrt{x+7} + x+7$
 $2x-16 = 4\sqrt{x+7}$
 $(2x-16)^2 = (4\sqrt{x+7})^2$
 $4x^2 - 64x + 256 = 16(x+7)$
 $4x^2 - 64x + 256 = 16x + 112$
 $4x^2 - 80x + 144 = 0$
 $x^2 - 20x + 36 = 0$
 $(x-2)(x-18) = 0$
 $x = 2 \text{ or } x = 18$
Check 2: $\sqrt{3(2)-5} - \sqrt{2+7} = \sqrt{1} - \sqrt{9} = 1 - 3 = -2 \neq 2$
Check 18: $\sqrt{3(18)-5} - \sqrt{18+7} = \sqrt{49} - \sqrt{25} = 7 - 5 = 2 = 2$
Discard $x = 2$ as extraneous.
The solution set is $\{18\}$.

33. $\sqrt{3-2\sqrt{x}} = \sqrt{x}$
 $(\sqrt{3-2\sqrt{x}})^2 = (\sqrt{x})^2$
 $3-2\sqrt{x} = x$
 $-2\sqrt{x} = x-3$
 $(-2\sqrt{x})^2 = (x-3)^2$
 $4x = x^2 - 6x + 9$
 $0 = x^2 - 10x + 9$
 $0 = (x-1)(x-9)$
 $x = 1 \text{ or } x = 9$

Check 1: $\sqrt{3-2\sqrt{1}} = \sqrt{1}$
 $\sqrt{3-2} = 1$
 $\sqrt{1} = 1$
Discard $x = 9$ as extraneous.
The solution set is $\{1\}$.

Check 9: $\sqrt{3-2\sqrt{9}} = \sqrt{9}$
 $\sqrt{3-2 \cdot 3} = 3$
 $\sqrt{-3} \neq 3$

34. $\sqrt{10+3\sqrt{x}} = \sqrt{x}$

$$\left(\sqrt{10+3\sqrt{x}}\right)^2 = (\sqrt{x})^2$$

$$10+3\sqrt{x} = x$$

$$3\sqrt{x} = x - 10$$

$$(3\sqrt{x})^2 = (x-10)^2$$

$$9x = x^2 - 20x + 100$$

$$0 = x^2 - 29x + 100$$

$$0 = (x-4)(x-25)$$

$$x = 4 \quad \text{or} \quad x = 25$$

Check 4:

$$\sqrt{10+3\sqrt{4}} = \sqrt{4}$$

$$\sqrt{10+3 \cdot 2} = 2$$

$$\sqrt{16} = 2$$

$$4 \neq 2$$

Check 25:

$$\sqrt{10+3\sqrt{25}} = \sqrt{25}$$

$$\sqrt{10+3 \cdot 5} = 5$$

$$\sqrt{25} = 5$$

$$5 = 5$$

Discard $x = 4$ as extraneous.

The solution set is $\{25\}$.

35. $(3x+1)^{1/2} = 4$

$$\left((3x+1)^{1/2}\right)^2 = (4)^2$$

$$3x+1 = 16$$

$$3x = 15$$

$$x = 5$$

Check: $(3(5)+1)^{1/2} = 16^{1/2} = 4$

The solution set is $\{5\}$.

36. $(3x-5)^{1/2} = 2$

$$\left((3x-5)^{1/2}\right)^2 = (2)^2$$

$$3x-5 = 4$$

$$3x = 9$$

$$x = 3$$

Check: $(3(3)-5)^{1/2} = 4^{1/2} = 2$

The solution set is $\{3\}$.

37. $(5x-2)^{1/3} = 2$

$$\left((5x-2)^{1/3}\right)^3 = (2)^3$$

$$5x-2 = 8$$

$$5x = 10$$

$$x = 2$$

Check: $(5(2)-2)^{1/3} = 8^{1/3} = 2$

The solution set is $\{2\}$.

38. $(2x+1)^{1/3} = -1$

$$\left((2x+1)^{1/3}\right)^3 = (-1)^3$$

$$2x+1 = -1$$

$$2x = -2$$

$$x = -1$$

Check: $(2(-1)+1)^{1/3} = (-1)^{1/3} = -1$

The solution set is $\{-1\}$.

39. $(x^2 + 9)^{1/2} = 5$

$$\left((x^2 + 9)^{1/2}\right)^2 = (5)^2$$

$$x^2 + 9 = 25$$

$$x^2 = 16$$

$$x = \pm\sqrt{16} = \pm 4$$

Check -4: $\left((-4)^2 + 9\right)^{1/2} = 25^{1/2} = 5$

Check 4: $\left((4)^2 + 9\right)^{1/2} = 25^{1/2} = 5$

The solution set is $\{-4, 4\}$.

40. $(x^2 - 16)^{1/2} = 9$

$$\left((x^2 - 16)^{1/2}\right)^2 = (9)^2$$

$$x^2 - 16 = 81$$

$$x^2 = 97$$

$$x = \pm\sqrt{97}$$

Check $-\sqrt{97}$: $\left(\left(-\sqrt{97}\right)^2 - 16\right)^{1/2} = 81^{1/2} = 9$

Check $\sqrt{97}$: $\left(\left(\sqrt{97}\right)^2 - 16\right)^{1/2} = 81^{1/2} = 9$

The solution set is $\{-\sqrt{97}, \sqrt{97}\}$.

Chapter 1: Equations and Inequalities

41. $x^{3/2} - 3x^{1/2} = 0$

$$x^{1/2}(x-3) = 0$$

$$x^{1/2} = 0 \text{ or } x-3 = 0$$

$$x=0 \text{ or } x=3$$

$$\text{Check 0: } 0^{3/2} - 3 \cdot 0^{1/2} = 0 - 0 = 0$$

$$\text{Check 3: } 3^{3/2} - 3 \cdot 3^{1/2} = 3\sqrt{3} - 3\sqrt{3} = 0$$

The solution set is $\{0,3\}$.

42. $x^{3/4} - 9x^{1/4} = 0$

$$x^{1/4}(x^{1/2} - 9) = 0$$

$$x^{1/4} = 0 \text{ or } x^{1/2} = 9$$

$$x=0 \text{ or } x=81$$

$$\text{Check 0: } 0^{3/4} - 9 \cdot 0^{1/4} = 0 - 0 = 0$$

$$\text{Check 81: } 81^{3/4} - 9 \cdot 81^{1/4} = 27 - 27 = 0$$

The solution set is $\{0,81\}$.

43. $x^4 - 5x^2 + 4 = 0$

$$(x^2 - 4)(x^2 - 1) = 0$$

$$x^2 - 4 = 0 \text{ or } x^2 - 1 = 0$$

$$x = \pm 2 \text{ or } x = \pm 1$$

The solution set is $\{-2,-1,1,2\}$.

44. $x^4 - 10x^2 + 25 = 0$

$$(x^2 - 5)(x^2 - 5) = 0$$

$$x^2 - 5 = 0$$

$$x = \pm\sqrt{5}$$

The solution set is $\{-\sqrt{5}, \sqrt{5}\}$.

45. $3x^4 - 2x^2 - 1 = 0$

$$(3x^2 + 1)(x^2 - 1) = 0$$

$$3x^2 + 1 = 0 \text{ or } x^2 - 1 = 0$$

$$3x^2 = -1 \text{ or } x^2 = 1$$

$$\text{Not real or } x = \pm 1$$

The solution set is $\{-1,1\}$.

46. $2x^4 - 5x^2 - 12 = 0$

$$(2x^2 + 3)(x^2 - 4) = 0$$

$$2x^2 + 3 = 0 \text{ or } x^2 - 4 = 0$$

$$2x^2 = -3 \text{ or } x^2 = 4$$

$$\text{Not real or } x = \pm 2$$

The solution set is $\{-2,2\}$.

47. $x^6 + 7x^3 - 8 = 0$

$$(x^3 + 8)(x^3 - 1) = 0$$

$$x^3 + 8 = 0 \text{ or } x^3 - 1 = 0$$

$$x^3 = -8 \text{ or } x^3 = 1$$

$$x = -2 \text{ or } x = 1$$

The solution set is $\{-2,1\}$.

48. $x^6 - 7x^3 - 8 = 0$

$$(x^3 - 8)(x^3 + 1) = 0$$

$$x^3 - 8 = 0 \text{ or } x^3 + 1 = 0$$

$$x^3 = 8 \text{ or } x^3 = -1$$

$$x = 2 \text{ or } x = -1$$

The solution set is $\{-1,2\}$.

49. $(x+2)^2 + 7(x+2) + 12 = 0$

Let $u = x+2$, so that $u^2 = (x+2)^2$.

$$u^2 + 7u + 12 = 0$$

$$(u+3)(u+4) = 0$$

$$u+3 = 0 \text{ or } u+4 = 0$$

$$u = -3 \text{ or } u = -4$$

$$x+2 = -3 \text{ or } x+2 = -4$$

$$x = -5 \text{ or } x = -6$$

The solution set is $\{-6,-5\}$.

50. $(2x+5)^2 - (2x+5) - 6 = 0$

Let $u = 2x+5$ so that $u^2 = (2x+5)^2$.

$$u^2 - u - 6 = 0$$

$$(u-3)(u+2) = 0$$

$$u-3 = 0 \text{ or } u+2 = 0$$

$$u = 3 \text{ or } u = -2$$

$$2x+5 = 3 \text{ or } 2x+5 = -2$$

$$x = -1 \text{ or } x = -\frac{7}{2}$$

The solution set is $\left\{-\frac{7}{2}, -1\right\}$.

51. $(3x+4)^2 - 6(3x+4) + 9 = 0$

Let $u = 3x+4$ so that $u^2 = (3x+4)^2$.

$$u^2 - 6u + 9 = 0$$

$$(u-3)^2 = 0$$

$$u-3 = 0$$

$$u = 3$$

$$3x+4=3$$

$$x = -\frac{1}{3}$$

The solution set is $\left\{-\frac{1}{3}\right\}$.

52. $(2-x)^2 + (2-x) - 20 = 0$

Let $u = 2-x$ so that $u^2 = (2-x)^2$.

$$u^2 + u - 20 = 0$$

$$(u+5)(u-4) = 0$$

$$u+5=0 \quad \text{or} \quad u-4=0$$

$$u=-5 \quad \text{or} \quad u=4$$

$$2-x=-5 \quad \text{or} \quad 2-x=4$$

$$x=7 \quad \text{or} \quad x=-2$$

The solution set is $\{-2, 7\}$.

53. $2(s+1)^2 - 5(s+1) = 3$

Let $u = s+1$ so that $u^2 = (s+1)^2$.

$$2u^2 - 5u = 3$$

$$2u^2 - 5u - 3 = 0$$

$$(2u+1)(u-3) = 0$$

$$2u+1=0 \quad \text{or} \quad u-3=0$$

$$u = -\frac{1}{2} \quad \text{or} \quad u = 3$$

$$s+1 = -\frac{1}{2} \quad \text{or} \quad s+1 = 3$$

$$s = -\frac{3}{2} \quad \text{or} \quad s = 2$$

The solution set is $\left\{-\frac{3}{2}, 2\right\}$.

54. $3(1-y)^2 + 5(1-y) + 2 = 0$

Let $u = 1-y$ so that $u^2 = (1-y)^2$.

$$3u^2 + 5u + 2 = 0$$

$$(3u+2)(u+1) = 0$$

$$3u+2=0 \quad \text{or} \quad u+1=0$$

$$u = -\frac{2}{3} \quad \text{or} \quad u = -1$$

$$1-y = -\frac{2}{3} \quad \text{or} \quad 1-y = -1$$

$$y = \frac{5}{3} \quad \text{or} \quad y = 2$$

The solution set is $\left\{\frac{5}{3}, 2\right\}$.

55. $x - 4x\sqrt{x} = 0$

$$x(1 - 4\sqrt{x}) = 0$$

$$x = 0 \quad \text{or} \quad 1 - 4\sqrt{x} = 0$$

$$1 = 4\sqrt{x}$$

$$\frac{1}{4} = \sqrt{x}$$

$$\left(\frac{1}{4}\right)^2 = (\sqrt{x})^2$$

$$\frac{1}{16} = x$$

Check:

$$x = 0: 0 - 4(0)\sqrt{0} = 0$$

$$0 = 0$$

$$x = \frac{1}{16}: \left(\frac{1}{16}\right) - 4\left(\frac{1}{16}\right)\sqrt{\frac{1}{16}} = 0$$

$$\frac{1}{16} - 4\left(\frac{1}{16}\right)\left(\frac{1}{4}\right) = 0$$

$$\frac{1}{16} - \frac{1}{16} = 0$$

$$0 = 0$$

The solution set is $\left\{0, \frac{1}{16}\right\}$.

56. $x + 8\sqrt{x} = 0$

$$8\sqrt{x} = -x$$

$$(8\sqrt{x})^2 = (-x)^2$$

$$64x = x^2$$

$$0 = x^2 - 64x$$

$$0 = x(x-64)$$

$$x = 0 \quad \text{or} \quad x = 64$$

Check: $x = 0: 0 + 8\sqrt{0} = 0$

$$0 = 0$$

$$x = 64: 64 + 8\sqrt{64} = 0$$

$$64 + 64 \neq 0$$

The solution set is $\{0\}$.

Chapter 1: Equations and Inequalities

57. $x + \sqrt{x} = 20$

Let $u = \sqrt{x}$ so that $u^2 = x$.

$$u^2 + u = 20$$

$$u^2 + u - 20 = 0$$

$$(u+5)(u-4) = 0$$

$$u+5=0 \quad \text{or} \quad u-4=0$$

$$u=-5 \quad \text{or} \quad u=4$$

$$\sqrt{x}=-5 \quad \text{or} \quad \sqrt{x}=4$$

$$\text{not possible or } x=16$$

$$\text{Check: } 16 + \sqrt{16} = 20$$

$$16+4=20$$

The solution set is $\{16\}$.

58. $x + \sqrt{x} = 6$

Let $u = \sqrt{x}$ so that $u^2 = x$.

$$u^2 + u = 6$$

$$u^2 + u - 6 = 0$$

$$(u+3)(u-2) = 0$$

$$u+3=0 \quad \text{or} \quad u-2=0$$

$$u=-3 \quad \text{or} \quad u=2$$

$$\sqrt{x}=-3 \quad \text{or} \quad \sqrt{x}=2$$

$$\text{not possible or } x=4$$

$$\text{Check: } 4 + \sqrt{4} = 6$$

$$4+2=6$$

The solution set is $\{4\}$.

59. $t^{1/2} - 2t^{1/4} + 1 = 0$

Let $u = t^{1/4}$ so that $u^2 = t^{1/2}$.

$$u^2 - 2u + 1 = 0$$

$$(u-1)^2 = 0$$

$$u-1=0$$

$$u=1$$

$$t^{1/4} = 1$$

$$t=1$$

$$\text{Check: } 1^{1/2} - 2(1)^{1/4} + 1 = 0$$

$$1-2+1=0$$

$$0=0$$

The solution set is $\{1\}$.

60. $z^{1/2} - 4z^{1/4} + 4 = 0$

Let $u = z^{1/4}$ so that $u^2 = z^{1/2}$.

$$u^2 - 4u + 4 = 0$$

$$(u-2)^2 = 0$$

$$u-2=0$$

$$u=2$$

$$z^{1/4} = 2$$

$$z=16$$

$$\text{Check: } 16^{1/2} - 4(16)^{1/4} + 4 = 0$$

$$4-8+4=0$$

$$0=0$$

The solution set is $\{16\}$.

61. $4x^{1/2} - 9x^{1/4} + 4 = 0$

Let $u = x^{1/4}$ so that $u^2 = x^{1/2}$.

$$4u^2 - 9u + 4 = 0$$

$$u = \frac{-(-9) \pm \sqrt{(-9)^2 - 4(4)(4)}}{2(4)} = \frac{9 \pm \sqrt{17}}{8}$$

$$x^{1/4} = \frac{9 \pm \sqrt{17}}{8}$$

$$x = \left(\frac{9 \pm \sqrt{17}}{8} \right)^4$$

$$\text{Check } x = \left(\frac{9 + \sqrt{17}}{8} \right)^4 :$$

$$4 \left(\left(\frac{9 + \sqrt{17}}{8} \right)^4 \right)^{1/2} - 9 \left(\left(\frac{9 + \sqrt{17}}{8} \right)^4 \right)^{1/4} + 4 = 0$$

$$4 \left(\frac{9 + \sqrt{17}}{8} \right)^2 - 9 \left(\frac{9 + \sqrt{17}}{8} \right) + 4 = 0$$

$$4 \frac{(9 + \sqrt{17})^2}{64} - 9 \left(\frac{9 + \sqrt{17}}{8} \right) + 4 = 0$$

$$64 \left(4 \frac{(9 + \sqrt{17})^2}{64} - 9 \left(\frac{9 + \sqrt{17}}{8} \right) + 4 \right) = (0)(64)$$

$$\begin{aligned}
 & 4(9 + \sqrt{17})^2 - 72(9 + \sqrt{17}) + 256 = 0 \\
 & 4(81 + 18\sqrt{17} + 17) - 72(9 + \sqrt{17}) + 256 = 0 \\
 & 324 + 72\sqrt{17} + 68 - 648 - 72\sqrt{17} + 256 = 0 \\
 & 0 = 0 \\
 \text{Check } x = & \left(\frac{9 - \sqrt{17}}{8} \right)^4 : \\
 & 4 \left(\left(\frac{9 - \sqrt{17}}{8} \right)^4 \right)^{1/2} - 9 \left(\left(\frac{9 - \sqrt{17}}{8} \right)^4 \right)^{1/4} + 4 = 0 \\
 & 4 \left(\frac{9 - \sqrt{17}}{8} \right)^2 - 9 \left(\frac{9 - \sqrt{17}}{8} \right) + 4 = 0 \\
 & 4(81 - 18\sqrt{17} + 17) - 72(9 - \sqrt{17}) + 256 = 0 \\
 & 324 - 72\sqrt{17} + 68 - 648 + 72\sqrt{17} + 256 = 0 \\
 & 0 = 0
 \end{aligned}$$

The solution set is $\left\{ \left(\frac{9 - \sqrt{17}}{8} \right)^4, \left(\frac{9 + \sqrt{17}}{8} \right)^4 \right\}$.

62. $x^{1/2} - 3x^{1/4} + 2 = 0$
Let $u = x^{1/4}$ so that $u^2 = x^{1/2}$.

$$\begin{aligned}
 u^2 - 3u + 2 &= 0 \\
 (u-2)(u-1) &= 0 \\
 u = 2 &\quad \text{or} \quad u = 1 \\
 x^{1/4} = 2 &\quad \text{or} \quad x^{1/4} = 1 \\
 x = 16 &\quad \text{or} \quad x = 1
 \end{aligned}$$

Check:

$$\begin{aligned}
 x = 16: 16^{1/2} - 3(16)^{1/4} + 2 &= 0 \\
 4 - 6 + 2 &= 0 \\
 0 &= 0
 \end{aligned}$$

$$\begin{aligned}
 x = 1: 1^{1/2} - 3(1)^{1/4} + 2 &= 0 \\
 1 - 3 + 2 &= 0 \\
 0 &= 0
 \end{aligned}$$

The solution set is $\{1, 16\}$.

63. $\sqrt[4]{5x^2 - 6} = x$

$$\begin{aligned}
 (\sqrt[4]{5x^2 - 6})^4 &= x^4 \\
 5x^2 - 6 &= x^4 \\
 0 &= x^4 - 5x^2 + 6
 \end{aligned}$$

Let $u = x^2$ so that $u^2 = x^4$.

$$\begin{aligned}
 0 &= u^2 - 5u + 6 \\
 0 &= (u-3)(u-2) \\
 u = 3 &\quad \text{or} \quad u = 2 \\
 x^2 = 3 &\quad \text{or} \quad x^2 = 2 \\
 x = \pm\sqrt{3} &\quad \text{or} \quad x = \pm\sqrt{2}
 \end{aligned}$$

Check:

$$\begin{aligned}
 x = -\sqrt{3}: \sqrt[4]{5(-\sqrt{3})^2 - 6} &= -\sqrt{3} \\
 \sqrt[4]{15-6} &= -\sqrt{3} \\
 \sqrt[4]{9} &\neq -\sqrt{3}
 \end{aligned}$$

$$\begin{aligned}
 x = \sqrt{3}: \sqrt[4]{5(\sqrt{3})^2 - 6} &= \sqrt{3} \\
 \sqrt[4]{15-6} &= \sqrt{3} \\
 \sqrt[4]{9} &= \sqrt{3} \\
 \sqrt{3} &= \sqrt{3}
 \end{aligned}$$

$$\begin{aligned}
 x = -\sqrt{2}: \sqrt[4]{5(-\sqrt{2})^2 - 6} &= -\sqrt{2} \\
 \sqrt[4]{10-6} &= -\sqrt{2} \\
 \sqrt[4]{4} &\neq -\sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 x = \sqrt{2}: \sqrt[4]{5(\sqrt{2})^2 - 6} &= \sqrt{2} \\
 \sqrt[4]{10-6} &= \sqrt{2} \\
 \sqrt[4]{4} &= \sqrt{2} \\
 \sqrt{2} &= \sqrt{2}
 \end{aligned}$$

The solution set is $\{\sqrt{2}, \sqrt{3}\}$.

64. $\sqrt[4]{4 - 5x^2} = x$

$$(\sqrt[4]{4 - 5x^2})^4 = x^4$$

$$4 - 5x^2 = x^4$$

$$0 = x^4 + 5x^2 - 4$$

Let $u = x^2$ so that $u^2 = x^4$.

$$0 = u^2 + 5u - 4$$

$$u = \frac{-5 \pm \sqrt{5^2 - 4(1)(-4)}}{2} = \frac{-5 \pm \sqrt{41}}{2}$$

$$x^2 = \frac{-5 \pm \sqrt{41}}{2}$$

$$x = \pm\sqrt{\frac{-5 \pm \sqrt{41}}{2}}$$

Chapter 1: Equations and Inequalities

Since $-5 - \sqrt{41} < 0$, $x = \pm\sqrt{\frac{-5 - \sqrt{41}}{2}}$ is not real.

Since x is a fourth root, $x = -\sqrt{\frac{-5 + \sqrt{41}}{2}}$ is also not real. Therefore, we have only one possible solution to check: $x = \sqrt{\frac{-5 + \sqrt{41}}{2}}$:

Check $x = \sqrt{\frac{-5 + \sqrt{41}}{2}}$:

$$\sqrt[4]{4 - 5\left(\pm\sqrt{\frac{-5 + \sqrt{41}}{2}}\right)^2} = \sqrt{\frac{-5 + \sqrt{41}}{2}}$$

$$\sqrt[4]{4 - 5\left(\frac{-5 + \sqrt{41}}{2}\right)} = \sqrt{\frac{-5 + \sqrt{41}}{2}}$$

$$\sqrt[4]{\frac{8 - 5(-5 + \sqrt{41})}{2}} = \sqrt{\frac{-5 + \sqrt{41}}{2}}$$

$$\sqrt[4]{\frac{33 - 5\sqrt{41}}{2}} = \sqrt{\frac{-5 + \sqrt{41}}{2}}$$

$$\sqrt[4]{\frac{66 - 10\sqrt{41}}{4}} = \sqrt{\frac{-5 + \sqrt{41}}{2}}$$

$$\sqrt[4]{\frac{25 - 10\sqrt{41} + 41}{4}} = \sqrt{\frac{-5 + \sqrt{41}}{2}}$$

$$\sqrt[4]{\frac{(-5 + \sqrt{41})^2}{4}} = \sqrt{\frac{-5 + \sqrt{41}}{2}}$$

$$\sqrt{\frac{-5 + \sqrt{41}}{2}} = \sqrt{\frac{-5 + \sqrt{41}}{2}}$$

The solution set is $\left\{ \sqrt{\frac{-5 + \sqrt{41}}{2}} \right\}$.

65. $x^2 + 3x + \sqrt{x^2 + 3x} = 6$

Let $u = \sqrt{x^2 + 3x}$ so that $u^2 = x^2 + 3x$.

$$u^2 + u = 6$$

$$u^2 + u - 6 = 0$$

$$(u+3)(u-2) = 0$$

$$\begin{array}{lll} u = -3 & \text{or} & u = 2 \\ \sqrt{x^2 + 3x} = -3 & \text{or} & \sqrt{x^2 + 3x} = 2 \\ \text{Not possible} & \text{or} & x^2 + 3x = 4 \\ & & x^2 + 3x - 4 = 0 \\ & & (x+4)(x-1) = 0 \\ & & x = -4 \text{ or } x = 1 \end{array}$$

Check $x = -4$:

$$\begin{aligned} (-4)^2 + 3(-4) + \sqrt{(-4)^2 + 3(-4)} &= 6 \\ 16 - 12 + \sqrt{16 - 12} &= 6 \\ 16 - 12 + \sqrt{4} &= 6 \\ 6 &= 6 \end{aligned}$$

Check $x = 1$:

$$\begin{aligned} (1)^2 + 3(1) + \sqrt{(1)^2 + 3(1)} &= 6 \\ 1 + 3 + \sqrt{1 + 3} &= 6 \\ 4 + \sqrt{4} &= 6 \\ 6 &= 6 \end{aligned}$$

The solution set is $\{-4, 1\}$.

66. $x^2 - 3x - \sqrt{x^2 - 3x} = 2$

Let $u = \sqrt{x^2 - 3x}$ so that $u^2 = x^2 - 3x$.

$$u^2 - u = 2$$

$$u^2 - u - 2 = 0$$

$$(u+1)(u-2) = 0$$

$$u = -1 \quad \text{or} \quad u = 2$$

$$\sqrt{x^2 - 3x} = -1 \quad \text{or} \quad \sqrt{x^2 - 3x} = 2$$

$$\text{Not possible} \quad \text{or} \quad x^2 - 3x = 4$$

$$x^2 - 3x - 4 = 0$$

$$(x-4)(x+1) = 0$$

$$x = 4 \text{ or } x = -1$$

Check $x = 4$:

$$\begin{aligned} (4)^2 - 3(4) - \sqrt{(4)^2 - 3(4)} &= 16 - 12 - \sqrt{4} \\ &= 4 - 2 = 2 \end{aligned}$$

Check $x = -1$:

$$\begin{aligned} (-1)^2 - 3(-1) - \sqrt{(-1)^2 - 3(-1)} &= 1 + 3 - \sqrt{4} \\ &= 4 - 2 = 2 \end{aligned}$$

The solution set is $\{-1, 4\}$.

Section 1.4: Radical Equations; Equations Quadratic in Form; Factorable Equations

67. $\frac{1}{(x+1)^2} = \frac{1}{x+1} + 2$

Let $u = \frac{1}{x+1}$ so that $u^2 = \left(\frac{1}{x+1}\right)^2$.

$$u^2 = u + 2$$

$$u^2 - u - 2 = 0$$

$$(u+1)(u-2) = 0$$

$$u = -1 \quad \text{or} \quad u = 2$$

$$\frac{1}{x+1} = -1 \quad \text{or} \quad \frac{1}{x+1} = 2$$

$$1 = -x - 1 \quad \text{or} \quad 1 = 2x + 2$$

$$x = -2 \quad \text{or} \quad -2x = 1$$

$$x = -\frac{1}{2}$$

Check:

$$x = -2: \frac{1}{(-2+1)^2} = \frac{1}{-2+1} + 2$$

$$1 = -1 + 2$$

$$1 = 1$$

$$x = -\frac{1}{2}: \frac{1}{\left(-\frac{1}{2}+1\right)^2} = \frac{1}{\left(-\frac{1}{2}+1\right)} + 2$$

$$4 = 2 + 2$$

$$4 = 4$$

The solution set is $\{-2, -\frac{1}{2}\}$.

68. $\frac{1}{(x-1)^2} + \frac{1}{x-1} = 12$

Let $u = \frac{1}{x-1}$ so that $u^2 = \left(\frac{1}{x-1}\right)^2$.

$$u^2 + u = 12$$

$$u^2 + u - 12 = 0$$

$$(u+4)(u-3) = 0$$

$$u = -4 \quad \text{or} \quad u = 3$$

$$\frac{1}{x-1} = -4 \quad \text{or} \quad \frac{1}{x-1} = 3$$

$$1 = -4x + 4 \quad \text{or} \quad 1 = 3x - 3$$

$$4x = 3 \quad \text{or} \quad 4 = 3x$$

$$x = \frac{3}{4} \quad \text{or} \quad x = \frac{4}{3}$$

Check:

$$x = \frac{3}{4}: \frac{1}{\left(\frac{3}{4}-1\right)^2} + \frac{1}{\left(\frac{3}{4}-1\right)} = 12$$

$$\frac{1}{\left(\frac{1}{16}\right)} + \frac{1}{\left(-\frac{1}{4}\right)} = 12$$

$$16 - 4 = 12$$

$$12 = 12$$

$$x = \frac{4}{3}: \frac{1}{\left(\frac{4}{3}-1\right)^2} + \frac{1}{\left(\frac{4}{3}-1\right)} = 12$$

$$\frac{1}{\left(\frac{1}{9}\right)} + \frac{1}{\left(\frac{1}{3}\right)} = 12$$

$$9 + 3 = 12$$

$$12 = 12$$

The solution set is $\left\{\frac{3}{4}, \frac{4}{3}\right\}$.

69. $3x^{-2} - 7x^{-1} - 6 = 0$

Let $u = x^{-1}$ so that $u^2 = x^{-2}$.

$$3u^2 - 7u - 6 = 0$$

$$(3u+2)(u-3) = 0$$

$$u = -\frac{2}{3} \quad \text{or} \quad u = 3$$

$$x^{-1} = -\frac{2}{3} \quad \text{or} \quad x^{-1} = 3$$

$$(x^{-1})^{-1} = \left(-\frac{2}{3}\right)^{-1} \quad \text{or} \quad (x^{-1})^{-1} = (3)^{-1}$$

$$x = -\frac{3}{2} \quad \text{or} \quad x = \frac{1}{3}$$

Check:

$$x = -\frac{3}{2}: 3\left(-\frac{3}{2}\right)^{-2} - 7\left(-\frac{3}{2}\right)^{-1} - 6 = 0$$

$$3\left(\frac{4}{9}\right) - 7\left(-\frac{2}{3}\right) - 6 = 0$$

$$\frac{4}{3} + \frac{14}{3} - 6 = 0$$

$$0 = 0$$

$$x = \frac{1}{3}: 3\left(\frac{1}{3}\right)^{-2} - 7\left(\frac{1}{3}\right)^{-1} - 6 = 0$$

$$3(9) - 7(3) - 6 = 0$$

$$27 - 21 - 6 = 0$$

$$0 = 0$$

The solution set is $\left\{-\frac{3}{2}, \frac{1}{3}\right\}$.

Chapter 1: Equations and Inequalities

70. $2x^{-2} - 3x^{-1} - 4 = 0$

Let $u = x^{-1}$ so that $u^2 = x^{-2}$.

$$2u^2 - 3u - 4 = 0$$

$$u = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(2)(-4)}}{2(2)} = \frac{3 \pm \sqrt{41}}{4}$$

$$u = \frac{3 + \sqrt{41}}{4} \quad \text{or} \quad u = \frac{3 - \sqrt{41}}{4}$$

$$x^{-1} = \frac{3 + \sqrt{41}}{4} \quad \text{or} \quad x^{-1} = \frac{3 - \sqrt{41}}{4}$$

$$(x^{-1})^{-1} = \left(\frac{3 + \sqrt{41}}{4}\right)^{-1} \quad \text{or} \quad (x^{-1})^{-1} = \left(\frac{3 - \sqrt{41}}{4}\right)^{-1}$$

$$x = \frac{4}{3 + \sqrt{41}} \left(\frac{3 - \sqrt{41}}{3 - \sqrt{41}} \right) \quad \text{or} \quad x = \frac{4}{3 - \sqrt{41}} \left(\frac{3 + \sqrt{41}}{3 + \sqrt{41}} \right)$$

$$= \frac{12 - 4\sqrt{41}}{-32} \quad = \frac{12 + 4\sqrt{41}}{-32}$$

$$= \frac{-3 + \sqrt{41}}{8} \quad = \frac{-3 - \sqrt{41}}{8}$$

Check $x = \frac{-3 + \sqrt{41}}{8}$:

$$2\left(\frac{-3 + \sqrt{41}}{8}\right)^{-2} - 3\left(\frac{-3 + \sqrt{41}}{8}\right)^{-1} - 4 = 0$$

$$2\left(\frac{64}{(-3 + \sqrt{41})^2}\right) - 3\left(\frac{8}{-3 + \sqrt{41}}\right) - 4 = 0$$

$$2(64) - 3(8)(-3 + \sqrt{41}) - 4(-3 + \sqrt{41})^2 = 0$$

$$128 + 72 - 24\sqrt{41} - 4(9 - 6\sqrt{41} + 41) = 0$$

$$128 + 72 - 24\sqrt{41} - 36 + 24\sqrt{41} - 164 = 0$$

$$0 = 0$$

Check $x = \frac{-3 - \sqrt{41}}{8}$:

$$2\left(\frac{-3 - \sqrt{41}}{8}\right)^{-2} - 3\left(\frac{-3 - \sqrt{41}}{8}\right)^{-1} - 4 = 0$$

$$2\left(\frac{64}{(-3 - \sqrt{41})^2}\right) - 3\left(\frac{8}{-3 - \sqrt{41}}\right) - 4 = 0$$

$$2(64) - 3(8)(-3 - \sqrt{41}) - 4(-3 - \sqrt{41})^2 = 0$$

$$128 + 72 + 24\sqrt{41} - 4(9 + 6\sqrt{41} + 41) = 0$$

$$128 + 72 + 24\sqrt{41} - 36 - 24\sqrt{41} - 164 = 0$$

$$0 = 0$$

The solution set is $\left\{ \frac{-3 - \sqrt{41}}{8}, \frac{-3 + \sqrt{41}}{8} \right\}$.

71. $2x^{2/3} - 5x^{1/3} - 3 = 0$

Let $u = x^{1/3}$ so that $u^2 = x^{2/3}$.

$$2u^2 - 5u - 3 = 0$$

$$(2u+1)(u-3) = 0$$

$$u = -\frac{1}{2} \quad \text{or} \quad u = 3$$

$$x^{1/3} = -\frac{1}{2} \quad \text{or} \quad x^{1/3} = 3$$

$$(x^{1/3})^3 = \left(-\frac{1}{2}\right)^3 \quad \text{or} \quad (x^{1/3})^3 = (3)^3$$

$$x = -\frac{1}{8} \quad \text{or} \quad x = 27$$

Check $x = -\frac{1}{8}$: $2\left(-\frac{1}{8}\right)^{2/3} - 5\left(-\frac{1}{8}\right)^{1/3} - 3 = 0$

$$2\left(\frac{1}{4}\right) - 5\left(-\frac{1}{2}\right) - 3 = 0$$

$$\frac{1}{2} + \frac{5}{2} - 3 = 0$$

$$3 - 3 = 0$$

$$0 = 0$$

Check $x = 27$: $2(27)^{2/3} - 5(27)^{1/3} - 3 = 0$

$$2(9) - 5(3) - 3 = 0$$

$$18 - 15 - 3 = 0$$

$$3 - 3 = 0$$

$$0 = 0$$

The solution set is $\left\{ -\frac{1}{8}, 27 \right\}$.

72. $3x^{4/3} + 5x^{2/3} - 2 = 0$

Let $u = x^{2/3}$ so that $u^2 = x^{4/3}$.

$$3u^2 + 5u - 2 = 0$$

$$(3u-1)(u+2) = 0$$

$$u = \frac{1}{3} \quad \text{or} \quad u = -2$$

$$x^{2/3} = \frac{1}{3} \quad \text{or} \quad x^{2/3} = -2$$

$$(x^{2/3})^3 = \left(\frac{1}{3}\right)^3 \quad \text{or} \quad (x^{2/3})^3 = (-2)^3$$

$$x^2 = \frac{1}{27} \quad \text{or} \quad x^2 = -8$$

$$x = \pm\sqrt{\frac{1}{27}} \quad \text{not real}$$

$$\text{Check: } 3\left(\pm\sqrt{\frac{1}{27}}\right)^{4/3} + 5\left(\pm\sqrt{\frac{1}{27}}\right)^{2/3} - 2 = 0$$

$$3\left(\frac{1}{27}\right)^{2/3} + 5\left(\pm\frac{1}{27}\right)^{1/3} - 2 = 0$$

$$3\left(\frac{1}{3}\right)^2 + 5\left(\frac{1}{3}\right) - 2 = 0$$

$$3\left(\frac{1}{9}\right) + \frac{5}{3} - 2 = 0$$

$$\frac{1}{3} + \frac{5}{3} - 2 = 0$$

$$2 - 2 = 0$$

$$0 = 0$$

$$\text{Note: } \pm\sqrt{\frac{1}{27}} = \pm\sqrt{\frac{3}{81}} = \pm\frac{\sqrt{3}}{9}$$

$$\text{The solution set is } \left\{-\frac{\sqrt{3}}{9}, \frac{\sqrt{3}}{9}\right\}.$$

73. $\left(\frac{v}{v+1}\right)^2 + \frac{2v}{v+1} = 8$

$$\left(\frac{v}{v+1}\right)^2 + 2\left(\frac{v}{v+1}\right) = 8$$

$$\text{Let } u = \frac{v}{v+1} \text{ so that } u^2 = \left(\frac{v}{v+1}\right)^2.$$

$$u^2 + 2u = 8$$

$$u^2 + 2u - 8 = 0$$

$$(u+4)(u-2) = 0$$

$$u = -4 \quad \text{or} \quad u = 2$$

$$\frac{v}{v+1} = -4 \quad \text{or} \quad \frac{v}{v+1} = 2$$

$$v = -4v - 4 \quad \text{or} \quad v = 2v + 2$$

$$v = -\frac{4}{5} \quad \text{or} \quad v = -2$$

$$\text{Check } v = -\frac{4}{5}: \left(\frac{-\frac{4}{5}}{-\frac{4}{5}+1}\right)^2 + \frac{2\left(-\frac{4}{5}\right)}{\left(-\frac{4}{5}\right)+1} = 8$$

$$\left(\frac{\frac{16}{25}}{\frac{1}{25}}\right) + \left(\frac{-\frac{8}{5}}{\frac{1}{5}}\right) = 8 \\ 16 - 8 = 8 \\ 8 = 8$$

$$\text{Check } v = -2: \left(\frac{-2}{-2+1}\right)^2 + \frac{2(-2)}{(-2)+1} = 8 \\ 4 + 4 = 8 \\ 8 = 8$$

The solution set is $\left\{-2, -\frac{4}{5}\right\}$.

74. $\left(\frac{y}{y-1}\right)^2 = 6\left(\frac{y}{y-1}\right) + 7$

$$\text{Let } u = \frac{y}{y-1} \text{ so that } u^2 = \left(\frac{y}{y-1}\right)^2.$$

$$u^2 = 6u + 7$$

$$u^2 - 6u - 7 = 0$$

$$(u-7)(u+1) = 0$$

$$u = -1 \quad \text{or} \quad u = 7$$

$$\frac{y}{y-1} = -1 \quad \text{or} \quad \frac{y}{y-1} = 7$$

$$y = -y + 1 \quad \text{or} \quad y = 7y - 7$$

$$2y = 1 \quad \text{or} \quad -6y = -7$$

$$y = \frac{1}{2} \quad \text{or} \quad y = \frac{7}{6}$$

Chapter 1: Equations and Inequalities

Check $y = \frac{1}{2}$: $\left(\frac{\frac{1}{2}}{\frac{1}{2}-1}\right)^2 = 6\left(\frac{\frac{1}{2}}{\frac{1}{2}-1}\right) + 7$

$$\frac{\frac{1}{4}}{\frac{1}{4}} = 6 \cdot \frac{\frac{1}{2}}{\left(-\frac{1}{2}\right)} + 7$$

$$1 = 6(-1) + 7$$

$$1 = 1$$

Check $y = \frac{7}{6}$: $\left(\frac{\frac{7}{6}}{\frac{7}{6}-1}\right)^2 = 6\left(\frac{\frac{7}{6}}{\frac{7}{6}-1}\right) + 7$

$$\frac{\left(\frac{49}{36}\right)}{\left(\frac{1}{36}\right)} = 6\left(\frac{\left(\frac{7}{6}\right)}{\left(\frac{1}{6}\right)}\right) + 7$$

$$49 = 42 + 7$$

$$49 = 49$$

The solution set is $\left\{\frac{1}{2}, \frac{7}{6}\right\}$.

75. $x^3 - 9x = 0$
 $x(x^2 - 9) = 0$
 $x(x-3)(x+3) = 0$
 $x = 0 \text{ or } x-3=0 \text{ or } x+3=0$
 $x = 3 \quad x = -3$

The solution set is $\{-3, 0, 3\}$.

76. $x^4 - x^2 = 0$
 $x^2(x^2 - 1) = 0$
 $x^2(x-1)(x+1) = 0$
 $x^2 = 0 \text{ or } x-1=0 \text{ or } x+1=0$
 $x = 0 \quad x = 1 \quad x = -1$

The solution set is $\{-1, 0, 1\}$.

77. $4x^3 = 3x^2$
 $4x^3 - 3x^2 = 0$
 $x^2(4x-3) = 0$
 $x^2 = 0 \text{ or } 4x-3 = 0$
 $x = 0 \quad 4x = 3$
 $x = \frac{3}{4}$

The solution set is $\left\{0, \frac{3}{4}\right\}$.

78. $x^5 = 4x^3$
 $x^5 - 4x^3 = 0$
 $x^3(x^2 - 4) = 0$
 $x^3(x-2)(x+2) = 0$
 $x^3 = 0 \text{ or } x-2=0 \text{ or } x+2=0$
 $x = 0 \quad x = 2 \quad x = -2$

The solution set is $\{-2, 0, 2\}$.

79. $x^3 + x^2 - 20x = 0$
 $x(x^2 + x - 20) = 0$
 $x(x+5)(x-4) = 0$
 $x = 0 \text{ or } x+5=0 \text{ or } x-4=0$
 $x = -5 \quad x = 4$

The solution set is $\{-5, 0, 4\}$.

80. $x^3 + 6x^2 - 7x = 0$
 $x(x^2 + 6x - 7) = 0$
 $x(x+7)(x-1) = 0$
 $x = 0 \text{ or } x+7=0 \text{ or } x-1=0$
 $x = -7 \quad x = 1$

The solution set is $\{-7, 0, 1\}$.

81. $x^3 + x^2 - x - 1 = 0$
 $x^2(x+1) - 1(x+1) = 0$
 $(x+1)(x^2 - 1) = 0$
 $(x+1)(x-1)(x+1) = 0$
 $x+1 = 0 \text{ or } x-1 = 0$
 $x = -1 \quad x = 1$

The solution set is $\{-1, 1\}$.

Section 1.4: Radical Equations; Equations Quadratic in Form; Factorable Equations

82. $x^3 + 4x^2 - x - 4 = 0$

$$x^2(x+4) - 1(x+4) = 0$$

$$(x+4)(x^2 - 1) = 0$$

$$(x+4)(x-1)(x+1) = 0$$

$$x+4=0 \quad \text{or} \quad x-1=0 \quad \text{or} \quad x+1=0$$

$$x=-4$$

$$x=1$$

$$x=-1$$

The solution set is $\{-4, -1, 1\}$.

83. $x^3 - 3x^2 - 4x + 12 = 0$

$$x^2(x-3) - 4(x-3) = 0$$

$$(x-3)(x^2 - 4) = 0$$

$$(x-3)(x-2)(x+2) = 0$$

$$x-3=0 \quad \text{or} \quad x-2=0 \quad \text{or} \quad x+2=0$$

$$x=3$$

$$x=2$$

$$x=-2$$

The solution set is $\{-2, 2, 3\}$.

84. $x^3 - 3x^2 - x + 3 = 0$

$$x^2(x-3) - 1(x-3) = 0$$

$$(x-3)(x^2 - 1) = 0$$

$$(x-3)(x-1)(x+1) = 0$$

$$x-3=0 \quad \text{or} \quad x-1=0 \quad \text{or} \quad x+1=0$$

$$x=3$$

$$x=1$$

$$x=-1$$

The solution set is $\{-1, 1, 3\}$.

85. $2x^3 + 4 = x^2 + 8x$

$$2x^3 - x^2 - 8x + 4 = 0$$

$$x^2(2x-1) - 4(2x-1) = 0$$

$$(2x-1)(x^2 - 4) = 0$$

$$(2x-1)(x-2)(x+2) = 0$$

$$2x-1=0 \quad \text{or} \quad x-2=0 \quad \text{or} \quad x+2=0$$

$$2x=1$$

$$x=2$$

$$x=-2$$

$$x=\frac{1}{2}$$

The solution set is $\left\{-2, \frac{1}{2}, 2\right\}$.

86. $3x^3 + 4x^2 = 27x + 36$

$$3x^3 + 4x^2 - 27x - 36 = 0$$

$$x^2(3x+4) - 9(3x+4) = 0$$

$$(3x+4)(x^2 - 9) = 0$$

$$(3x+4)(x-3)(x+3) = 0$$

$$3x+4=0 \quad \text{or} \quad x-3=0 \quad \text{or} \quad x+3=0$$

$$3x=-4 \quad x=3 \quad x=-3$$

$$x=-\frac{4}{3}$$

The solution set is $\left\{-3, -\frac{4}{3}, 3\right\}$.

87. $5x^3 + 45x = 2x^2 + 18$

$$5x^3 - 2x^2 + 45x - 18 = 0$$

$$x^2(5x-2) + 9(5x-2) = 0$$

$$(5x-2)(x^2 + 9) = 0$$

$$5x-2=0 \quad \text{or} \quad x^2 + 9 = 0$$

$$5x=2 \quad x^2 = -9$$

$$x=\frac{2}{5} \quad \text{no real solutions}$$

The solution set is $\left\{\frac{2}{5}\right\}$.

88. $3x^3 + 12x = 5x^2 + 20$

$$3x^3 - 5x^2 + 12x - 20 = 0$$

$$x^2(3x-5) + 4(3x-5) = 0$$

$$(3x-5)(x^2 + 4) = 0$$

$$3x-5=0 \quad \text{or} \quad x^2 + 4 = 0$$

$$3x=5 \quad x^2 = -4$$

$$x=\frac{5}{3} \quad \text{no real solutions}$$

The solution set is $\left\{\frac{5}{3}\right\}$.

89. $x(x^2 - 3x)^{1/3} + 2(x^2 - 3x)^{4/3} = 0$

$$(x^2 - 3x)^{1/3} [x + 2(x^2 - 3x)] = 0$$

$$(x^2 - 3x)^{1/3} (x + 2x^2 - 6x) = 0$$

$$(x^2 - 3x)^{1/3} (2x^2 - 5x) = 0$$

Chapter 1: Equations and Inequalities

$$\begin{aligned} (x^2 - 3x)^{1/3} &= 0 \quad \text{or} \quad 2x^2 - 5x = 0 \\ x^2 - 3x &= 0 \quad \text{or} \quad 2x^2 - 5x = 0 \\ x(x-3) &= 0 \quad \text{or} \quad x(2x-5) = 0 \\ x = 0 \text{ or } x &= 3 \quad \text{or} \quad x = 0 \text{ or } x = \frac{5}{2} \end{aligned}$$

The solution set is $\left\{0, \frac{5}{2}, 3\right\}$.

$$\begin{aligned} 90. \quad 3x(x^2 + 2x)^{1/2} - 2(x^2 + 2x)^{3/2} &= 0 \\ (x^2 + 2x)^{1/2} [3x - 2(x^2 + 2x)] &= 0 \\ (x^2 + 2x)^{1/2} (3x - 2x^2 - 4x) &= 0 \\ (x^2 + 2x)^{1/2} (-2x^2 - x) &= 0 \\ (x^2 + 2x)^{1/2} &= 0 \quad \text{or} \quad -2x^2 - x = 0 \\ x^2 + 2x &= 0 \quad \text{or} \quad 2x^2 + x = 0 \\ x(x+2) &= 0 \quad \text{or} \quad x(2x+1) = 0 \\ x = 0 \text{ or } x &= -2 \quad \text{or} \quad x = 0 \text{ or } x = -\frac{1}{2} \end{aligned}$$

Check $x = 0$:

$$\begin{aligned} 3 \cdot 0(0^2 + 2 \cdot 0)^{1/2} - 2(0^2 + 2 \cdot 0)^{3/2} &= 0 \\ 3 \cdot 0(0)^{1/2} - 2(0)^{3/2} &= 0 \\ 0 &= 0 \end{aligned}$$

Check $x = -2$:

$$\begin{aligned} 3(-2)((-2)^2 + 2(-2))^{1/2} - 2((-2)^2 + 2(-2))^{3/2} &= 0 \\ 3(-2)(4-4)^{1/2} - 2(4-4)^{3/2} &= 0 \\ 3(-2)(0)^{1/2} - 2(0)^{3/2} &= 0 \\ 3(-2)(0) - 2(0) &= 0 \\ 0 &= 0 \end{aligned}$$

Check $x = -\frac{1}{2}$:

$$\begin{aligned} 3\left(-\frac{1}{2}\right)\left(\left(-\frac{1}{2}\right)^2 + 2\left(-\frac{1}{2}\right)\right)^{1/2} - 2\left(\left(-\frac{1}{2}\right)^2 + 2\left(-\frac{1}{2}\right)\right)^{3/2} &= 0 \\ 3\left(-\frac{1}{2}\right)\left(\frac{1}{4}-1\right)^{1/2} - 2\left(\frac{1}{4}-1\right)^{3/2} &= 0 \\ 3\left(-\frac{1}{2}\right)\left(-\frac{3}{4}\right)^{1/2} - 2\left(-\frac{3}{4}\right)^{3/2} &= 0 \\ \text{Not real} & \end{aligned}$$

The solution set is $\{-2, 0\}$.

$$\begin{aligned} 91. \quad x - 4x^{1/2} + 2 &= 0 \\ \text{Let } u = x^{1/2} \text{ so that } u^2 &= x^2. \\ u^2 - 4u + 2 &= 0 \\ u = \frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(2)}}{2} &= \frac{4 \pm \sqrt{8}}{2} = \frac{4 \pm 2\sqrt{2}}{2} = 2 \pm \sqrt{2} \\ u = 2 + \sqrt{2} \quad \text{or} \quad u &= 2 - \sqrt{2} \\ x^{1/2} = 2 + \sqrt{2} \quad \text{or} \quad x^{1/2} &= 2 - \sqrt{2} \\ (x^{1/2})^2 = (2 + \sqrt{2})^2 \quad \text{or} \quad (x^{1/2})^2 &= (2 - \sqrt{2})^2 \\ x = (2 + \sqrt{2})^2 \quad \text{or} \quad x &= (2 - \sqrt{2})^2 \end{aligned}$$

Check $x = (2 + \sqrt{2})^2$:

$$\begin{aligned} (2 + \sqrt{2})^2 - 4(2 + \sqrt{2}) + 2 &= 0 \\ 4 + 4\sqrt{2} + 2 - 8 - 4\sqrt{2} + 2 &= 0 \\ 0 &= 0 \end{aligned}$$

Check $x = (2 - \sqrt{2})^2$:

$$\begin{aligned} (2 - \sqrt{2})^2 - 4(2 - \sqrt{2}) + 2 &= 0 \\ 4 - 4\sqrt{2} + 2 - 8 + 4\sqrt{2} + 2 &= 0 \\ 0 &= 0 \end{aligned}$$

The solution set is

$$\left\{(2 - \sqrt{2})^2, (2 + \sqrt{2})^2\right\} \approx \{0.34, 11.66\}.$$

$$\begin{aligned} 92. \quad x^{2/3} + 4x^{1/3} + 2 &= 0 \\ \text{Let } u = x^{1/3} \text{ so that } u^2 &= x^{2/3}. \\ u^2 + 4u + 2 &= 0 \\ u = \frac{-4 \pm \sqrt{4^2 - 4(1)(2)}}{2(1)} &= \frac{-4 \pm \sqrt{8}}{2} = \frac{-4 \pm 2\sqrt{2}}{2} = -2 \pm \sqrt{2} \\ u = -2 + \sqrt{2} \quad \text{or} \quad u &= -2 - \sqrt{2} \\ x^{1/3} = -2 + \sqrt{2} \quad \text{or} \quad x^{1/3} &= -2 - \sqrt{2} \\ x = (-2 + \sqrt{2})^3 \quad \text{or} \quad x &= (-2 - \sqrt{2})^3 \end{aligned}$$

Check $x = (-2 + \sqrt{2})^3$:

$$\begin{aligned} \left((-2 + \sqrt{2})^3\right)^{2/3} + 4\left((-2 + \sqrt{2})^3\right)^{1/3} + 2 &= 0 \\ (-2 + \sqrt{2})^2 + 4(-2 + \sqrt{2}) + 2 &= 0 \\ 4 - 4\sqrt{2} + 2 - 8 + 4\sqrt{2} + 2 &= 0 \\ 0 &= 0 \end{aligned}$$

Check $x = (-2 - \sqrt{2})^3$:

$$\begin{aligned} & \left((-2 - \sqrt{2})^3\right)^{2/3} + 4\left((-2 - \sqrt{2})^3\right)^{1/3} + 2 = 0 \\ & (-2 - \sqrt{2})^2 + 4(-2 - \sqrt{2}) + 2 = 0 \\ & 4 + 4\sqrt{2} + 2 - 8 - 4\sqrt{2} + 2 = 0 \\ & 0 = 0 \end{aligned}$$

The solution set is

$$\left\{(-2 - \sqrt{2})^3, (-2 + \sqrt{2})^3\right\} \approx \{-39.80, -0.20\}.$$

93. $x^4 + \sqrt{3}x^2 - 3 = 0$

Let $u = x^2$ so that $u^2 = x^4$.

$$u^2 + \sqrt{3}u - 3 = 0$$

$$\begin{aligned} u &= \frac{-\sqrt{3} \pm \sqrt{(\sqrt{3})^2 - 4(1)(-3)}}{2(1)} = \frac{-\sqrt{3} \pm \sqrt{15}}{2} \\ u &= \frac{-\sqrt{3} + \sqrt{15}}{2} \quad \text{or} \quad u = \frac{-\sqrt{3} - \sqrt{15}}{2} \\ x^2 &= \frac{-\sqrt{3} + \sqrt{15}}{2} \quad \text{or} \quad x^2 = \frac{-\sqrt{3} - \sqrt{15}}{2} \\ x &= \pm \sqrt{\frac{-\sqrt{3} + \sqrt{15}}{2}} \quad \text{or} \quad x = \pm \sqrt{\frac{-\sqrt{3} - \sqrt{15}}{2}} \\ &\quad \text{Not real} \end{aligned}$$

Check $x = \sqrt{\frac{-\sqrt{3} + \sqrt{15}}{2}}$:

$$\begin{aligned} & \left(\sqrt{\frac{-\sqrt{3} + \sqrt{15}}{2}}\right)^4 + \sqrt{3}\left(\sqrt{\frac{-\sqrt{3} + \sqrt{15}}{2}}\right)^2 - 3 = 0 \\ & \left(\frac{-\sqrt{3} + \sqrt{15}}{2}\right)^2 + \sqrt{3}\left(\frac{-\sqrt{3} + \sqrt{15}}{2}\right) - 3 = 0 \\ & \frac{3 - 2\sqrt{3}\sqrt{15} + 15}{4} + \frac{\sqrt{3}(-\sqrt{3}) + \sqrt{3}\sqrt{15}}{2} - 3 = 0 \\ & \frac{18 - 2\sqrt{45}}{4} + \frac{-3 + \sqrt{45}}{2} - 3 = 0 \\ & \frac{9 - \sqrt{45}}{2} + \frac{-3 + \sqrt{45}}{2} - 3 = 0 \\ & \frac{9 - \sqrt{45} - 3 + \sqrt{45}}{2} - 3 = 0 \\ & 3 - 3 = 0 \\ & 0 = 0 \end{aligned}$$

Check $x = -\sqrt{\frac{-\sqrt{3} + \sqrt{15}}{2}}$:

$$\begin{aligned} & \left(-\sqrt{\frac{-\sqrt{3} + \sqrt{15}}{2}}\right)^4 + \sqrt{3}\left(-\sqrt{\frac{-\sqrt{3} + \sqrt{15}}{2}}\right)^2 - 3 = 0 \\ & \left(\frac{-\sqrt{3} + \sqrt{15}}{2}\right)^2 + \sqrt{3}\left(\frac{-\sqrt{3} + \sqrt{15}}{2}\right) - 3 = 0 \\ & \frac{3 - 2\sqrt{3}\sqrt{15} + 15}{4} + \frac{\sqrt{3}(-\sqrt{3}) + \sqrt{3}\sqrt{15}}{2} - 3 = 0 \\ & \frac{18 - 2\sqrt{45}}{4} + \frac{-3 + \sqrt{45}}{2} - 3 = 0 \\ & \frac{9 - \sqrt{45}}{2} + \frac{-3 + \sqrt{45}}{2} - 3 = 0 \\ & \frac{9 - \sqrt{45} - 3 + \sqrt{45}}{2} - 3 = 0 \\ & 3 - 3 = 0 \\ & 0 = 0 \end{aligned}$$

The solution set is

$$\left\{-\sqrt{\frac{-\sqrt{3} + \sqrt{15}}{2}}, \sqrt{\frac{-\sqrt{3} + \sqrt{15}}{2}}\right\} \approx \{-1.03, 1.03\}.$$

94. $x^4 + \sqrt{2}x^2 - 2 = 0$

Let $u = x^2$ so that $u^2 = x^4$.

$$\begin{aligned} u^2 + \sqrt{2}u - 2 &= 0 \\ u &= \frac{-\sqrt{2} \pm \sqrt{(\sqrt{2})^2 - 4(1)(-2)}}{2(1)} = \frac{-\sqrt{2} \pm \sqrt{10}}{2} \\ u &= \frac{-\sqrt{2} + \sqrt{10}}{2} \quad \text{or} \quad u = \frac{-\sqrt{2} - \sqrt{10}}{2} \\ x^2 &= \frac{-\sqrt{2} + \sqrt{10}}{2} \quad \text{or} \quad x^2 = \frac{-\sqrt{2} - \sqrt{10}}{2} \\ x &= \pm \sqrt{\frac{-\sqrt{2} + \sqrt{10}}{2}} \quad \text{or} \quad x = \pm \sqrt{\frac{-\sqrt{2} - \sqrt{10}}{2}} \\ &\quad \text{Not real} \end{aligned}$$

Check $x = \sqrt{\frac{-\sqrt{2} + \sqrt{10}}{2}}$:

$$\begin{aligned} & \left(\sqrt{\frac{-\sqrt{2} + \sqrt{10}}{2}}\right)^4 + \sqrt{2}\left(\sqrt{\frac{-\sqrt{2} + \sqrt{10}}{2}}\right)^2 - 2 = 0 \\ & \left(\frac{-\sqrt{2} + \sqrt{10}}{2}\right)^2 + \sqrt{2}\left(\frac{-\sqrt{2} + \sqrt{10}}{2}\right) - 2 = 0 \\ & \frac{12 - 2\sqrt{20}}{4} + \frac{-2 + \sqrt{20}}{2} - 2 = 0 \\ & \frac{6 - \sqrt{20}}{2} - 2 + \frac{\sqrt{20}}{2} - 2 = 0 \\ & 2 - 2 = 0 \\ & 0 = 0 \end{aligned}$$

Check $x = -\sqrt{\frac{-\sqrt{2} + \sqrt{10}}{2}}$:

$$\left(-\sqrt{\frac{-\sqrt{2} + \sqrt{10}}{2}}\right)^4 + \sqrt{2}\left(-\sqrt{\frac{-\sqrt{2} + \sqrt{10}}{2}}\right)^2 - 2 = 0$$

$$\left(\frac{-\sqrt{2} + \sqrt{10}}{2}\right)^2 + \sqrt{2}\left(\frac{-\sqrt{2} + \sqrt{10}}{2}\right) - 2 = 0$$

$$\frac{12 - 2\sqrt{20}}{4} + \frac{-2 + \sqrt{20}}{2} - 2 = 0$$

$$\frac{6 - \sqrt{20} - 2 + \sqrt{20}}{2} - 2 = 0$$

$$2 - 2 = 0$$

$$0 = 0$$

The solution set is

$$\left\{-\sqrt{\frac{-\sqrt{2} + \sqrt{10}}{2}}, \sqrt{\frac{-\sqrt{2} + \sqrt{10}}{2}}\right\} \approx \{-0.93, 0.93\}.$$

95. $\pi(1+t)^2 = \pi + 1 + t$

Let $u = 1+t$ so that $u^2 = (1+t)^2$.

$$\pi u^2 = \pi + u$$

$$\pi u^2 - u - \pi = 0$$

$$u = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(\pi)(-\pi)}}{2(\pi)} = \frac{1 \pm \sqrt{1+4\pi^2}}{2\pi}$$

$$1+t = \frac{1 \pm \sqrt{1+4\pi^2}}{2\pi}$$

$$t = -1 + \frac{1 \pm \sqrt{1+4\pi^2}}{2\pi}$$

Check $t = -1 + \frac{1+\sqrt{1+4\pi^2}}{2\pi}$:

$$\pi\left(\frac{1+\sqrt{1+4\pi^2}}{2\pi}\right)^2 = \pi + \frac{1+\sqrt{1+4\pi^2}}{2\pi}$$

$$\pi\left(\frac{1+2\sqrt{1+4\pi^2}+1+4\pi^2}{4\pi^2}\right) = \pi + \frac{1+\sqrt{1+4\pi^2}}{2\pi}$$

$$\frac{2+2\sqrt{1+4\pi^2}+4\pi^2}{4\pi} = \frac{2\pi^2+1+\sqrt{1+4\pi^2}}{2\pi}$$

$$\frac{1+\sqrt{1+4\pi^2}+2\pi^2}{2\pi} = \frac{2\pi^2+1+\sqrt{1+4\pi^2}}{2\pi}$$

Check $t = -1 + \frac{1-\sqrt{1+4\pi^2}}{2\pi}$:

$$\pi\left(\frac{1-\sqrt{1+4\pi^2}}{2\pi}\right)^2 = \pi + \frac{1-\sqrt{1+4\pi^2}}{2\pi}$$

$$\pi\left(\frac{1-2\sqrt{1+4\pi^2}+1+4\pi^2}{4\pi^2}\right) = \pi + \frac{1-\sqrt{1+4\pi^2}}{2\pi}$$

$$\frac{2-2\sqrt{1+4\pi^2}+4\pi^2}{4\pi} = \frac{2\pi^2+1-\sqrt{1+4\pi^2}}{2\pi}$$

$$\frac{1-\sqrt{1+4\pi^2}+2\pi^2}{2\pi} = \frac{2\pi^2+1-\sqrt{1+4\pi^2}}{2\pi}$$

The solution set is

$$\left\{-1 + \frac{1-\sqrt{1+4\pi^2}}{2\pi}, -1 + \frac{1+\sqrt{1+4\pi^2}}{2\pi}\right\}$$

$$\approx \{-1.85, 0.17\}.$$

96. $\pi(1+r)^2 = 2 + \pi(1+r)$

Let $u = 1+r$ so that $u^2 = (1+r)^2$.

$$\pi u^2 = 2 + \pi u$$

$$\pi u^2 - \pi u - 2 = 0$$

$$u = \frac{-(-\pi) \pm \sqrt{(-\pi)^2 + 4(\pi)(-2)}}{2(\pi)}$$

$$= \frac{\pi \pm \sqrt{\pi^2 - 8\pi}}{2\pi}$$

$$1+r = \frac{\pi \pm \sqrt{\pi^2 + 8\pi}}{2\pi}$$

$$r = -1 + \frac{\pi \pm \sqrt{\pi^2 + 8\pi}}{2\pi}$$

Check $r = -1 + \frac{\pi + \sqrt{\pi^2 + 8\pi}}{2\pi}$:

$$\pi\left(\frac{\pi + \sqrt{\pi^2 + 8\pi}}{2\pi}\right)^2 = 2 + \pi\left(\frac{\pi + \sqrt{\pi^2 + 8\pi}}{2\pi}\right)$$

$$\pi\left(\frac{\pi^2 + 2\pi\sqrt{\pi^2 + 8\pi} + \pi^2 + 8\pi}{4\pi^2}\right) = 2 + \pi\left(\frac{\pi + \sqrt{\pi^2 + 8\pi}}{2\pi}\right)$$

$$\frac{2\pi^2 + 2\pi\sqrt{\pi^2 + 8\pi} + 8\pi}{4\pi} = 2 + \frac{\pi + \sqrt{\pi^2 + 8\pi}}{2}$$

$$\frac{\pi + \sqrt{\pi^2 + 8\pi} + 4}{2} = \frac{4 + \pi + \sqrt{\pi^2 + 8\pi}}{2}$$

Check $r = -1 + \frac{\pi - \sqrt{\pi^2 + 8\pi}}{2\pi}$:

$$\begin{aligned} \pi \left(\frac{\pi - \sqrt{\pi^2 + 8\pi}}{2\pi} \right)^2 &= 2 + \pi \left(\frac{\pi - \sqrt{\pi^2 + 8\pi}}{2\pi} \right) \\ \pi \left(\frac{\pi^2 - 2\pi\sqrt{\pi^2 + 8\pi} + \pi^2 + 8\pi}{4\pi^2} \right) &= 2 + \pi \left(\frac{\pi - \sqrt{\pi^2 + 8\pi}}{2\pi} \right) \\ \frac{2\pi^2 - 2\pi\sqrt{\pi^2 + 8\pi} + 8\pi}{4\pi} &= 2 + \frac{\pi - \sqrt{\pi^2 + 8\pi}}{2} \\ \frac{\pi - \sqrt{\pi^2 + 8\pi} + 4}{2} &= \frac{4 + \pi - \sqrt{\pi^2 + 8\pi}}{2} \end{aligned}$$

The solution set is

$$\left\{ -1 + \frac{\pi - \sqrt{\pi^2 + 8\pi}}{2\pi}, -1 + \frac{\pi + \sqrt{\pi^2 + 8\pi}}{2\pi} \right\} \\ \approx \{-1.44, 0.44\}.$$

97. $k^2 - k = 12$

$$k^2 - k - 12 = 0$$

$$(k-4)(k+3) = 0$$

$$k = 4 \quad \text{or} \quad k = -3$$

$$\frac{x+3}{x-3} = 4 \quad \text{or} \quad \frac{x+3}{x-3} = -3$$

$$x+3 = 4x-12 \quad \text{or} \quad x+3 = -3x+9$$

$$3x = 15 \quad \text{or} \quad 4x = 6$$

$$x = 5 \quad \text{or} \quad x = \frac{6}{4} = 1.5$$

Neither of these values causes a denominator to equal zero, so the solution set is $\{1.5, 5\}$.

98. $k^2 - 3k = 28$

$$k^2 - 3k - 28 = 0$$

$$(k+4)(k-7) = 0$$

$$k = -4 \quad \text{or} \quad k = 7$$

$$\frac{x+3}{x-4} = -4 \quad \text{or} \quad \frac{x+3}{x-4} = 7$$

$$x+3 = -4x+16 \quad \text{or} \quad x+3 = 7x-28$$

$$5x = 13 \quad \text{or} \quad -6x = -31$$

$$x = \frac{13}{5} = 2.6 \quad \text{or} \quad x = \frac{31}{6} \approx 5.17$$

Neither of these values causes a denominator to equal zero, so the solution set is

$$\left\{ \frac{13}{5}, \frac{31}{6} \right\} \approx \{2.6, 5.17\}.$$

99. Solve the equation $\frac{\sqrt{s}}{4} + \frac{s}{1100} = 4$.

$$\begin{aligned} \frac{s}{1100} + \frac{\sqrt{s}}{4} - 4 &= 0 \\ (1100)\left(\frac{s}{1100} + \frac{\sqrt{s}}{4} - 4\right) &= (0)(1100) \\ s + 275\sqrt{s} - 4400 &= 0 \end{aligned}$$

Let $u = \sqrt{s}$, so that $u^2 = s$.

$$\begin{aligned} u^2 + 275u - 4400 &= 0 \\ u &= \frac{-275 \pm \sqrt{275^2 - 4(1)(-4400)}}{2} \\ &= \frac{-275 \pm \sqrt{93,225}}{2} \end{aligned}$$

$$u \approx 15.1638 \text{ or } u \approx -290.1638$$

Since $u = \sqrt{s}$, it must be positive, so $s = u^2 \approx (15.1638)^2 \approx 229.94$

The distance to the water's surface is approximately 229.94 feet.

100. $T = \sqrt[4]{\frac{LH^2}{25}}$

Let $T = 4$ and $H = 10$, and solve for L .

$$\begin{aligned} 4 &= \sqrt[4]{\frac{L(10)^2}{25}} \\ 4 &= \sqrt[4]{4L} \end{aligned}$$

$$(4)^4 = \left(\sqrt[4]{4L}\right)^4$$

$$256 = 4L$$

$$64 = L$$

The crushing load is 64 tons.

101. $T = 2\pi\sqrt{\frac{l}{32}}$

Let $T = 16.5$ and solve for l .

Chapter 1: Equations and Inequalities

$$\begin{aligned} 16.5 &= 2\pi \sqrt{\frac{l}{32}} \\ \frac{16.5}{2\pi} &= \sqrt{\frac{l}{32}} \\ \left(\frac{16.5}{2\pi}\right)^2 &= \left(\sqrt{\frac{l}{32}}\right)^2 \\ \left(\frac{16.5}{2\pi}\right)^2 &= \frac{l}{32} \\ l &= 32 \left(\frac{16.5}{2\pi}\right)^2 \approx 220.7 \end{aligned}$$

The length was approximately 220.7 feet.

- 102.** Answers will vary. One example: $\sqrt{x+1} = -1$.
103. Answers will vary. One example: $x - \sqrt{x} - 2 = 0$.
104. Answers will vary.

- 105.** Jane did not check her solutions and included the extraneous solution, $x = -1$.


$$\begin{aligned} \sqrt{2x+3} - x &= 0 \\ \sqrt{2x+3} &= x \\ \left(\sqrt{2x+3}\right)^2 &= x^2 \\ 2x+3 &= x^2 \\ x^2 - 2x - 3 &= 0 \\ (x-3)(x+1) &= 0 \\ x = 3 \text{ or } x &= -1 \end{aligned}$$

Check:

$$\begin{array}{ll} \sqrt{2(3)+3}-3=0 & \sqrt{2(-1)+3}-(-1)=0 \\ \sqrt{9}-3=0 & \sqrt{1}+1=0 \\ 3-3=0 & 1+1=0 \\ 0=0 \quad \text{T} & 2 \neq 0 \end{array}$$

The solution set is $\{3\}$.




Section 1.5



- 2.** False. -5 is to the left of -2 on the number line, so $-5 > -2$.

3. closed interval
4. multiplication properties (for inequalities)
5. True. This follows from the addition property for inequalities.
6. True. This follows from the addition property for inequalities.
7. True;. This follows from the multiplication property for inequalities.
8. False. Since both sides of the inequality are being divided by a negative number, the sense, or direction, of the inequality must be reversed.
That is, $\frac{a}{c} > \frac{b}{c}$.
9. True
10. False; either or both endpoints could be any real number.
11. b
12. c
13. Interval: $[0, 2]$
Inequality: $0 \leq x \leq 2$
14. Interval: $(-1, 2)$
Inequality: $-1 < x < 2$
15. Interval: $[2, \infty)$
Inequality: $x \geq 2$
16. Interval: $(-\infty, 0]$
Inequality: $x \leq 0$
17. Interval: $[0, 3)$
Inequality: $0 \leq x < 3$
18. Interval: $(-1, 1]$
Inequality: $-1 < x \leq 1$
19. a. $3 < 5$
 $3+3 < 5+3$
 $6 < 8$
- b. $3 < 5$
 $3-5 < 5-5$
 $-2 < 0$

- c. $3 < 5$
 $3(3) < 3(5)$
 $9 < 15$
- d. $3 < 5$
 $-2(3) > -2(5)$
 $-6 > -10$
- 20.** a. $2 > 1$
 $2+3 > 1+3$
 $5 > 4$
- b. $2 > 1$
 $2-5 > 1-5$
 $-3 > -4$
- c. $2 > 1$
 $3(2) > 3(1)$
 $6 > 3$
- d. $2 > 1$
 $-2(2) < -2(1)$
 $-4 < -2$
- 21.** a. $4 > -3$
 $4+3 > -3+3$
 $7 > 0$
- b. $4 > -3$
 $4-5 > -3-5$
 $-1 > -8$
- c. $4 > -3$
 $3(4) > 3(-3)$
 $12 > -9$
- d. $4 > -3$
 $-2(4) < -2(-3)$
 $-8 < 6$
- 22.** a. $-3 > -5$
 $-3+3 > -5+3$
 $0 > -2$
- b. $-3 > -5$
 $-3-5 > -5-5$
 $-8 > -10$
- c. $-3 > -5$
 $3(-3) > 3(-5)$
 $-9 > -15$
- d. $-3 > -5$
 $-2(-3) < -2(-5)$
 $6 < 10$
- 23.** a. $2x+1 < 2$
 $2x+1+3 < 2+3$
 $2x+4 < 5$
- b. $2x+1 < 2$
 $2x+1-5 < 2-5$
 $2x-4 < -3$
- c. $2x+1 < 2$
 $3(2x+1) < 3(2)$
 $6x+3 < 6$
- d. $2x+1 < 2$
 $-2(2x+1) > -2(2)$
 $-4x-2 > -4$
- 24.** a. $1-2x > 5$
 $1-2x+3 > 5+3$
 $4-2x > 8$
- b. $1-2x > 5$
 $1-2x-5 > 5-5$
 $-4-2x > 0$
- c. $1-2x > 5$
 $3(1-2x) > 3(5)$
 $3-6x > 15$
- d. $1-2x > 5$
 $-2(1-2x) < -2(5)$
 $-2+4x < -10$
- 25.** $[0, 4]$

- 26.** $(-1, 5)$

- 27.** $[4, 6)$

- 28.** $(-2, 0)$


Chapter 1: Equations and Inequalities

29. $[4, \infty)$



30. $(-\infty, 5]$



31. $(-\infty, -4)$




32. $(1, \infty)$



33. $2 \leq x \leq 5$



34. $1 < x < 2$



35. $-3 < x < -2$



36. $0 \leq x < 1$




37. $x \geq 4$



38. $x \leq 2$



39. $x < -3$



40. $x > -8$



41. If $x < 5$, then $x - 5 < 0$.

42. If $x < -4$, then $x + 4 < 0$.

43. If $x > -4$, then $x + 4 > 0$.

44. If $x > 6$, then $x - 6 > 0$.

45. If $x \geq -4$, then $3x \geq -12$.

46. If $x \leq 3$, then $2x \leq 6$.

47. If $x > 6$, then $-2x < -12$.

48. If $x > -2$, then $-4x < 8$.

49. If $x \geq 5$, then $-4x \leq -20$.

50. If $x \leq -4$, then $-3x \geq 12$.

51. If $2x > 6$, then $x > 3$.

52. If $3x \leq 12$, then $x \leq 4$.

53. If $-\frac{1}{2}x \leq 3$, then $x \geq -6$.

54. If $-\frac{1}{4}x > 1$, then $x < -4$.

55. $x + 1 < 5$

$$x + 1 - 1 < 5 - 1$$

$$x < 4$$

$$\{x | x < 4\} \text{ or } (-\infty, 4)$$



56. $x - 6 < 1$

$$x - 6 + 6 < 1 + 6$$

$$x < 7$$

The solution set is $\{x | x < 7\}$ or $(-\infty, 7)$.




57. $1 - 2x \leq 3$

$$-2x \leq 2$$

$$x \geq -1$$

The solution set is $\{x | x \geq -1\}$ or $[-1, \infty)$.




58. $2 - 3x \leq 5$

$$-3x \leq 3$$

$$x \geq -1$$

The solution set is $\{x | x \geq -1\}$ or $[-1, \infty)$.



59. $3x - 7 > 2$

$$3x > 9$$

$$x > 3$$

The solution set is $\{x | x > 3\}$ or $(3, \infty)$.




60. $2x + 5 > 1$

$$2x > -4$$

$$x > -2$$

The solution set is $\{x | x > -2\}$ or $(-2, \infty)$.




61. $3x - 1 \geq 3 + x$

$$2x \geq 4$$

$$x \geq 2$$


The solution set is $\{x | x \geq 2\}$ or $[2, \infty)$.



62. $2x - 2 \geq 3 + x$

$$x \geq 5$$

The solution set is $\{x | x \geq 5\}$ or $[5, \infty)$.



63. $-2(x + 3) < 8$

$$-2x - 6 < 8$$

$$-2x < 14$$

$$x > -7$$

The solution set is $\{x | x > -7\}$ or $(-7, \infty)$.



64. $-3(1 - x) < 12$

$$-3 + 3x < 12$$

$$3x < 15$$

$$x < 5$$

The solution set is $\{x | x < 5\}$ or $(-\infty, 5)$.



65. $4 - 3(1 - x) \leq 3$


$$4 - 3 + 3x \leq 3$$

$$3x + 1 \leq 3$$

$$3x \leq 2$$

$$x \leq \frac{2}{3}$$

The solution set is $\{x | x \leq \frac{2}{3}\}$ or $(-\infty, \frac{2}{3}]$.



66. $8 - 4(2 - x) \leq -2x$

$$8 - 8 + 4x \leq -2x$$

$$4x \leq -2x$$

$$6x \leq 0$$

$$x \leq 0$$

The solution set is $\{x | x \leq 0\}$ or $(-\infty, 0]$.




67. $\frac{1}{2}(x - 4) > x + 8$

$$\frac{1}{2}x - 2 > x + 8$$

$$-\frac{1}{2}x > 10$$

$$x < -20$$

The solution set is $\{x | x < -20\}$ or $(-\infty, -20)$.



Chapter 1: Equations and Inequalities

68. $3x + 4 > \frac{1}{3}(x - 2)$


$$3x + 4 > \frac{1}{3}x - \frac{2}{3}$$

$$9x + 12 > x - 2$$

$$8x > -14$$

$$x > -\frac{7}{4}$$

The solution set is $\left\{x \mid x > -\frac{7}{4}\right\}$ or $(-\frac{7}{4}, \infty)$.




69. $\frac{x}{2} \geq 1 - \frac{x}{4}$

$$2x \geq 4 - x$$

$$3x \geq 4$$

$$x \geq \frac{4}{3}$$

The solution set is $\left\{x \mid x \geq \frac{4}{3}\right\}$ or $[\frac{4}{3}, \infty)$.




70. $\frac{x}{3} \geq 2 + \frac{x}{6}$

$$2x \geq 12 + x$$

$$x \geq 12$$

The solution set is $\{x \mid x \geq 12\}$ or $[12, \infty)$.




71. $0 \leq 2x - 6 \leq 4$

$$6 \leq 2x \leq 10$$

$$3 \leq x \leq 5$$

The solution set is $\{x \mid 3 \leq x \leq 5\}$ or $[3, 5]$.



72. $4 \leq 2x + 2 \leq 10$

$$2 \leq 2x \leq 8$$

$$1 \leq x \leq 4$$

The solution set is $\{x \mid 1 \leq x \leq 4\}$ or $[1, 4]$.




73. $-5 \leq 4 - 3x \leq 2$

$$-9 \leq -3x \leq -2$$

$$3 \geq x \geq \frac{2}{3}$$

The solution set is $\left\{x \mid \frac{2}{3} \leq x \leq 3\right\}$ or $[\frac{2}{3}, 3]$.




74. $-3 \leq 3 - 2x \leq 9$

$$-6 \leq -2x \leq 6$$

$$3 \geq x \geq -3$$

The solution set is $\{x \mid -3 \leq x \leq 3\}$ or $[-3, 3]$.




75. $-3 < \frac{2x - 1}{4} < 0$

$$-12 < 2x - 1 < 0$$

$$-11 < 2x < 1$$

$$-\frac{11}{2} < x < \frac{1}{2}$$

The solution set is $\left\{x \mid -\frac{11}{2} < x < \frac{1}{2}\right\}$ or $(-\frac{11}{2}, \frac{1}{2})$.




76. $0 < \frac{3x + 2}{2} < 4$

$$0 < 3x + 2 < 8$$

$$-2 < 3x < 6$$

$$-\frac{2}{3} < x < 2$$

The solution set is $\left\{x \mid -\frac{2}{3} < x < 2\right\}$ or $(-\frac{2}{3}, 2)$.



77. $1 < 1 - \frac{1}{2}x < 4$

$$0 < -\frac{1}{2}x < 3$$

$$0 > x > -6 \quad \text{or} \quad -6 < x < 0$$

The solution set is $\{x \mid -6 < x < 0\}$ or $(-6, 0)$.



78. $0 < 1 - \frac{1}{3}x < 1$

$$-1 < -\frac{1}{3}x < 0$$

$$3 > x > 0 \quad \text{or} \quad 0 < x < 3$$

The solution set is $\{x \mid 0 < x < 3\}$ or $(0, 3)$.



79. $(x+2)(x-3) > (x-1)(x+1)$

$$x^2 - x - 6 > x^2 - 1$$

$$-x - 6 > -1$$

$$-x > 5$$

$$x < -5$$

The solution set is $\{x \mid x < -5\}$ or $(-\infty, -5)$.



80. $(x-1)(x+1) > (x-3)(x+4)$

$$x^2 - 1 > x^2 + x - 12$$

$$-1 > x - 12$$

$$-x > -11$$

$$x < 11$$

The solution set is $\{x \mid x < 11\}$ or $(-\infty, 11)$.



81. $x(4x+3) \leq (2x+1)^2$


$$4x^2 + 3x \leq 4x^2 + 4x + 1$$

$$3x \leq 4x + 1$$

$$-x \leq 1$$

$$x \geq -1$$

The solution set is $\{x \mid x \geq -1\}$ or $[-1, \infty)$.




82. $x(9x-5) \leq (3x-1)^2$

$$9x^2 - 5x \leq 9x^2 - 6x + 1$$

$$-5x \leq -6x + 1$$

$$x \leq 1$$

The solution set is $\{x \mid x \leq 1\}$ or $(-\infty, 1]$.




83. $\frac{1}{2} \leq \frac{x+1}{3} < \frac{3}{4}$

$$6 \leq 4x + 4 < 9$$

$$2 \leq 4x < 5$$

$$\frac{1}{2} \leq x < \frac{5}{4}$$

The solution set is $\{x \mid \frac{1}{2} \leq x < \frac{5}{4}\}$ or $\left[\frac{1}{2}, \frac{5}{4}\right)$.




84. $\frac{1}{3} < \frac{x+1}{2} \leq \frac{2}{3}$

$$2 < 3x + 3 \leq 4$$

$$-1 < 3x \leq 1$$

$$-\frac{1}{3} < x \leq \frac{1}{3}$$

The solution set is $\{x \mid -\frac{1}{3} < x \leq \frac{1}{3}\}$ or $\left(-\frac{1}{3}, \frac{1}{3}\right]$.




85. $(4x+2)^{-1} < 0$

$$\frac{1}{4x+2} < 0$$

$$4x+2 < 0$$

$$x < -\frac{1}{2}$$

The solution set is $\{x \mid x < -\frac{1}{2}\}$ or $(-\infty, -\frac{1}{2})$.



Chapter 1: Equations and Inequalities

86. $(2x-1)^{-1} > 0$

$$\frac{1}{2x-1} > 0$$


Since $\frac{1}{2x-1} > 0$, this means $2x-1 > 0$.

Therefore,

$$2x-1 > 0$$

$$x > \frac{1}{2}$$

The solution set is $\left\{x \mid x > \frac{1}{2}\right\}$ or $\left(\frac{1}{2}, \infty\right)$.



87. $(2-7x)^{-1} \geq 5$

$$\frac{1}{2-7x} - 5 \geq 0$$

$$\frac{1}{2-7x} - 5 \geq 0$$

$$\frac{1-5(2-7x)}{2-7x} \geq 0$$

$$\frac{-9+35x}{2-7x} \geq 0$$

The zeros and values where the expression is undefined are $x = \frac{9}{35}$ and $x = \frac{2}{7}$

Interval	$(-\infty, \frac{9}{35})$	$(\frac{9}{35}, \frac{2}{7})$	$(\frac{2}{7}, \infty)$
Number Chosen	0	$\frac{27}{100}$	1
Value of f	$-\frac{9}{2}$	$\frac{45}{11}$	$-\frac{26}{5}$
Conclusion	Negative	Positive	Negative

We want to know where $f(x) \geq 0$, so the solution set is $\left\{x \mid x \geq \frac{9}{35} \text{ or } x < \frac{2}{7}\right\}$ or, using interval notation, $[\frac{9}{35}, \frac{2}{7})$. Note that $\frac{2}{7}$ is not in the solution set because $\frac{2}{7}$ is not in the domain of f .

The solution set is $[\frac{9}{35}, \frac{2}{7})$.



88. $2(3x+5)^{-1} \leq -3$

$$\frac{2}{(3x+5)} + 3 \leq 0$$

$$\frac{2+3(3x+5)}{2(3x+5)} \leq 0$$

$$\frac{17+9x}{(3x+5)} \leq 0$$

The zeros and values where the expression is undefined are $x = -\frac{17}{9}$ and $x = -\frac{5}{3}$

Interval	$(-\infty, -\frac{17}{9})$	$(-\frac{17}{9}, -\frac{5}{3})$	$(-\frac{5}{3}, \infty)$
Number Chosen	-2	-1.7	0
Value of f	1	-17	$\frac{17}{5}$
Conclusion	Positive	Negative	Positive

We want to know where $f(x) \leq 0$, so the solution set is $\left\{x \mid x \geq -\frac{17}{9} \text{ or } x < -\frac{5}{3}\right\}$ or, using interval notation, $[-\frac{17}{9}, -\frac{5}{3})$. Note that $-\frac{5}{3}$ is not in the solution set because $\frac{2}{7}$ is not in the domain of f .

The solution set is $[-\frac{17}{9}, -\frac{5}{3})$.



89. $0 < \frac{2}{x} < \frac{3}{5}$

$$0 < \frac{2}{x} \text{ and } \frac{2}{x} < \frac{3}{5}$$

Since $\frac{2}{x} > 0$, this means that $x > 0$. Therefore,


$$\frac{2}{x} < \frac{3}{5}$$

$$5x\left(\frac{2}{x}\right) < 5x\left(\frac{3}{5}\right)$$

$$10 < 3x$$

$$\frac{10}{3} < x$$

The solution set is $\left\{x \mid x > \frac{10}{3}\right\}$ or $\left(\frac{10}{3}, \infty\right)$.



90. $0 < \frac{4}{x} < \frac{2}{3}$

$$0 < \frac{4}{x} \text{ and } \frac{4}{x} < \frac{2}{3}$$

Since $\frac{4}{x} > 0$, this means that $x > 0$. Therefore,


$$\frac{4}{x} < \frac{2}{3}$$

$$3x\left(\frac{4}{x}\right) < 3x\left(\frac{2}{3}\right)$$

$$12 < 2x$$

$$6 < x$$

The solution set is $\{x | x > 6\}$ or $(6, \infty)$.



91. $0 < (2x-4)^{-1} < \frac{1}{2}$

$$0 < \frac{1}{2x-4} < \frac{1}{2}$$

$$0 < \frac{1}{2x-4} \text{ and } \frac{1}{2x-4} < \frac{1}{2}$$

Since $\frac{1}{2x-4} > 0$, this means that $2x-4 > 0$.

Therefore,

$$\frac{1}{2x-4} < \frac{1}{2}$$


$$\frac{1}{2(x-2)} < \frac{1}{2}$$

$$2(x-2)\left(\frac{1}{2(x-2)}\right) < 2(x-2)\left(\frac{1}{2}\right)$$

$$1 < x-2$$

$$3 < x$$

The solution set is $\{x | x > 3\}$ or $(3, \infty)$.



92. $0 < (3x+6)^{-1} < \frac{1}{3}$

$$0 < \frac{1}{3x+6} < \frac{1}{3}$$

$$0 < \frac{1}{3x+6} \text{ and } \frac{1}{3x+6} < \frac{1}{3}$$

Since $\frac{1}{3x+6} > 0$, this means that $3x+6 > 0$.

Therefore,

$$\frac{1}{3x+6} < \frac{1}{3}$$


$$\frac{1}{3(x+2)} < \frac{1}{3}$$

$$3(x+2)\left(\frac{1}{3(x+2)}\right) < 3(x+2)\left(\frac{1}{3}\right)$$

$$1 < x+2$$

$$-1 < x$$

The solution set is $\{x | x > -1\}$ or $(-1, \infty)$.



93. If $-1 < x < 1$, then

$$-1+4 < x+4 < 1+4$$

$$3 < x+4 < 5$$

So, $a = 3$ and $b = 5$.

94. If $-3 < x < 2$, then

$$-3-6 < x-6 < 2-6$$

$$-9 < x-6 < -4$$

So, $a = -9$ and $b = -4$.

95. If $2 < x < 3$, then

$$-4(2) < -4(x) < -4(3)$$

$$-12 < -4x < -8$$

So, $a = -12$ and $b = -8$.

96. If $-4 < x < 0$, then

$$\frac{1}{2}(-4) < \frac{1}{2}(x) < \frac{1}{2}(0)$$

$$-2 < \frac{1}{2}x < 0$$

So, $a = -2$ and $b = 0$.

97. If $0 < x < 4$, then

$$2(0) < 2(x) < 2(4)$$

$$0 < 2x < 8$$

$$0+3 < 2x+3 < 8+3$$

$$3 < 2x+3 < 11$$

So, $a = 3$ and $b = 11$.

98. If $-3 < x < 3$, then

$$-2(-3) > -2(x) > -2(3)$$

$$6 > -2x > -6$$

$$6+1 > -2x+1 > -6+1$$

$$7 > 1 - 2x > -5$$

$$-5 < 1 - 2x < 7$$

So, $a = -5$ and $b = 7$.

Chapter 1: Equations and Inequalities

- 99.** If $-3 < x < 0$, then

$$-3 + 4 < x + 4 < 0 + 4$$

$$1 < x + 4 < 4$$

$$1 > \frac{1}{x+4} > \frac{1}{4}$$

$$\frac{1}{4} < \frac{1}{x+4} < 1$$

$$\text{So, } a = \frac{1}{4} \text{ and } b = 1.$$

- 100.** If $2 < x < 4$, then

$$2 - 6 < x - 6 < 4 - 6$$

$$-4 < x - 6 < -2$$

$$-\frac{1}{4} > \frac{1}{x-6} > -\frac{1}{2}$$

$$-\frac{1}{2} < \frac{1}{x-6} < -\frac{1}{4}$$

$$\text{So, } a = -\frac{1}{2} \text{ and } b = -\frac{1}{4}.$$

- 101.** If $6 < 3x < 12$, then

$$\frac{6}{3} < \frac{3x}{3} < \frac{12}{3}$$

$$2 < x < 4$$

$$2^2 < x^2 < 4^2$$

$$4 < x^2 < 16$$

$$\text{So, } a = 4 \text{ and } b = 16.$$

- 102.** If $0 < 2x < 6$, then

$$\frac{0}{2} < \frac{2x}{2} < \frac{6}{2}$$

$$0 < x < 3$$

$$0^2 < x^2 < 3^2$$

$$0 < x^2 < 9$$

$$\text{So, } a = 0 \text{ and } b = 9.$$

- 103.** $\sqrt{3x+6}$

We need $3x+6 \geq 0$

$$3x \geq -6$$

$$x \geq -2$$

To the domain is $\{x | x \geq -2\}$ or $[-2, \infty)$.

- 104.** $\sqrt{8+2x}$

We need $8+2x \geq 0$

$$2x \geq -8$$

$$x \geq -4$$

To the domain is $\{x | x \geq -4\}$ or $[-4, \infty)$.

- 105.** $21 < \text{young adult's age} < 30$

- 106.** $40 \leq \text{middle-aged} < 60$

- 107.** **a.** Let $x = \text{age at death}$.

$$x - 30 \geq 51.9$$

$$x \geq 81.9$$

Therefore, the average life expectancy for a 30-year-old male in 2014 will be greater than or equal to 81.9 years.

- b.** Let $x = \text{age at death}$.

$$x - 30 \geq 55.6$$

$$x \geq 85.6$$

Therefore, the average life expectancy for a 30-year-old female in 2014 will be greater than or equal to 85.6 years.

- c.** By the given information, a female can expect to live $85.6 - 81.9 = 3.7$ years longer.

- 108.** $V = 20T$

$$80^\circ \leq T \leq 120^\circ$$

$$80^\circ \leq \frac{V}{20} \leq 120^\circ$$

$$1600 \leq V \leq 2400$$

The volume ranges from 1600 to 2400 cubic centimeters, inclusive.

- 109.** Let P represent the selling price and C represent the commission.

Calculating the commission:

$$C = 45,000 + 0.25(P - 900,000)$$

$$= 45,000 + 0.25P - 225,000$$

$$= 0.25P - 180,000$$

Calculate the commission range, given the price range:

$$900,000 \leq P \leq 1,100,000$$

$$0.25(900,000) \leq 0.25P \leq 0.25(1,100,000)$$

$$225,000 \leq 0.25P \leq 275,000$$

$$225,000 - 180,000 \leq 0.25P - 180,000 \leq 275,000 - 180,000$$

$$45,000 \leq C \leq 95,000$$

The agent's commission ranges from \$45,000 to \$95,000, inclusive.

$$\frac{45,000}{900,000} = 0.05 = 5\% \text{ to } \frac{95,000}{1,100,000} = 0.086 = 8.6\%,$$

inclusive.

As a percent of selling price, the commission ranges from 5% to 8.6%, inclusive.

- 110.** Let C represent the commission.

Calculate the commission range:

$$25 + 0.4(200) \leq C \leq 25 + 0.4(3000)$$

$$105 \leq C \leq 1225$$

The commissions are at least \$105 and at most \$1225.

- 111.** Let W = weekly wages and T = tax withheld. Calculating the withholding tax range, given the range of weekly wages:

$$900 \leq W \leq 1100$$

$$900 - 753 \leq W - 753 \leq 1100 - 753$$

$$147 \leq W - 753 \leq 347$$

$$0.25(147) \leq 0.25(W - 753) \leq 0.25(347)$$

$$36.75 \leq 0.25(W - 753) \leq 86.75$$

$$36.75 + 97.75 \leq 0.25(W - 753) + 97.75 \leq 86.75 + 97.75$$

$$134.50 \leq T \leq 184.50$$

The amount withheld varies from \$134.50 to \$184.50, inclusive.

- 112.** Let x represent the length of time Sue should exercise on the seventh day.

$$200 \leq 40 + 45 + 0 + 50 + 25 + 35 + x \leq 300$$

$$200 \leq 195 + x \leq 300$$

$$5 \leq x \leq 105$$

Sue will stay within the ACSM guidelines by exercising from 5 to 105 minutes.

- 113.** Let K represent the monthly usage in kilowatt-hours and let C represent the monthly customer bill.

Calculating the bill: $C = 0.0821K + 15.37$

Calculating the range of kilowatt-hours, given the range of bills:

$$72.84 \leq C \leq 237.04$$

$$72.84 \leq 0.0821K + 15.37 \leq 237.04$$

$$57.47 \leq 0.0821K \leq 221.67$$

$$700.00 \leq K \leq 2700.00$$

The usage varies from 700.00 kilowatt-hours to 2700.00 kilowatt-hours, inclusive.

- 114.** Let W represent the amount of water used (in thousands of gallons). Let C represent the customer charge (in dollars).

Calculating the charge:

$$C = 57.07 + 5.81(W - 10)$$

$$= 57.07 + 5.81W - 58.10$$

$$= 5.81W - 1.03$$

Calculating the range of water usage, given the range of charges:

$$97.74 \leq C \leq 150.03$$

$$97.74 \leq 5.81W - 1.03 \leq 150.03$$

$$98.77 \leq 5.81W \leq 151.06$$

$$17 \leq W \leq 26$$

The range of water usage ranged from 17,000 to 26,000 gallons.

- 115.** Let C represent the dealer's cost and M represent the markup over dealer's cost. If the price is \$18,000, then

$$18,000 = C + MC = C(1 + M)$$

$$\text{Solving for } C \text{ yields: } C = \frac{18,000}{1 + M}$$

Calculating the range of dealer costs, given the range of markups:

$$0.12 \leq M \leq 0.18$$

$$1.12 \leq 1 + M \leq 1.18$$

$$\frac{1}{1.12} \geq \frac{1}{1 + M} \geq \frac{1}{1.18}$$

$$\frac{18,000}{1.12} \geq \frac{18,000}{1 + M} \geq \frac{18,000}{1.18}$$

$$16,071.43 \geq C \geq 15,254.24$$

The dealer's cost varies from \$15,254.24 to \$16,071.43, inclusive.

- 116.** Let T represent the test scores of the people in the top 2.5%.

$$T > 1.96(12) + 100 = 123.52$$

People in the top 2.5% will have test scores greater than 123.52. That is, $T > 123.52$ or $(123.52, \infty)$.

- 117. a.** Let T represent the score on the last test and G represent the course grade.

Calculating the course grade and solving for the last test:

$$G = \frac{68 + 82 + 87 + 89 + T}{5}$$

$$G = \frac{326 + T}{5}$$

$$5G = 326 + T$$

$$T = 5G - 326$$

Calculating the range of scores on the last test, given the grade range:

$$80 \leq G < 90$$

$$400 \leq 5G < 450$$

$$74 \leq 5G - 326 < 124$$

$$74 \leq T < 124$$

To get a grade of B, you need at least a 74 on the fifth test.

Chapter 1: Equations and Inequalities

- b. Let T represent the score on the last test and G represent the course grade.

Calculating the course grade and solving for the last test:

$$G = \frac{68+82+87+89+2T}{6}$$

$$G = \frac{326+2T}{6}$$

$$G = \frac{163+T}{3}$$

$$T = 3G - 163$$

Calculating the range of scores on the last test, given the grade range:

$$80 \leq G < 90$$

$$240 \leq 3G < 270$$

$$77 \leq 3G - 163 < 107$$

$$77 \leq T < 107$$

To get a grade of B, you need at least a 77 on the fifth test.

118. Let C represent the number of calories in a serving of regular Miracle Whip®, and let F represent the grams of fat in a serving of regular Miracle Whip®.

One possibility for a “light” classification is that the 20 calories in a serving of Miracle Whip® Light is less than or equal to one-third the calories in regular Miracle Whip®. That is,

$$20 \leq \frac{1}{3}C.$$

The second possibility for a “light” classification is that the 1.5 grams of fat in a serving of Miracle Whip® Light is less than or equal to one-half the grams of fat in regular Miracle Whip®.

$$\text{That is, } 1.5 \leq \frac{1}{2}F.$$

We have:

$$20 \leq \frac{1}{3}C \quad \text{or} \quad 1.5 \leq \frac{1}{2}F$$

$$60 \leq C \quad \text{or} \quad 3 \leq F$$

A serving of regular Miracle Whip® either contains at least 60 calories or at least 3 grams of fat, or both.

119. Since $a < b$,

$$\frac{a}{2} < \frac{b}{2} \quad \text{and} \quad \frac{a}{2} < \frac{b}{2}$$

$$\frac{a}{2} + \frac{a}{2} < \frac{b}{2} + \frac{b}{2} \quad \text{and} \quad \frac{a}{2} + \frac{b}{2} < \frac{b}{2} + \frac{b}{2}$$

$$a < \frac{a+b}{2} \quad \text{and} \quad \frac{a+b}{2} < b$$

$$\text{Thus, } a < \frac{a+b}{2} < b.$$

120. From problem 115, $a < \frac{a+b}{2} < b$, so

$$d\left(a, \frac{a+b}{2}\right) = \frac{a+b}{2} - a = \frac{a+b-2a}{2} = \frac{b-a}{2} \quad \text{and}$$

$$d\left(b, \frac{a+b}{2}\right) = b - \frac{a+b}{2} = \frac{2b-a-b}{2} = \frac{b-a}{2}.$$

Therefore, $\frac{a+b}{2}$ is equidistant from a and b .

121. If $0 < a < b$, then

$$ab > a^2 > 0 \quad \text{and} \quad b^2 > ab > 0$$

$$(\sqrt{ab})^2 > a^2 \quad \text{and} \quad b^2 > (\sqrt{ab})^2$$

$$\sqrt{ab} > a \quad \text{and} \quad b > \sqrt{ab}$$

$$\text{Thus, } a < \sqrt{ab} < b.$$

122. Show that $\sqrt{ab} < \frac{a+b}{2}$.

$$\frac{a+b}{2} - \sqrt{ab} = \frac{1}{2}(a - 2\sqrt{ab} + b)$$

$$= \frac{1}{2}(\sqrt{a} - \sqrt{b})^2 > 0, \text{ since } a \neq b.$$

$$\text{Therefore, } \sqrt{ab} < \frac{a+b}{2}.$$

123. For $0 < a < b$, $\frac{1}{h} = \frac{1}{2}\left(\frac{1}{a} + \frac{1}{b}\right)$

$$h \cdot \frac{1}{h} = \frac{1}{2}\left(\frac{b+a}{ab}\right) \cdot h$$

$$1 = \frac{1}{2}\left(\frac{b+a}{ab}\right) \cdot h$$

$$h = \frac{2ab}{a+b}$$

$$\begin{aligned} h-a &= \frac{2ab}{a+b} - a = \frac{2ab - a(a+b)}{a+b} \\ &= \frac{2ab - a^2 - ab}{a+b} = \frac{ab - a^2}{a+b} \\ &= \frac{a(b-a)}{a+b} > 0 \end{aligned}$$

Therefore, $h > a$.

$$\begin{aligned} b-h &= b - \frac{2ab}{a+b} = \frac{b(a+b) - 2ab}{a+b} \\ &= \frac{ab + b^2 - 2ab}{a+b} = \frac{b^2 - ab}{a+b} \\ &= \frac{b(b-a)}{a+b} > 0 \end{aligned}$$

Therefore, $h < b$, and we have $a < h < b$.

124. Show that $h = \frac{(\text{geometric mean})^2}{\text{arithmetic mean}} = \frac{(\sqrt{ab})^2}{\left(\frac{1}{2}(a+b)\right)}$

From Problem 119, we know:

$$\begin{aligned} \frac{1}{h} &= \frac{1}{2} \left(\frac{1}{a} + \frac{1}{b} \right) \\ \frac{2}{h} &= \frac{1}{a} + \frac{1}{b} = \frac{b+a}{ab} \\ \frac{h}{2} &= \frac{ab}{a+b} \\ h &= 2 \cdot \frac{ab}{a+b} = \frac{(\sqrt{ab})^2}{\left(\frac{1}{2}(a+b)\right)} \end{aligned}$$

125. Since $0 < a < b$, then $a - b < 0$ and $ab > 0$.

Therefore, $\frac{a-b}{ab} < 0$. So,

$$\begin{aligned} \frac{a}{ab} - \frac{b}{ab} &< 0 \\ \frac{1}{b} - \frac{1}{a} &< 0 \\ \frac{1}{b} &< \frac{1}{a} \end{aligned}$$

Now, since $b > 0$, then $\frac{1}{b} > 0$, so we have

$$0 < \frac{1}{b} < \frac{1}{a}.$$

126. Answers will vary. One possibility:

No solution: $4x+6 \leq 2(x-5)+2x$

One solution: $3x+5 \leq 2(x+3)+1 \leq 3(x+2)-1$

127. Since $x^2 \geq 0$, we have

$$x^2 + 1 \geq 0 + 1$$

$$x^2 + 1 \geq 1$$

Therefore, the expression $x^2 + 1$ can never be less than -5 .

- 128 – 129. Answers will vary.

Section 1.6

1. $| -2 | = 2$

2. True

3. $\{ -5, 5 \}$

4. $\{ x \mid -5 < x < 5 \}$

5. True

6. True

7. d

8. a

9. $| 2x | = 6$

$2x = 6$ or $2x = -6$

$x = 3$ or $x = -3$

The solution set is $\{ -3, 3 \}$.

10. $| 3x | = 12$

$3x = 12$ or $3x = -12$

$x = 4$ or $x = -4$

The solution set is $\{ -4, 4 \}$.

11. $| 2x+3 | = 5$

$2x+3 = 5$ or $2x+3 = -5$

$2x = 2$ or $2x = -8$

$x = 1$ or $x = -4$

The solution set is $\{ -4, 1 \}$.

Chapter 1: Equations and Inequalities

12. $|3x - 1| = 2$

$$3x - 1 = 2 \text{ or } 3x - 1 = -2$$

$$3x = 3 \text{ or } 3x = -1$$

$$x = 1 \text{ or } x = -\frac{1}{3}$$

The solution set is $\left\{-\frac{1}{3}, 1\right\}$.

13. $|1 - 4t| + 8 = 13$

$$|1 - 4t| = 5$$

$$1 - 4t = 5 \text{ or } 1 - 4t = -5$$

$$-4t = 4 \text{ or } -4t = -6$$

$$t = -1 \text{ or } t = \frac{3}{2}$$

The solution set is $\left\{-1, \frac{3}{2}\right\}$.

14. $|1 - 2z| + 6 = 9$

$$|1 - 2z| = 3$$

$$1 - 2z = 3 \text{ or } 1 - 2z = -3$$

$$-2z = 2 \text{ or } -2z = -4$$

$$z = -1 \text{ or } z = 2$$

The solution set is $\{-1, 2\}$.

15. $|-2x| = |8|$

$$|-2x| = 8$$

$$-2x = 8 \text{ or } -2x = -8$$

$$x = -4 \text{ or } x = 4$$

The solution set is $\{-4, 4\}$.

16. $|-x| = |1|$

$$|-x| = 1$$

$$-x = 1 \text{ or } -x = -1$$

The solution set is $\{-1, 1\}$.

17. $|-2|x| = 4$

$$2x = 4$$

$$x = 2$$

The solution set is $\{2\}$.

18. $|3|x| = 9$

$$3x = 9$$

$$x = 3$$

The solution set is $\{3\}$.

19. $\frac{2}{3}|x| = 9$

$$|x| = \frac{27}{2}$$

$$x = \frac{27}{2} \text{ or } x = -\frac{27}{2}$$

The solution set is $\left\{-\frac{27}{2}, \frac{27}{2}\right\}$.

20. $\frac{3}{4}|x| = 9$

$$|x| = 12$$

$$x = 12 \text{ or } x = -12$$

The solution set is $\{-12, 12\}$.

21. $\left|\frac{x}{3} + \frac{2}{5}\right| = 2$

$$\frac{x}{3} + \frac{2}{5} = 2 \text{ or } \frac{x}{3} + \frac{2}{5} = -2$$

$$5x + 6 = 30 \text{ or } 5x + 6 = -30$$

$$5x = 24 \text{ or } 5x = -36$$

$$x = \frac{24}{5} \text{ or } x = -\frac{36}{5}$$

The solution set is $\left\{-\frac{36}{5}, \frac{24}{5}\right\}$.

22. $\left|\frac{x}{2} - \frac{1}{3}\right| = 1$

$$\frac{x}{2} - \frac{1}{3} = 1 \text{ or } \frac{x}{2} - \frac{1}{3} = -1$$

$$3x - 2 = 6 \text{ or } 3x - 2 = -6$$

$$3x = 8 \text{ or } 3x = -4$$

$$x = \frac{8}{3} \text{ or } x = -\frac{4}{3}$$

The solution set is $\left\{-\frac{4}{3}, \frac{8}{3}\right\}$.

23. $|u - 2| = -\frac{1}{2}$

No solution, since absolute value always yields a non-negative number.

24. $|2 - v| = -1$

No solution, since absolute value always yields a non-negative number.

25. $4 - |2x| = 3$

$$-|2x| = -1$$

$$|2x| = 1$$

$$2x = 1 \text{ or } 2x = -1$$

$$x = \frac{1}{2} \text{ or } x = -\frac{1}{2}$$

The solution set is $\left\{-\frac{1}{2}, \frac{1}{2}\right\}$.

26. $5 - \left|\frac{1}{2}x\right| = 3$

$$-\left|\frac{1}{2}x\right| = -2$$

$$\left|\frac{1}{2}x\right| = 2$$

$$\frac{1}{2}x = 2 \text{ or } \frac{1}{2}x = -2$$

$$x = 4 \text{ or } x = -4$$

The solution set is $\{-4, 4\}$.

27. $|x^2 - 9| = 0$

$$x^2 - 9 = 0$$

$$x^2 = 9$$

$$x = \pm 3$$

The solution set is $\{-3, 3\}$.

28. $|x^2 - 16| = 0$

$$x^2 - 16 = 0$$

$$x^2 = 16$$

$$x = \pm 4$$

The solution set is $\{-4, 4\}$.

29. $|x^2 - 2x| = 3$

$$x^2 - 2x = 3 \quad \text{or} \quad x^2 - 2x = -3$$

$$x^2 - 2x - 3 = 0 \quad \text{or} \quad x^2 - 2x + 3 = 0$$

$$(x-3)(x+1) = 0 \quad \text{or} \quad x = \frac{2 \pm \sqrt{4-12}}{2}$$

$$x = 3 \text{ or } x = -1 \text{ or } x = \frac{2 \pm \sqrt{-8}}{2} \text{ no real sol.}$$

The solution set is $\{-1, 3\}$.

30. $|x^2 + x| = 12$

$$x^2 + x = 12 \quad \text{or} \quad x^2 + x = -12$$

$$x^2 + x - 12 = 0 \quad \text{or} \quad x^2 + x + 12 = 0$$

$$(x-3)(x+4) = 0 \quad \text{or} \quad x = \frac{-1 \pm \sqrt{1-48}}{2}$$

$$x = 3 \text{ or } x = -4 \text{ or } x = \frac{1 \pm \sqrt{-47}}{2} \text{ no real sol.}$$

The solution set is $\{-4, 3\}$.

31. $|x^2 + x - 1| = 1$

$$x^2 + x - 1 = 1 \quad \text{or} \quad x^2 + x - 1 = -1$$

$$x^2 + x - 2 = 0 \quad \text{or} \quad x^2 + x = 0$$

$$(x-1)(x+2) = 0 \quad \text{or} \quad x(x+1) = 0$$

$$x = 1, x = -2 \quad \text{or} \quad x = 0, x = -1$$

The solution set is $\{-2, -1, 0, 1\}$.

32. $|x^2 + 3x - 2| = 2$

$$x^2 + 3x - 2 = 2 \quad \text{or} \quad x^2 + 3x - 2 = -2$$

$$x^2 + 3x = 4 \quad \text{or} \quad x^2 + 3x = 0$$

$$x^2 + 3x - 4 = 0 \quad \text{or} \quad x(x+3) = 0$$

$$(x+4)(x-1) = 0 \quad \text{or} \quad x = 0, x = -3$$

$$x = -4, x = 1$$

The solution set is $\{-4, -3, 0, 1\}$.

33. $\left|\frac{3x-2}{2x-3}\right| = 2$

$$\frac{3x-2}{2x-3} = 2 \quad \text{or} \quad \frac{3x-2}{2x-3} = -2$$

$$3x-2 = 2(2x-3) \quad \text{or} \quad 3x-2 = -2(2x-3)$$

$$3x-2 = 4x-6 \quad \text{or} \quad 3x-2 = -4x+6$$

$$-x = -4 \quad \text{or} \quad 7x = 8$$

$$x = 4 \quad \text{or} \quad x = \frac{8}{7}$$

Neither of these values cause the denominator to equal zero, so the solution set is $\left\{\frac{8}{7}, 4\right\}$.

Chapter 1: Equations and Inequalities

34. $\left| \frac{2x+1}{3x+4} \right| = 1$

$$\frac{2x+1}{3x+4} = 1 \quad \text{or} \quad \frac{2x+1}{3x+4} = -1$$

$$2x+1 = 1(3x+4) \quad \text{or} \quad 2x+1 = -1(3x+4)$$

$$2x+1 = 3x+4 \quad \text{or} \quad 2x+1 = -3x-4$$

$$-x = 3 \quad \text{or} \quad 5x = -5$$

$$x = -3 \quad \text{or} \quad x = -1$$

Neither of these values cause the denominator to equal zero, so the solution set is $\{-3, -1\}$.

35. $|x^2 + 3x| = |x^2 - 2x|$

$$x^2 + 3x = x^2 - 2x \quad \text{or} \quad x^2 + 3x = -(x^2 - 2x)$$

$$3x = -2x \quad \text{or} \quad x^2 + 3x = -x^2 + 2x$$

$$5x = 0 \quad \text{or} \quad 2x^2 + x = 0$$

$$x = 0 \quad \text{or} \quad x(2x+1) = 0$$

$$x = 0 \quad \text{or} \quad x = 0 \quad \text{or} \quad x = -\frac{1}{2}$$

The solution set is $\left\{-\frac{1}{2}, 0\right\}$.

36. $|x^2 - 2x| = |x^2 + 6x|$

$$x^2 - 2x = x^2 + 6x \quad \text{or} \quad x^2 - 2x = -(x^2 + 6x)$$

$$-2x = 6x \quad \text{or} \quad x^2 - 2x = -x^2 - 6x$$

$$-8x = 0 \quad \text{or} \quad 2x^2 + 4x = 0$$

$$x = 0 \quad \text{or} \quad 2x(x+2) = 0$$

$$x = 0 \quad \text{or} \quad x = 0 \text{ or } x = -2$$


The solution set is $\{-2, 0\}$.

37. $|2x| < 8$

$$-8 < 2x < 8$$

$$-4 < x < 4$$

$$\{x \mid -4 < x < 4\} \text{ or } (-4, 4)$$




38. $|3x| < 15$

$$-15 < 3x < 15$$

$$-5 < x < 5$$

$$\{x \mid -5 < x < 5\} \text{ or } (-5, 5)$$




39. $|3x| > 12$

$$3x < -12 \quad \text{or} \quad 3x > 12$$

$$x < -4 \quad \text{or} \quad x > 4$$

$$\{x \mid x < -4 \text{ or } x > 4\} \text{ or } (-\infty, -4) \cup (4, \infty)$$




40. $|2x| > 6$

$$2x < -6 \quad \text{or} \quad 2x > 6$$

$$x < -3 \quad \text{or} \quad x > 3$$

$$\{x \mid x < -3 \text{ or } x > 3\} \text{ or } (-\infty, -3) \cup (3, \infty)$$




41. $|x-2| + 2 < 3$

$$|x-2| < 1$$

$$-1 < x-2 < 1$$

$$1 < x < 3$$

$$\{x \mid 1 < x < 3\} \text{ or } (1, 3)$$




42. $|x+4| + 3 < 5$

$$|x+4| < 2$$

$$-2 < x+4 < 2$$

$$-6 < x < -2$$

$$\{x \mid -6 < x < -2\} \text{ or } (-6, -2)$$




43. $|3t-2| \leq 4$

$$-4 \leq 3t-2 \leq 4$$

$$-2 \leq 3t \leq 6$$

$$-\frac{2}{3} \leq t \leq 2$$

$$\left\{t \mid -\frac{2}{3} \leq t \leq 2\right\} \text{ or } \left[-\frac{2}{3}, 2\right]$$



44. $|2u+5| \leq 7$

$$-7 \leq 2u+5 \leq 7$$

$$-12 \leq 2u \leq 2$$

$$-6 \leq u \leq 1$$

$$\{u | -6 \leq u \leq 1\} \text{ or } [-6, 1]$$




44. $|2x-3| \geq 2$

$$2x-3 \leq -2 \text{ or } 2x-3 \geq 2$$

$$2x \leq 1 \text{ or } 2x \geq 5$$

$$x \leq \frac{1}{2} \text{ or } x \geq \frac{5}{2}$$

$$\left\{x \mid x \leq \frac{1}{2} \text{ or } x \geq \frac{5}{2}\right\} \text{ or } \left(-\infty, \frac{1}{2}\right] \cup \left[\frac{5}{2}, \infty\right)$$




46. $|3x+4| \geq 2$

$$3x+4 \leq -2 \text{ or } 3x+4 \geq 2$$

$$3x \leq -6 \text{ or } 3x \geq -2$$

$$x \leq -2 \text{ or } x \geq -\frac{2}{3}$$

$$\left\{x \mid x \leq -2 \text{ or } x \geq -\frac{2}{3}\right\} \text{ or } (-\infty, -2] \cup \left[-\frac{2}{3}, \infty\right)$$



47. $|1-4x|-7 < -2$

$$|1-4x| < 5$$


$$-5 < 1-4x < 5$$

$$-6 < -4x < 4$$

$$\frac{-6}{-4} > x > \frac{4}{-4}$$

$$\frac{3}{2} > x > -1 \text{ or } -1 < x < \frac{3}{2}$$

$$\left\{x \mid -1 < x < \frac{3}{2}\right\} \text{ or } \left(-1, \frac{3}{2}\right)$$



48. $|1-2x|-4 < -1$

$$|1-2x| < 3$$

$$-3 < 1-2x < 3$$

$$-4 < -2x < 2$$

$$\frac{-4}{-2} > x > \frac{2}{-2}$$

$$2 > x > -1 \text{ or } -1 < x < 2$$

$$\{x | -1 < x < 2\} \text{ or } (-1, 2)$$




49. $|1-2x| > 3$

$$1-2x < -3 \text{ or } 1-2x > 3$$

$$-2x < -4 \text{ or } -2x > 2$$

$$x > 2 \text{ or } x < -1$$

$$\{x | x < -1 \text{ or } x > 2\} \text{ or } (-\infty, -1) \cup (2, \infty)$$




50. $|2-3x| > 1$

$$2-3x < -1 \text{ or } 2-3x > 1$$

$$-3x < -3 \text{ or } -3x > -1$$

$$x > 1 \text{ or } x < \frac{1}{3}$$

$$\left\{x \mid x < \frac{1}{3} \text{ or } x > 1\right\} \text{ or } \left(-\infty, \frac{1}{3}\right) \cup (1, \infty)$$




51. $|-4x| + |-5| \leq 1$

$$|-4x| + 5 \leq 1$$

$$|-4x| \leq -4$$

This is impossible since absolute value always yields a non-negative number. The inequality has no solution.



52. $|-x| - 4 \leq 2$

$$|-x| - 4 \leq 2$$


$$|-x| \leq 6$$

$$-6 \leq -x \leq 6$$

$$6 \geq x \geq -6$$

Chapter 1: Equations and Inequalities

$\{x \mid -6 \leq x \leq 6\}$ or $[-6, 6]$




53. $| -2x | > | -3 |$

$$| 2x | > 3$$

$$2x < -3 \text{ or } 2x > 3$$

$$x < -\frac{3}{2} \text{ or } x > \frac{3}{2}$$

$$\left\{ x \mid x < -\frac{3}{2} \text{ or } x > \frac{3}{2} \right\} \text{ or } \left(-\infty, -\frac{3}{2} \right) \cup \left(\frac{3}{2}, \infty \right)$$




54. $| -x - 2 | \geq 1$

$$-x - 2 \leq -1 \text{ or } -x - 2 \geq 1$$

$$-x \leq 1 \text{ or } -x \geq 3$$

$$x \geq -1 \text{ or } x \leq -3$$

$$\{x \mid x \leq -3 \text{ or } x \geq -1\} \text{ or } (-\infty, -3] \cup [-1, \infty)$$



55. $-| 2x - 1 | \geq -3$


$$| 2x - 1 | \leq 3$$

$$-3 \leq 2x - 1 \leq 3$$

$$-2 \leq 2x \leq 4$$

$$-1 \leq x \leq 2$$

$$\{x \mid -1 \leq x \leq 2\} \text{ or } [-1, 2]$$



56. $-| 1 - 2x | \geq -3$


$$| 1 - 2x | \leq 3$$

$$-3 \leq 1 - 2x \leq 3$$

$$-4 \leq -2x \leq 2$$


$$2 \geq x \geq -1$$

$$\{x \mid -1 \leq x \leq 2\} \text{ or } [-1, 2]$$



57. $| 2x | < -1$

This is impossible since absolute value always yields a non-negative number. No solution.



58. $| 3x | \geq 0$

Absolute value yields a non-negative number, so this inequality is true for all real numbers, $(-\infty, \infty)$.




59. $| 5x | \geq -1$

Absolute value yields a non-negative number, so this inequality is true for all real numbers, $(-\infty, \infty)$.



60. $| 6x | < -2$

This is impossible since absolute value always yields a non-negative number. No solution.



61. $\left| \frac{2x+3}{3} - \frac{1}{2} \right| < 1$

$$-1 < \frac{2x+3}{3} - \frac{1}{2} < 1$$

$$6(-1) < 6\left(\frac{2x+3}{3} - \frac{1}{2}\right) < 6(1)$$

$$-6 < 2(2x+3) - 3 < 6$$


$$-6 < 4x + 6 - 3 < 6$$

$$-6 < 4x + 3 < 6$$

$$-9 < 4x < 3$$

$$-\frac{9}{4} < x < \frac{3}{4}$$

$$\left\{ x \mid -\frac{9}{4} < x < \frac{3}{4} \right\} \text{ or } \left(-\frac{9}{4}, \frac{3}{4} \right)$$



62. $3 - | x + 1 | < \frac{1}{2}$


$$-| x + 1 | < -\frac{5}{2}$$

$$| x + 1 | > \frac{5}{2}$$

$$x + 1 < -\frac{5}{2} \text{ or } x + 1 > \frac{5}{2}$$

$$x < -\frac{7}{2} \text{ or } x > \frac{3}{2}$$

$$\left\{ x \mid x < -\frac{7}{2} \text{ or } x > \frac{3}{2} \right\} \text{ or } \left(-\infty, -\frac{7}{2} \right) \cup \left(\frac{3}{2}, \infty \right)$$



63. $5 + |x - 1| > \frac{1}{2}$
 $|x - 1| > -\frac{9}{2}$


Absolute value yields a non-negative number, so this inequality is true for all real numbers, $(-\infty, \infty)$.



64. $\left| \frac{2x-3}{2} + \frac{1}{3} \right| > 1$

$$\begin{aligned} \frac{2x-3}{2} + \frac{1}{3} &< -1 & \text{or} & \quad \frac{2x-3}{2} + \frac{1}{3} > 1 \\ 6\left(\frac{2x-3}{2} + \frac{1}{3}\right) &< 6(-1) & \text{or} & \quad 6\left(\frac{2x-3}{2} + \frac{1}{3}\right) > 6(1) \\ 3(2x-3) + 2 &< -6 & \text{or} & \quad 3(2x-3) + 2 > 6 \\ 6x - 9 + 2 &< -6 & \text{or} & \quad 6x - 9 + 2 > 6 \\ 6x - 7 &< -6 & \text{or} & \quad 6x - 7 > 6 \\ 6x &< 1 & \text{or} & \quad 6x > 13 \\ x &< \frac{1}{6} & \text{or} & \quad x > \frac{13}{6} \end{aligned}$$

$$\left\{ x \mid x < \frac{1}{6} \text{ or } x > \frac{13}{6} \right\} \text{ or } \left(-\infty, \frac{1}{6} \right) \cup \left(\frac{13}{6}, \infty \right)$$



65. A temperature x that differs from 98.6°F by at least 1.5°F .

$$\begin{aligned} |x - 98.6^\circ| &\geq 1.5^\circ \\ x - 98.6^\circ &\leq -1.5^\circ \text{ or } x - 98.6^\circ \geq 1.5^\circ \\ x &\leq 97.1^\circ \text{ or } x \geq 100.1^\circ \end{aligned}$$

The temperatures that are considered unhealthy are those that are less than 97.1°F or greater than 100.1°F , inclusive.

66. A voltage x that differs from 110 volts by at most 5 volts.

$$\begin{aligned} |x - 110| &\leq 5 \\ -5 &\leq x - 110 \leq 5 \end{aligned}$$

$$105 \leq x \leq 115$$

The actual voltage is between 105 and 115 volts, inclusive.

67. The true average number of books read x should differ from 13.6 by less than 1.8 books.

$$\begin{aligned} |x - 13.6| &< 1.85 \\ -1.85 &< x - 13.6 < 1.85 \\ 11.8 &< x < 15.4 \end{aligned}$$

Gallup is 99% confident that the actual average number of books read per year is between 11.8 and 15.4 books.

68. The speed x varies from 707 mph by up to 55 mph.

a. $|x - 707| \leq 55$
 b. $-55 \leq x - 707 \leq 55$
 $-55 \leq x - 707 \leq 55$
 $652 \leq x \leq 762$

The speed of sound is between 652 and 762 miles per hour, depending on conditions.

69. x differs from 3 by less than $\frac{1}{2}$.

$$\begin{aligned} |x - 3| &< \frac{1}{2} \\ -\frac{1}{2} &< x - 3 < \frac{1}{2} \\ \frac{5}{2} &< x < \frac{7}{2} \\ \left\{ x \mid \frac{5}{2} < x < \frac{7}{2} \right\} \end{aligned}$$

70. x differs from -4 by less than 1

$$\begin{aligned} |x - (-4)| &< 1 \\ |x + 4| &< 1 \\ -1 &< x + 4 < 1 \\ -5 &< x < -3 \\ \{x \mid -5 < x < -3\} \end{aligned}$$

71. x differs from -3 by more than 2.

$$\begin{aligned} |x - (-3)| &> 2 \\ |x + 3| &> 2 \\ x + 3 &< -2 \text{ or } x + 3 > 2 \\ x &< -5 \text{ or } x > -1 \\ \{x \mid x < -5 \text{ or } x > -1\} \end{aligned}$$

72. x differs from 2 by more than 3.

$$\begin{aligned} |x - 2| &> 3 \\ x - 2 &< -3 \text{ or } x - 2 > 3 \\ x &< -1 \text{ or } x > 5 \\ \{x \mid x < -1 \text{ or } x > 5\} \end{aligned}$$

Chapter 1: Equations and Inequalities

73. $|x-1| < 3$

$$-3 < x-1 < 3$$

$$-3+5 < (x-1)+5 < 3+5$$

$$2 < x+4 < 8$$

$$a = 2, b = 8$$

74. $|x+2| < 5$

$$-5 < x+2 < 5$$

$$-5-4 < (x+2)-4 < 5-4$$

$$-9 < x-2 < 1$$

$$a = -9, b = 1$$

75. $|x+4| \leq 2$

$$-2 \leq x+4 \leq 2$$

$$-6 \leq x \leq -2$$

$$-12 \leq 2x \leq -4$$

$$-15 \leq 2x-3 \leq -7$$

$$a = -15, b = -7$$

76. $|x-3| \leq 1$

$$-1 \leq x-3 \leq 1$$

$$2 \leq x \leq 4$$

$$6 \leq 3x \leq 12$$

$$7 \leq 3x+1 \leq 13$$

$$a = 7, b = 13$$

77. $|x-2| \leq 7$

$$-7 \leq x-2 \leq 7$$

$$-5 \leq x \leq 9$$

$$-15 \leq x-10 \leq -1$$

$$-\frac{1}{15} \geq \frac{1}{x-10} \geq -1$$

$$-1 \leq \frac{1}{x-10} \leq -\frac{1}{15}$$

$$a = -1, b = -\frac{1}{15}$$

78. $|x+1| \leq 3$

$$-3 \leq x+1 \leq 3$$

$$-4 \leq x \leq 2$$

$$1 \leq x+5 \leq 7$$

$$1 \geq \frac{1}{x+5} \geq \frac{1}{7}$$

$$\frac{1}{7} \leq \frac{1}{x+5} \leq 1$$

$$a = \frac{1}{7}, b = 1$$

79. Given that $a > 0, b > 0$, and $\sqrt{a} < \sqrt{b}$, show that $a < b$.

$$\text{Note that } b-a = (\sqrt{b} + \sqrt{a})(\sqrt{b} - \sqrt{a}).$$

Since $\sqrt{a} < \sqrt{b}$ means $\sqrt{b} - \sqrt{a} > 0$, we have

$$b-a = (\sqrt{b} + \sqrt{a})(\sqrt{b} - \sqrt{a}) > 0.$$

Therefore, $b-a > 0$ which means $a < b$.

80. Show that $a \leq |a|$.

We know $0 \leq |a|$. So if $a < 0$, then we have

$a < 0 \leq |a|$ which means $a \leq |a|$. Now, if

$a \geq 0$, then $|a| = a$. So $a \leq |a|$.

81. Prove $|a+b| \leq |a| + |b|$.

$$\text{Note that } |a+b|^2 = |a+b| \cdot |a+b|.$$

Case 1: If $a+b \geq 0$, then $|a+b| = a+b$, so

$$\begin{aligned} |a+b| \cdot |a+b| &= (a+b)(a+b) \\ &= a^2 + 2ab + b^2 \\ &\leq |a|^2 + 2|a| \cdot |b| + |b|^2 \\ &= (|a| + |b|)^2 \quad \text{by problem 78} \end{aligned}$$

$$\text{Thus, } (|a+b|)^2 \leq (|a| + |b|)^2$$

$$|a+b| \leq |a| + |b|.$$

Case 2: If $a+b < 0$, then $|a+b| = -(a+b)$, so

$$\begin{aligned} |a+b| \cdot |a+b| &= (-(a+b))(-(a+b)) \\ &= (a+b)(a+b) \\ &= a^2 + 2ab + b^2 \\ &\leq |a|^2 + 2|a| \cdot |b| + |b|^2 \\ &= (|a| + |b|)^2 \quad \text{by problem 78} \end{aligned}$$

$$\text{Thus, } (|a+b|)^2 \leq (|a| + |b|)^2$$

$$|a+b| \leq |a| + |b|$$

82. Prove $|a-b| \geq |a| - |b|$.

$$|a| = |(a-b) + b| \leq |a-b| + |b| \quad \text{by the Triangle}$$

Inequality, so $|a| \leq |a-b| + |b|$ which means

$$|a| - |b| \leq |a-b|. \quad \text{Therefore, } |a-b| \geq |a| - |b|.$$

83. Given that $a > 0$,

$$\begin{aligned}x^2 &< a \\x^2 - a &< 0 \\(x + \sqrt{a})(x - \sqrt{a}) &< 0\end{aligned}$$

If $x < -\sqrt{a}$, then $x + \sqrt{a} < 0$ and $x - \sqrt{a} < -2\sqrt{a} < 0$. Therefore, $(x + \sqrt{a})(x - \sqrt{a}) > 0$, which is a contradiction.

If $-\sqrt{a} < x < \sqrt{a}$, then $0 < x + \sqrt{a} < 2\sqrt{a}$ and $-2\sqrt{a} < x - \sqrt{a} < 0$.

Therefore, $(x + \sqrt{a})(x - \sqrt{a}) < 0$.

If $x > \sqrt{a}$, then $x + \sqrt{a} > 2\sqrt{a} > 0$ and $x - \sqrt{a} > 0$. Therefore, $(x + \sqrt{a})(x - \sqrt{a}) > 0$, which is a contradiction. So the solution set for $x^2 < a$ is $\{x | -\sqrt{a} < x < \sqrt{a}\}$.

84. Given that $a > 0$,

$$\begin{aligned}x^2 &> a \\x^2 - a &> 0 \\(x + \sqrt{a})(x - \sqrt{a}) &> 0\end{aligned}$$

If $x < -\sqrt{a}$, then $x + \sqrt{a} < 0$ and $x - \sqrt{a} < -2\sqrt{a} < 0$. Therefore, $(x + \sqrt{a})(x - \sqrt{a}) > 0$.

If $-\sqrt{a} < x < \sqrt{a}$, then $0 < x + \sqrt{a} < 2\sqrt{a}$ and $-2\sqrt{a} < x - \sqrt{a} < 0$. Therefore, $(x + \sqrt{a})(x - \sqrt{a}) < 0$, which is a contradiction.

If $x > \sqrt{a}$, then $x + \sqrt{a} > 2\sqrt{a} > 0$ and $x - \sqrt{a} > 0$. Therefore, $(x + \sqrt{a})(x - \sqrt{a}) > 0$.

So the solution set for $x^2 > a$ is $\{x | x < -\sqrt{a} \text{ or } x > \sqrt{a}\}$.

85. $x^2 < 1$

$$\begin{aligned}-\sqrt{1} &< x < \sqrt{1} \\-1 &< x < 1\end{aligned}$$

The solution set is $\{x | -1 < x < 1\}$.

86. $x^2 < 4$

$$\begin{aligned}-\sqrt{4} &< x < \sqrt{4} \\-2 &< x < 2\end{aligned}$$

The solution set is $\{x | -2 < x < 2\}$.

87. $x^2 \geq 9$

$$\begin{aligned}x \leq -\sqrt{9} \text{ or } x &\geq \sqrt{9} \\x \leq -3 \text{ or } x &\geq 3\end{aligned}$$

The solution set is $\{x | x \leq -3 \text{ or } x \geq 3\}$.

88. $x^2 \geq 1$

$$\begin{aligned}x \leq -\sqrt{1} \text{ or } x &\geq \sqrt{1} \\x \leq -1 \text{ or } x &\geq 1\end{aligned}$$

The solution set is $\{x | x \leq -1 \text{ or } x \geq 1\}$.

89. $x^2 \leq 16$

$$\begin{aligned}-\sqrt{16} \leq x &\leq \sqrt{16} \\-4 \leq x &\leq 4\end{aligned}$$

The solution set is $\{x | -4 \leq x \leq 4\}$.

90. $x^2 \leq 9$

$$\begin{aligned}-\sqrt{9} \leq x &\leq \sqrt{9} \\-3 \leq x &\leq 3\end{aligned}$$

The solution set is $\{x | -3 \leq x \leq 3\}$.

91. $x^2 > 4$

$$\begin{aligned}x < -\sqrt{4} \text{ or } x &> \sqrt{4} \\x < -2 \text{ or } x &> 2\end{aligned}$$

The solution set is $\{x | x < -2 \text{ or } x > 2\}$.

92. $x^2 \geq 16$

$$\begin{aligned}x \leq -\sqrt{16} \text{ or } x &\geq \sqrt{16} \\x \leq -4 \text{ or } x &\geq 4\end{aligned}$$

The solution set is $\{x | x < -4 \text{ or } x > 4\}$.

93. $|3x - |2x + 1|| = 4$

$$\begin{aligned}3x - |2x + 1| &= 4 \text{ or } 3x - |2x + 1| = -4 \\3x - |2x + 1| &= 4 \\3x - 4 &= |2x + 1|\end{aligned}$$

Chapter 1: Equations and Inequalities

$$\begin{aligned} 2x+1 &= 3x-4 \quad \text{or} \quad 2x+1 = -(3x-4) \\ -x &= -5 \quad \text{or} \quad 2x+1 = -3x+4 \\ x &= 5 \quad \text{or} \quad 5x = 3 \\ x &= 5 \quad \text{or} \quad x = \frac{3}{5} \end{aligned}$$

or

$$\begin{aligned} 3x - |2x+1| &= -4 \\ 3x + 4 &= |2x+1| \end{aligned}$$

$$\begin{aligned} 2x+1 &= 3x+4 \quad \text{or} \quad 2x+1 = -(3x+4) \\ -x &= 3 \quad \text{or} \quad 2x+1 = -3x-4 \\ x &= -3 \quad \text{or} \quad 5x = -5 \\ x &= -3 \quad \text{or} \quad x = -1 \end{aligned}$$

The only values that check in the original equation are $x = 5$ and $x = -1$.

The solution set is $\{-1, 5\}$.

$$\begin{aligned} 94. \quad |x+|3x-2|| &= 2 \\ x+|3x-2| &= 2 \quad \text{or} \quad x+|3x-2| = -2 \\ x+|3x-2| &= 2 \\ |3x-2| &= 2-x \\ 3x-2 &= 2-x \quad \text{or} \quad 3x-2 = -(2-x) \\ 4x &= 4 \quad \text{or} \quad 3x-2 = -2+x \\ x &= 1 \quad \text{or} \quad 2x = 0 \\ x &= 1 \quad \text{or} \quad x = 0 \end{aligned}$$

or

$$\begin{aligned} x+|3x-2| &= -2 \\ |3x-2| &= -2-x \\ 3x-2 &= -2-x \quad \text{or} \quad 3x-2 = -(-2-x) \\ 4x &= 0 \quad \text{or} \quad 3x-2 = 2+x \\ x &= 0 \quad \text{or} \quad 2x = 4 \\ x &= 0 \quad \text{or} \quad x = 2 \end{aligned}$$

The only values that check in the original equation are $x = 0$ and $x = 1$.

The solution set is $\{0, 1\}$.

95 – 97. Answers will vary.

Section 1.7

1. mathematical modeling
2. interest
3. uniform motion

4. False; the amount charged for the use of principal is the interest.
5. True; this is the uniform motion formula.
6. a
7. b
8. c
9. Let A represent the area of the circle and r the radius. The area of a circle is the product of π times the square of the radius: $A = \pi r^2$
10. Let C represent the circumference of a circle and r the radius. The circumference of a circle is the product of π times twice the radius: $C = 2\pi r$
11. Let A represent the area of the square and s the length of a side. The area of the square is the square of the length of a side: $A = s^2$
12. Let P represent the perimeter of a square and s the length of a side. The perimeter of a square is four times the length of a side: $P = 4s$
13. Let F represent the force, m the mass, and a the acceleration. Force equals the product of the mass times the acceleration: $F = ma$
14. Let P represent the pressure, F the force, and A the area. Pressure is the force per unit area:

$$P = \frac{F}{A}$$
15. Let W represent the work, F the force, and d the distance. Work equals force times distance: $W = Fd$
16. Let K represent the kinetic energy, m the mass, and v the velocity. Kinetic energy is one-half the product of the mass and the square of the velocity: $K = \frac{1}{2}mv^2$
17. C = total variable cost in dollars, x = number of dishwashers manufactured: $C = 150x$
18. R = total revenue in dollars, x = number of dishwashers sold: $R = 250x$

Section 1.7: Problem Solving: Interest, Mixture, Uniform Motion, and Constant Rate Job Applications

19. Let x represent the amount of money invested in bonds. Then $50,000 - x$ represents the amount of money invested in CD's. Since the total interest is to be \$6,000, we have:

$$0.15x + 0.07(50,000 - x) = 6,000$$

$$(100)(0.15x + 0.07(50,000 - x)) = (6,000)(100)$$

$$15x + 7(50,000 - x) = 600,000$$

$$15x + 350,000 - 7x = 600,000$$

$$8x + 350,000 = 600,000$$

$$8x = 250,000$$

$$x = 31,250$$

\$31,250 should be invested in bonds at 15% and \$18,750 should be invested in CD's at 7%.

20. Let x represent the amount of money invested in bonds. Then $50,000 - x$ represents the amount of money invested in CD's. Since the total interest is to be \$7,000, we have:

$$0.15x + 0.07(50,000 - x) = 7,000$$

$$(100)(0.15x + 0.07(50,000 - x)) = (7,000)(100)$$

$$15x + 7(50,000 - x) = 700,000$$

$$15x + 350,000 - 7x = 700,000$$

$$8x + 350,000 = 700,000$$

$$8x = 350,000$$

$$x = 43,750$$

\$43,750 should be invested in bonds at 15% and \$6,250 should be invested in CD's at 7%.

21. Let x represent the amount of money loaned at 8%. Then $12,000 - x$ represents the amount of money loaned at 18%. Since the total interest is to be \$1,000, we have:

$$0.08x + 0.18(12,000 - x) = 1,000$$

$$(100)(0.08x + 0.18(12,000 - x)) = (1,000)(100)$$

$$8x + 18(12,000 - x) = 100,000$$

$$8x + 216,000 - 18x = 100,000$$

$$-10x + 216,000 = 100,000$$

$$-10x = -116,000$$

$$x = 11,600$$

\$11,600 is loaned at 8% and \$400 is at 18%.

22. Let x represent the amount of money loaned at 16%. Then $1,000,000 - x$ represents the amount of money loaned at 19%. Since the total interest is to be \$1,000,000(0.18), we have:

$$0.16x + 0.19(1,000,000 - x) = 1,000,000(0.18)$$

$$0.16x + 190,000 - 0.19x = 180,000$$

$$-0.03x + 190,000 = 180,000$$

$$-0.03x = -10,000$$

$$x = \frac{-10,000}{-0.03}$$

$$x = \$333,333.33$$

Wendy can lend \$333,333.33 at 16%.

23. Let x represent the number of pounds of Earl Gray tea. Then $100 - x$ represents the number of pounds of Orange Pekoe tea.

$$5x + 3(100 - x) = 4.50(100)$$

$$5x + 300 - 3x = 450$$

$$2x + 300 = 450$$

$$2x = 150$$

$$x = 75$$

75 pounds of Earl Gray tea must be blended with 25 pounds of Orange Pekoe.

24. Let x represent the number of pounds of the first kind of coffee. Then $100 - x$ represents the number of pounds of the second kind of coffee.

$$2.75x + 5(100 - x) = 3.90(100)$$

$$2.75x + 500 - 5x = 390$$

$$-2.25x + 500 = 390$$

$$-2.25x = -110$$

$$x \approx 48.9$$

Approximately 49 pounds of the first kind of coffee must be blended with approximately 51 pounds of the second kind of coffee.

25. Let x represent the number of pounds of cashews. Then $x + 60$ represents the number of pounds in the mixture.

$$9x + 3.50(60) = 7.50(x + 60)$$

$$9x + 210 = 7.50x + 450$$

$$1.5x = 240$$

$$x = 160$$

160 pounds of cashews must be added to the 60 pounds of almonds.

26. Let x represent the number of caramels in the box. Then $30 - x$ represents the number of cremes in the box.

Chapter 1: Equations and Inequalities

$$\begin{aligned} \text{Revenue} - \text{Cost} &= \text{Profit} \\ 12.50 - (0.25x + 0.45(30 - x)) &= 3.00 \\ 12.50 - (0.25x + 13.5 - 0.45x) &= 3.00 \\ 12.50 - (13.5 - 0.20x) &= 3.00 \\ 12.50 - 13.50 + 0.20x &= 3.00 \\ -1.00 + 0.20x &= 3.00 \\ 0.20x &= 4.00 \\ x &= 20 \end{aligned}$$

The box should contain 20 caramels and 10 cremes.

27. Let r represent the speed of the current.

	Rate	Time	Distance
Upstream	$16 - r$	$\frac{20}{60} = \frac{1}{3}$	$\frac{16-r}{3}$
Downstream	$16 + r$	$\frac{15}{60} = \frac{1}{4}$	$\frac{16+r}{4}$

Since the distance is the same in each direction:

$$\begin{aligned} \frac{16-r}{3} &= \frac{16+r}{4} \\ 4(16-r) &= 3(16+r) \\ 64 - 4r &= 48 + 3r \\ 16 &= 7r \\ r &= \frac{16}{7} \approx 2.286 \end{aligned}$$

The speed of the current is approximately 2.286 miles per hour.

28. Let r represent the speed of the motorboat.

	Rate	Time	Distance
Upstream	$r - 3$	5	$5(r - 3)$
Downstream	$r + 3$	2.5	$2.5(r + 3)$

The distance is the same in each direction:

$$\begin{aligned} 5(r - 3) &= 2.5(r + 3) \\ 5r - 15 &= 2.5r + 7.5 \\ 2.5r &= 22.5 \\ r &= 9 \end{aligned}$$

The speed of the motorboat is 9 miles per hour.

29. Let r represent the speed of the current.

	Rate	Time	Distance
Upstream	$15 - r$	$\frac{10}{15-r}$	10
Downstream	$15 + r$	$\frac{10}{15+r}$	10

Since the total time is 1.5 hours, we have:

$$\begin{aligned} \frac{10}{15-r} + \frac{10}{15+r} &= 1.5 \\ 10(15+r) + 10(15-r) &= 1.5(15-r)(15+r) \\ 150 + 10r + 150 - 10r &= 1.5(225 - r^2) \\ 300 &= 1.5(225 - r^2) \\ 200 &= 225 - r^2 \\ r^2 - 25 &= 0 \\ (r-5)(r+5) &= 0 \\ r = 5 \text{ or } r &= -5 \end{aligned}$$

Speed must be positive, so disregard $r = -5$. The speed of the current is 5 miles per hour.

30. Let r represent the rate of the slower car. Then $r + 10$ represents the rate of the faster car.

	Rate	Time	Distance
Slower car	r	3.5	$3.5r$
Faster car	$r + 10$	3	$3(r + 10)$

$$3.5r = 3(r + 10)$$

$$3.5r = 3r + 30$$

$$0.5r = 30$$

$$r = 60$$

The slower car travels at a rate of 60 miles per hour. The faster car travels at a rate of 70 miles per hour. The distance is $(70)(3) = 210$ miles.

31. Let r represent Karen's normal walking speed.

	Rate	Time	Distance
With walkway	$r + 2.5$	$\frac{50}{r+2.5}$	50
Against walkway	$r - 2.5$	$\frac{50}{r-2.5}$	50

Since the total time is 40 seconds:

$$\begin{aligned} \frac{50}{r+2.5} + \frac{50}{r-2.5} &= 40 \\ 50(r-2.5) + 50(r+2.5) &= 40(r-2.5)(r+2.5) \\ 50r - 125 + 50r + 125 &= 40(r^2 - 6.25) \\ 100r &= 40r^2 - 250 \\ 0 &= 40r^2 - 100r - 250 \\ 0 &= 4r^2 - 10r - 25 \\ r &= \frac{-(-10) \pm \sqrt{(-10)^2 - 4(4)(-25)}}{2(4)} \\ &= \frac{10 \pm \sqrt{500}}{8} = \frac{10 \pm 10\sqrt{5}}{8} = \frac{5 \pm 5\sqrt{5}}{4} \\ r &\approx 4.05 \text{ or } r \approx -1.55 \end{aligned}$$

Speed must be positive, so disregard $r \approx -1.55$. Karen's normal walking speed is approximately 4.05 feet per second.

Section 1.7: Problem Solving: Interest, Mixture, Uniform Motion, and Constant Rate Job Applications

32. Let r represent the speed of the airport walkway.

	Rate	Time	Distance
Walking with	$1.5 + r$	$\frac{280}{1.5 + r}$	280
Standing still	r	$\frac{280}{r}$	280

Walking with the walkway takes 60 seconds less time than standing still on the walkway:

$$\begin{aligned} \frac{280}{1.5 + r} &= \frac{280}{r} - 60 \\ 280r &= 280(1.5 + r) - 60r(r + 1.5) \\ 280r &= 420 + 280r - 60r^2 - 90r \\ 60r^2 + 90r - 420 &= 0 \\ 2r^2 + 3r - 14 &= 0 \\ (2r + 7)(r - 2) &= 0 \\ 2r + 7 = 0 &\quad \text{or} \quad r - 2 = 0 \\ r = -\frac{7}{2} &\quad \text{or} \quad r = 2 \end{aligned}$$

Speed must be positive, so disregard $r = -\frac{7}{2}$.

The speed of the airport walkway is 2 meters per second.

33. Let w represent the width of a regulation doubles tennis court. Then $2w + 6$ represents the length. The area is 2808 square feet:

$$\begin{aligned} w(2w + 6) &= 2808 \\ 2w^2 + 6w &= 2808 \\ 2w^2 + 6w - 2808 &= 0 \\ w^2 + 3w - 1404 &= 0 \\ (w + 39)(w - 36) &= 0 \\ w + 39 = 0 &\quad \text{or} \quad w - 36 = 0 \\ w = -39 &\quad \text{or} \quad w = 36 \end{aligned}$$

The width must be positive, so disregard $w = -39$.

The width of a regulation doubles tennis court is 36 feet and the length is $2(36) + 6 = 78$ feet.

34. Let t represent the time it takes the HP LaserJet CP4025dn to complete the print job alone. Then $t + 16$ represents the time it takes the HP LaserJet M451dw to complete the print job alone.

	Time to do job	Part of job done in one minute
HP LJ 2420	t	$\frac{1}{t}$
HP LJ 1300	$t + 16$	$\frac{1}{t+16}$
Together	15	$\frac{1}{15}$

$$\frac{1}{t} + \frac{1}{t+16} = \frac{1}{15}$$

$$15(t + 16) + 15t = t(t + 16)$$

$$15t + 240 + 15t = t^2 + 16t$$

$$0 = t^2 - 14t - 240$$

$$0 = (t - 24)(t + 10)$$

$$t - 24 = 0 \quad \text{or} \quad t + 10 = 0$$

$$t = 24 \quad \text{or} \quad t = -10$$

Time must be positive, so disregard $t = -10$.

The HP LaserJet CP4025dn takes 24 minutes to complete the job alone, printing $\frac{840}{24} = 35$ pages per minute. The HP LaserJet M451dw takes $24 + 16 = 40$ minutes to complete the job alone, printing $\frac{840}{40} = 21$ pages per minute.

35. Let t represent the time it takes to do the job together.

	Time to do job	Part of job done in one minute
Trent	30	$\frac{1}{30}$
Lois	20	$\frac{1}{20}$
Together	t	$\frac{1}{t}$

$$\frac{1}{30} + \frac{1}{20} = \frac{1}{t}$$

$$2t + 3t = 60$$

$$5t = 60$$

$$t = 12$$

Working together, the job can be done in 12 minutes.

Chapter 1: Equations and Inequalities

36. Let t represent the time it takes April to do the job working alone.

	Time to do job	Part of job done in one hour
Patrice	10	$\frac{1}{10}$
April	t	$\frac{1}{t}$
Together	6	$\frac{1}{6}$

$$\frac{1}{10} + \frac{1}{t} = \frac{1}{6}$$

$$3t + 30 = 5t$$

$$2t = 30$$

$$t = 15$$

April would take 15 hours to paint the rooms.

37. l = length of the garden

w = width of the garden

- a. The length of the garden is to be twice its width. Thus, $l = 2w$.

The dimensions of the fence are $l + 4$ and $w + 4$.

The perimeter is 46 feet, so:

$$2(l + 4) + 2(w + 4) = 46$$

$$2(2w + 4) + 2(w + 4) = 46$$

$$4w + 8 + 2w + 8 = 46$$

$$6w + 16 = 46$$

$$6w = 30$$

$$w = 5$$

The dimensions of the garden are 5 feet by 10 feet.

- b. Area = $l \cdot w = 5 \cdot 10 = 50$ square feet
- c. If the dimensions of the garden are the same, then the length and width of the fence are also the same ($l + 4$). The perimeter is 46 feet, so:

$$2(l + 4) + 2(l + 4) = 46$$

$$2l + 8 + 2l + 8 = 46$$

$$4l + 16 = 46$$

$$4l = 30$$

$$l = 7.5$$

The dimensions of the garden are 7.5 feet by 7.5 feet.

- d. Area = $l \cdot w = 7.5(7.5) = 56.25$ square feet.

38. l = length of the pond
 w = width of the pond

- a. The pond is to be a square. Thus, $l = w$. The dimensions of the fenced area are $w + 6$ on each side. The perimeter is 100 feet, so:

$$4(w + 6) = 100$$

$$4w + 24 = 100$$

$$4w = 76$$

$$w = 19$$

The dimensions of the pond are 19 feet by 19 feet.

- b. The length of the pond is to be three times the width. Thus, $l = 3w$. The dimensions of the fenced area are $w + 6$ and $l + 6$. The perimeter is 100 feet, so:

$$2(w + 6) + 2(l + 6) = 100$$

$$2(w + 6) + 2(3w + 6) = 100$$

$$2w + 12 + 6w + 12 = 100$$

$$8w + 24 = 100$$


$$8w = 76$$

$$w = 9.5$$

$$l = 3(9.5) = 28.5$$

The dimensions of the pond are 9.5 feet by 28.5 feet.

- c. If the pond is circular, the diameter is d and the diameter of the circle with the pond and the deck is $d + 6$.



The perimeter is 100 feet, so:

$$\pi(d + 6) = 100$$

$$\pi d + 6\pi = 100$$

$$\pi d = 100 - 6\pi$$

$$d = \frac{100}{\pi} - 6 \approx 25.83$$

The diameter of the pond is 25.83 feet.

- d. $\text{Area}_{\text{square}} = l \cdot w = 19(19) = 361 \text{ ft}^2$.

$$\text{Area}_{\text{rectangle}} = l \cdot w = 28.5(9.5) = 270.75 \text{ ft}^2$$

$$\text{Area}_{\text{circle}} = \pi r^2 = \pi \left(\frac{25.83}{2} \right)^2 \approx 524 \text{ ft}^2$$

The circular pond has the largest area.

Section 1.7: Problem Solving: Interest, Mixture, Uniform Motion, and Constant Rate Job Applications

39. Let t represent the time it takes for the defensive back to catch the tight end.

	Time to run 100 yards	Time	Rate	Distance
Tight End	12 sec	t	$\frac{100}{12} = \frac{25}{3}$	$\frac{25}{3}t$
Def. Back	10 sec	t	$\frac{100}{10} = 10$	$10t$

Since the defensive back has to run 5 yards farther, we have:

$$\frac{25}{3}t + 5 = 10t$$

$$25t + 15 = 30t$$

$$15 = 5t$$

$$t = 3 \quad \rightarrow \quad 10t = 30$$

The defensive back will catch the tight end at the 45 yard line ($15 + 30 = 45$).

40. Let x represent the number of highway miles traveled. Then $30,000 - x$ represents the number of city miles traveled.

$$\begin{aligned} \frac{x}{40} + \frac{30,000 - x}{25} &= 900 \\ 200\left(\frac{x}{40} + \frac{30,000 - x}{25}\right) &= 200(900) \\ 5x + 240,000 - 8x &= 180,000 \\ -3x + 240,000 &= 180,000 \\ -3x &= -60,000 \\ x &= 20,000 \end{aligned}$$

Therese is allowed to claim 20,000 miles as a business expense.

41. Let x represent the number of gallons of pure water. Then $x+1$ represents the number of gallons in the 60% solution.

$$(\%)(\text{gallons}) + (\%)(\text{gallons}) = (\%)(\text{gallons})$$

$$\begin{aligned} 0(x) + 1(1) &= 0.60(x+1) \\ 1 &= 0.6x + 0.6 \\ 0.4 &= 0.6x \\ x &= \frac{4}{6} = \frac{2}{3} \end{aligned}$$

$\frac{2}{3}$ gallon of pure water should be added.

42. Let x represent the number of liters to be drained and replaced with pure antifreeze.

$$(\%)(\text{liters}) + (\%)(\text{liters}) = (\%)(\text{liters})$$

$$1(x) + 0.40(15 - x) = 0.60(15)$$

$$x + 6 - 0.40x = 9$$

$$0.60x = 3$$

$$x = 5$$

5 liters should be drained and replaced with pure antifreeze.

43. Let x represent the number of ounces of water to be evaporated; the amount of salt remains the same. Therefore, we get

$$0.04(32) = 0.06(32 - x)$$

$$1.28 = 1.92 - 0.06x$$

$$0.06x = 0.64$$

$$x = \frac{0.64}{0.06} = \frac{64}{6} = \frac{32}{3} = 10\frac{2}{3}$$

$10\frac{2}{3} \approx 10.67$ ounces of water need to be evaporated.

44. Let x represent the number of gallons of water to be evaporated; the amount of salt remains the same.

$$0.03(240) = 0.05(240 - x)$$

$$7.2 = 12 - 0.05x$$

$$0.05x = 4.8$$

$$x = \frac{4.8}{0.05} = 96$$

96 gallons of water need to be evaporated.

45. Let x represent the number of grams of pure gold. Then $60 - x$ represents the number of grams of 12 karat gold to be used.

$$x + \frac{1}{2}(60 - x) = \frac{2}{3}(60)$$

$$x + 30 - 0.5x = 40$$

$$0.5x = 10$$

$$x = 20$$

20 grams of pure gold should be mixed with 40 grams of 12 karat gold.

46. Let x represent the number of atoms of oxygen. $2x$ represents the number of atoms of hydrogen. $x+1$ represents the number of atoms of carbon.

$$x + 2x + x + 1 = 45$$

$$4x = 44$$

$$x = 11$$

There are 11 atoms of oxygen and 22 atoms of hydrogen in the sugar molecule.

Chapter 1: Equations and Inequalities

47. Let t represent the time it takes for Mike to catch up with Dan. Since the distances are the same, we have:

$$\frac{1}{6}t = \frac{1}{9}(t+1)$$

$$3t = 2t + 2$$

$$t = 2$$

Mike will pass Dan after 2 minutes, which is a distance of $\frac{1}{3}$ mile.

48. Let t represent the time of flight with the wind. The distance is the same in each direction:

$$330t = 270(5-t)$$

$$330t = 1350 - 270t$$

$$600t = 1350$$

$$t = 2.25$$

The distance the plane can fly and still return safely is $330(2.25) = 742.5$ miles.

49. Let t represent the time the auxiliary pump needs to run. Since the two pumps are emptying one tanker, we have:

$$\frac{3}{4} + \frac{t}{9} = 1$$

$$27 + 4t = 36$$

$$4t = 9$$

$$t = \frac{9}{4} = 2.25$$

The auxiliary pump must run for 2.25 hours. It must be started at 9:45 a.m.

50. Let x represent the number of pounds of pure cement. Then $x+20$ represents the number of pounds in the 40% mixture.

$$x + 0.25(20) = 0.40(x + 20)$$

$$x + 5 = 0.4x + 8$$

$$0.6x = 3$$

$$x = \frac{30}{6} = 5$$

5 pounds of pure cement should be added.

51. Let t represent the time for the tub to fill with the faucets on and the stopper removed. Since one tub is being filled, we have:

$$\frac{t}{15} + \left(-\frac{t}{20}\right) = 1$$

$$4t - 3t = 60$$

$$t = 60$$

60 minutes is required to fill the tub.

52. Let t be the time the 5 horsepower pump needs to run to finish emptying the pool. Since the two pumps are emptying one pool, we have:

$$\frac{t+2}{5} + \frac{2}{8} = 1$$

$$4(2+t) + 5 = 20$$

$$8 + 4t + 5 = 20$$

$$4t = 7$$

$$t = 1.75$$

The 5 horsepower pump must run for an additional 1.75 hours or 1 hour and 45 minutes to empty the pool.

53. Let t represent the time spent running. Then $5-t$ represents the time spent biking.

	Rate	Time	Distance
Run	6	t	$6t$
Bike	25	$5-t$	$25(5-t)$

The total distance is 87 miles:

$$6t + 25(5-t) = 87$$

$$6t + 125 - 25t = 87$$

$$-19t + 125 = 87$$

$$-19t = -38$$

$$t = 2$$

The time spent running is 2 hours, so the distance of the run is $6(2) = 12$ miles. The distance of the bicycle race is $25(5-2) = 75$ miles.

54. Let r represent the speed of the eastbound cyclist. Then $r+5$ represents the speed of the westbound cyclist.

	Rate	Time	Distance
Eastbound	r	6	$6r$
Westbound	$r+5$	6	$6(r+5)$

The total distance is 246 miles:

$$6r + 6(r+5) = 246$$

$$6r + 6r + 30 = 246$$

$$12r + 30 = 246$$

$$12r = 216$$

$$r = 18$$

The speed of the eastbound cyclist is 18 miles per hour, and the speed of the westbound cyclist is $18 + 5 = 23$ miles per hour.

55. Burke's rate is $\frac{100}{12}$ meters/sec. In 9.69 seconds, Burke will run $\frac{100}{12}(9.69) = 80.75$ meters. Bolt would win by 19.25 meters.

56. $A = 2\pi r^2 + 2\pi r h$. Since $A = 188.5$ square inches and $h = 7$ inches,

$$\begin{aligned} 2\pi r^2 + 2\pi r(7) &= 188.5 \\ 2\pi r^2 + 14\pi r - 188.5 &= 0 \\ r &= \frac{-14\pi \pm \sqrt{(14\pi)^2 - 4(2\pi)(-188.5)}}{2(2\pi)} \\ &= \frac{-14\pi \pm \sqrt{6671.9642}}{4\pi} \end{aligned}$$

$$r \approx 3 \text{ or } r \approx -10$$

The radius of the coffee can is approximately 3 inches.

57. Let x be the original selling price of the shirt.
Profit = Revenue – Cost

$$4 = x - 0.40x - 20 \rightarrow 24 = 0.60x \rightarrow x = 40$$

The original price should be \$40 to ensure a profit of \$4 after the sale.

If the sale is 50% off, the profit is:

$$40 - 0.50(40) - 20 = 40 - 20 - 20 = 0$$

At 50% off there will be no profit.

58. Answers will vary.

59. It is impossible to mix two solutions with a lower concentration and end up with a new solution with a higher concentration.

Algebraic Solution:

Let x = the number of liters of 25% solution.

$$(\%)(\text{liters}) + (\%)(\text{liters}) = (\%)(\text{liters})$$

$$0.25x + 0.48(20) = 0.58(20 + x)$$

$$0.25x + 9.6 = 10.6 + 0.58x$$

$$-0.33x = 1$$

$$x \approx -3.03 \text{ liters}$$

(not possible)

60. Let t_1 and t_2 represent the times for the two segments of the trip. Since Atlanta is halfway between Chicago and Miami, the distances are equal.

$$45t_1 = 55t_2$$

$$t_1 = \frac{55}{45}t_2$$

$$t_1 = \frac{11}{9}t_2$$

Computing the average speed:

$$\text{Avg Speed} = \frac{\text{Distance}}{\text{Time}} = \frac{45t_1 + 55t_2}{t_1 + t_2}$$

$$= \frac{45\left(\frac{11}{9}t_2\right) + 55t_2}{\frac{11}{9}t_2 + t_2} = \frac{55t_2 + 55t_2}{\left(\frac{11t_2 + 9t_2}{9}\right)}$$

$$= \frac{110t_2}{\left(\frac{20t_2}{9}\right)} = \frac{990t_2}{20t_2}$$

$$= \frac{99}{2} = 49.5 \text{ miles per hour}$$

The average speed for the trip from Chicago to Miami is 49.5 miles per hour.

61. The time traveled with the tail wind was:

$$t = \frac{919}{550} \approx 1.67091 \text{ hours}.$$

Since they were 20 minutes ($\frac{1}{3}$ hour) early, the

time in still air would have been:

$$\begin{aligned} 1.67091 \text{ hrs} + 20 \text{ min} &= (1.67091 + 0.33333) \text{ hrs} \\ &\approx 2.00424 \text{ hrs} \end{aligned}$$

Thus, with no wind, the ground speed is

$$\frac{919}{2.00424} \approx 458.53. \text{ Therefore, the tail wind is}$$

$$550 - 458.53 = 91.47 \text{ knots}.$$

Chapter 1 Review

$$1. 2 - \frac{x}{3} = 8$$

$$6 - x = 24$$

$$x = -18$$

The solution set is $\{-18\}$.

Chapter 1: Equations and Inequalities

2. $-2(5 - 3x) + 8 = 4 + 5x$
 $-10 + 6x + 8 = 4 + 5x$
 $6x - 2 = 4 + 5x$
 $x = 6$

The solution set is $\{6\}$.

3. $\frac{3x}{4} - \frac{x}{3} = \frac{1}{12}$
 $9x - 4x = 1$
 $5x = 1$
 $x = \frac{1}{5}$

The solution set is $\left\{\frac{1}{5}\right\}$.

7. $\frac{x}{x-1} = \frac{6}{5}$
 $5x = 6x - 6$
 $6 = x$

Since $x = 6$ does not cause a denominator to equal zero, the solution set is $\{6\}$.

5. $x(1-x) = 6$
 $x - x^2 = 6$
 $0 = x^2 - x + 6$
 $b^2 - 4ac = (-1)^2 - 4(1)(6)$
 $= 1 - 24 = -23$

Therefore, there are no real solutions.

6. $x(1+x) = 6$
 $x + x^2 = 6$
 $x^2 + x - 6 = 0$
 $(x+3)(x-2) = 0$
 $x = -3 \text{ or } x = 2$

The solution set is $\{-3, 2\}$.

7. $\frac{1}{2}\left(x - \frac{1}{3}\right) = \frac{3}{4} - \frac{x}{6}$
 $(12)\left(\frac{1}{2}\right)\left(x - \frac{1}{3}\right) = \left(\frac{3}{4} - \frac{x}{6}\right)(12)$
 $6x - 2 = 9 - 2x$
 $8x = 11$
 $x = \frac{11}{8}$

The solution set is $\left\{\frac{11}{8}\right\}$.

8. $\frac{1-3x}{4} = \frac{x+6}{3} + \frac{1}{2}$
 $(12)\left(\frac{1-3x}{4}\right) = \left(\frac{x+6}{3} + \frac{1}{2}\right)(12)$
 $3(1-3x) = 4(x+6) + 6$
 $3 - 9x = 4x + 24 + 6$
 $-13x = 27$
 $x = -\frac{27}{13}$

The solution set is $\left\{-\frac{27}{13}\right\}$.

9. $(x-1)(2x+3) = 3$
 $2x^2 + x - 3 = 3$
 $2x^2 + x - 6 = 0$
 $(2x-3)(x+2) = 0$
 $x = \frac{3}{2} \text{ or } x = -2$

The solution set is $\left\{-2, \frac{3}{2}\right\}$.

10. $2x+3 = 4x^2$
 $0 = 4x^2 - 2x - 3$
 $x = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(4)(-3)}}{2(4)}$
 $= \frac{2 \pm \sqrt{52}}{8} = \frac{2 \pm 2\sqrt{13}}{8} = \frac{1 \pm \sqrt{13}}{4}$

The solution set is $\left\{\frac{1-\sqrt{13}}{4}, \frac{1+\sqrt{13}}{4}\right\}$.

11. $\sqrt[3]{x^2 - 1} = 2$
 $(\sqrt[3]{x^2 - 1})^3 = (2)^3$
 $x^2 - 1 = 8$
 $x^2 = 9$
 $x = \pm 3$

Check $x = -3$: $\sqrt[3]{(-3)^2 - 1} = 2$
 $\sqrt[3]{9 - 1} = 2$
 $\sqrt[3]{8} = 2$
 $2 = 2$

Check $x = 3$: $\sqrt[3]{(3)^2 - 1} = 2$
 $\sqrt[3]{9 - 1} = 2$
 $\sqrt[3]{8} = 2$
 $2 = 2$

The solution set is $\{-3, 3\}$.

12. $\sqrt{1+x^3} = 3$

$$(\sqrt{1+x^3})^2 = (3)^2$$

$$1+x^3 = 9$$

$$x^3 = 8$$

$$x = \sqrt[3]{8} = 2$$

Check $x = 2$: $\sqrt{1+(2)^3} = 3$

$$\sqrt{9} = 3$$

$$3 = 3$$

The solution set is $\{2\}$.

13. $x(x+1)+2=0$

$$x^2 + x + 2 = 0$$

$$x = \frac{-1 \pm \sqrt{(1)^2 - 4(1)(2)}}{2(1)} = \frac{-1 \pm \sqrt{-7}}{2}$$

No real solutions.

14. $x^4 - 5x^2 + 4 = 0$

$$(x^2 - 4)(x^2 - 1) = 0$$

$$x^2 - 4 = 0 \text{ or } x^2 - 1 = 0$$

$$x = \pm 2 \text{ or } x = \pm 1$$

The solution set is $\{-2, -1, 1, 2\}$.

15. $\sqrt{2x-3} + x = 3$

$$\sqrt{2x-3} = 3 - x$$

$$2x - 3 = 9 - 6x + x^2$$

$$x^2 - 8x + 12 = 0$$

$$(x-2)(x-6) = 0$$

$$x = 2 \text{ or } x = 6$$

Check $x = 2$: $\sqrt{2(2)-3} + 2 = \sqrt{1} + 2 = 3$

Check $x = 6$: $\sqrt{2(6)-3} + 6 = \sqrt{9} + 6 = 9 \neq 3$

The solution set is $\{2\}$.

16. $\sqrt[4]{2x+3} = 2$

$$(\sqrt[4]{2x+3})^4 = 2^4$$

$$2x+3 = 16$$

$$2x = 13$$

$$x = \frac{13}{2}$$

Check $x = \frac{13}{2}$:

$$\sqrt[4]{2\left(\frac{13}{2}\right) + 3} = \sqrt[4]{13+3} = \sqrt[4]{16} = 2$$

The solution set is $\left\{\frac{13}{2}\right\}$.

17. $\sqrt{x+1} + \sqrt{x-1} = \sqrt{2x+1}$

$$(\sqrt{x+1} + \sqrt{x-1})^2 = (\sqrt{2x+1})^2$$

$$x+1 + 2\sqrt{x+1}\sqrt{x-1} + x-1 = 2x+1$$

$$2x + 2\sqrt{x+1}\sqrt{x-1} = 2x+1$$

$$2\sqrt{x+1}\sqrt{x-1} = 1$$

$$(2\sqrt{x+1}\sqrt{x-1})^2 = (1)^2$$

$$4(x+1)(x-1) = 1$$

$$4x^2 - 4 = 1$$

$$4x^2 = 5$$

$$x^2 = \frac{5}{4}$$

$$x = \pm \frac{\sqrt{5}}{2}$$

Check $x = \frac{\sqrt{5}}{2}$:

$$\sqrt{\frac{\sqrt{5}}{2} + 1} + \sqrt{\frac{\sqrt{5}}{2} - 1} = \sqrt{2\left(\frac{\sqrt{5}}{2}\right) + 1}$$

$$1.79890743995 = 1.79890743995$$

Check $x = -\frac{\sqrt{5}}{2}$:

$$\sqrt{-\frac{\sqrt{5}}{2} + 1} + \sqrt{-\frac{\sqrt{5}}{2} - 1} = \sqrt{2\left(-\frac{\sqrt{5}}{2}\right) + 1},$$

The second solution is not possible because it makes the radicand negative.

The solution set is $\left\{\frac{\sqrt{5}}{2}\right\}$.

Chapter 1: Equations and Inequalities

18. $\sqrt{2x-1} - \sqrt{x-5} = 3$

$$\sqrt{2x-1} = 3 + \sqrt{x-5}$$

$$(\sqrt{2x-1})^2 = (3 + \sqrt{x-5})^2$$

$$2x-1 = 9 + 6\sqrt{x-5} + x-5$$

$$x-5 = 6\sqrt{x-5}$$

$$(x-5)^2 = (6\sqrt{x-5})^2$$

$$x^2 - 10x + 25 = 36(x-5)$$

$$x^2 - 10x + 25 = 36x - 180$$

$$x^2 - 46x + 205 = 0$$

$$(x-41)(x-5) = 0$$

$$x = 41 \text{ or } x = 5$$

Check $x = 41$:

$$\sqrt{2(41)-1} - \sqrt{41-5} = \sqrt{81} - \sqrt{36} = 9 - 6 = 3$$

Check $x = 5$:

$$\sqrt{2(5)-1} - \sqrt{5-5} = \sqrt{9} - \sqrt{0} = 3 - 0 = 3$$

The solution set is $\{5, 41\}$.

19. $2x^{1/2} - 3 = 0$

$$2x^{1/2} = 3$$

$$(2x^{1/2})^2 = 3^2$$

$$4x = 9$$

$$x = \frac{9}{4}$$

Check $x = \frac{9}{4}$:

$$2\left(\frac{9}{4}\right)^{1/2} - 3 = 2\left(\frac{3}{2}\right) - 3 = 3 - 3 = 0$$

The solution set is $\left\{\frac{9}{4}\right\}$.

20. $x^{-6} - 7x^{-3} - 8 = 0$

Let $u = x^{-3}$ so that $u^2 = x^{-6}$.

$$u^2 - 7u - 8 = 0$$

$$(u-8)(u+1) = 0$$

$$u = 8 \quad \text{or} \quad u = -1$$

$$x^{-3} = 8 \quad \text{or} \quad x^{-3} = -1$$

$$(x^{-3})^{-1/3} = (8)^{-1/3} \quad \text{or} \quad (x^{-3})^{-1/3} = (-1)^{-1/3}$$

$$x = \frac{1}{2} \quad \text{or} \quad x = -1$$

Check $\frac{1}{2} : \left(\frac{1}{2}\right)^{-6} - 7\left(\frac{1}{2}\right)^{-3} - 8 = 64 - 56 - 8 = 0$

Check $-1 : (-1)^{-6} - 7(-1)^{-3} - 8 = 1 + 7 - 8 = 0$

The solution set is $\left\{-1, \frac{1}{2}\right\}$.

21.

$$x^2 + m^2 = 2mx + (nx)^2$$

$$x^2 + m^2 = 2mx + n^2x^2$$

$$x^2 - n^2x^2 - 2mx + m^2 = 0$$

$$(1-n^2)x^2 - 2mx + m^2 = 0$$

$$x = \frac{-(2m) \pm \sqrt{(-2m)^2 - 4(1-n^2)m^2}}{2(1-n^2)}$$

$$= \frac{2m \pm \sqrt{4m^2 - 4m^2 + 4m^2n^2}}{2(1-n^2)}$$

$$= \frac{2m \pm \sqrt{4m^2n^2}}{2(1-n^2)} = \frac{2m \pm 2mn}{2(1-n^2)}$$

$$= \frac{2m(1 \pm n)}{2(1-n^2)} = \frac{m(1 \pm n)}{1-n^2}$$

$$x = \frac{m(1+n)}{1-n^2} = \frac{m(1+n)}{(1+n)(1-n)} = \frac{m}{1-n}$$

or

$$x = \frac{m(1-n)}{1-n^2} = \frac{m(1-n)}{(1+n)(1-n)} = \frac{m}{1+n}$$

The solution set is $\left\{\frac{m}{1-n}, \frac{m}{1+n}\right\}$, $n \neq 1$, $n \neq -1$.

22. $10a^2x^2 - 2abx - 36b^2 = 0$

$$5a^2x^2 - abx - 18b^2 = 0$$

$$(5ax+9b)(ax-2b) = 0$$

$$5ax+9b = 0 \quad \text{or} \quad ax-2b = 0$$

$$5ax = -9b \quad \text{or} \quad ax = 2b$$

$$x = -\frac{9b}{5a} \quad x = \frac{2b}{a}$$

The solution set is $\left\{-\frac{9b}{5a}, \frac{2b}{a}\right\}$, $a \neq 0$.

23. $\sqrt{x^2 + 3x + 7} - \sqrt{x^2 - 3x + 9} + 2 = 0$
 $\sqrt{x^2 + 3x + 7} = \sqrt{x^2 - 3x + 9} - 2$
 $(\sqrt{x^2 + 3x + 7})^2 = (\sqrt{x^2 - 3x + 9} - 2)^2$
 $x^2 + 3x + 7 = x^2 - 3x + 9 - 4\sqrt{x^2 - 3x + 9} + 4$
 $6x - 6 = -4\sqrt{x^2 - 3x + 9}$
 $(6(x-1))^2 = (-4\sqrt{x^2 - 3x + 9})^2$
 $36(x^2 - 2x + 1) = 16(x^2 - 3x + 9)$
 $36x^2 - 72x + 36 = 16x^2 - 48x + 144$
 $20x^2 - 24x - 108 = 0$
 $5x^2 - 6x - 27 = 0$
 $(5x+9)(x-3) = 0$
 $x = -\frac{9}{5} \text{ or } x = 3$

Check $x = -\frac{9}{5}$:

$$\begin{aligned} & \sqrt{\left(-\frac{9}{5}\right)^2 + 3\left(-\frac{9}{5}\right) + 7} - \sqrt{\left(-\frac{9}{5}\right)^2 - 3\left(-\frac{9}{5}\right) + 9} + 2 \\ &= \sqrt{\frac{81}{25} - \frac{27}{5} + 7} - \sqrt{\frac{81}{25} + \frac{27}{5} + 9} + 2 \\ &= \sqrt{\frac{81-135+175}{25}} - \sqrt{\frac{81+135+225}{25}} + 2 \\ &= \sqrt{\frac{121}{25}} - \sqrt{\frac{441}{25}} + 2 = \frac{11}{5} - \frac{21}{5} + 2 = 0 \end{aligned}$$

Check $x = 3$:

$$\begin{aligned} & \sqrt{(3)^2 + 3(3) + 7} - \sqrt{(3)^2 - 3(3) + 9} + 2 \\ &= \sqrt{9+9+7} - \sqrt{9-9+9} + 2 \\ &= \sqrt{25} - \sqrt{9} + 2 = 2+2 \\ &= 4 \neq 0 \end{aligned}$$

The solution set is $\left\{-\frac{9}{5}\right\}$.

24. $|2x+3| = 7$
 $2x+3 = 7 \text{ or } 2x+3 = -7$
 $2x = 4 \text{ or } 2x = -10$
 $x = 2 \text{ or } x = -5$

The solution set is $\{-5, 2\}$.

25. $|2-3x| + 2 = 9$
 $|2-3x| = 7$
 $2-3x = 7 \text{ or } 2-3x = -7$
 $-3x = 5 \text{ or } -3x = -9$
 $x = -\frac{5}{3} \text{ or } x = 3$

The solution set is $\left\{-\frac{5}{3}, 3\right\}$

26. $2x^3 = 3x^2$
 $2x^3 - 3x^2 = 0$
 $x^2(2x-3) = 0$
 $x^2 = 0 \text{ or } 2x-3 = 0$
 $x = 0 \text{ or } x = \frac{3}{2}$

The solution set is $\left\{0, \frac{3}{2}\right\}$.

27. $2x^3 + 5x^2 - 8x - 20 = 0$
 $x^2(2x+5) - 4(2x+5) = 0$
 $(2x+5)(x^2 - 4) = 0$
 $2x+5 = 0 \text{ or } x^2 - 4 = 0$
 $2x = -5 \text{ or } x^2 = 4$
 $x = -\frac{5}{2} \text{ or } x = \pm 2$

The solution set is $\left\{-\frac{5}{2}, -2, 2\right\}$.

28. $\frac{2x-3}{5} + 2 \leq \frac{x}{2}$
 $2(2x-3) + 10(2) \leq 5x$
 $4x - 6 + 20 \leq 5x$
 $14 \leq x$
 $x \geq 14$

$\{x | x \geq 14\}$ or $[14, \infty)$



Chapter 1: Equations and Inequalities

29. $-9 \leq \frac{2x+3}{-4} \leq 7$

$$36 \geq 2x + 3 \geq -28$$

$$33 \geq 2x \geq -31$$

$$\frac{33}{2} \geq x \geq -\frac{31}{2}$$

$$-\frac{31}{2} \leq x \leq \frac{33}{2}$$

$$\left\{ x \mid -\frac{31}{2} \leq x \leq \frac{33}{2} \right\} \text{ or } \left[-\frac{31}{2}, \frac{33}{2} \right]$$



30. $2 < \frac{3-3x}{12} < 6$

$$24 < 3-3x < 72$$

$$21 < -3x < 69$$

$$-7 > x > -23$$

$$\{x \mid -23 < x < -7\} \text{ or } (-23, -7)$$



31. $|3x+4| < \frac{1}{2}$

$$-\frac{1}{2} < 3x+4 < \frac{1}{2}$$

$$-\frac{9}{2} < 3x < -\frac{7}{2}$$

$$-\frac{3}{2} < x < -\frac{7}{6}$$

$$\left\{ x \mid -\frac{3}{2} < x < -\frac{7}{6} \right\} \text{ or } \left(-\frac{3}{2}, -\frac{7}{6} \right)$$




32. $|2x-5| \geq 9$

$$2x-5 \leq -9 \text{ or } 2x-5 \geq 9$$

$$2x \leq -4 \text{ or } 2x \geq 14$$

$$x \leq -2 \text{ or } x \geq 7$$

$$\{x \mid x \leq -2 \text{ or } x \geq 7\} \text{ or } (-\infty, -2] \cup [7, \infty)$$



33. $2 + |2-3x| \leq 4$

$$|2-3x| \leq 2$$

$$-2 \leq 2-3x \leq 2$$

$$-4 \leq -3x \leq 0$$

$$\frac{4}{3} \geq x \geq 0$$

$$\left\{ x \mid 0 \leq x \leq \frac{4}{3} \right\} \text{ or } \left[0, \frac{4}{3} \right]$$



34. $1 - |2-3x| < -4$

$$-|2-3x| < -5$$

$$|2-3x| > 5$$

$$2-3x < -5 \text{ or } 2-3x > 5$$

$$7 < 3x \text{ or } -3 > 3x$$

$$\frac{7}{3} < x \text{ or } -1 > x$$

$$x < -1 \text{ or } x > \frac{7}{3}$$

$$\left\{ x \mid x < -1 \text{ or } x > \frac{7}{3} \right\} \text{ or } (-\infty, -1) \cup \left(\frac{7}{3}, \infty \right)$$



35. $(6+3i) - (2-4i) = (6-2) + (3-(-4))i = 4+7i$

36. $4(3-i) + 3(-5+2i) = 12-4i-15+6i = -3+2i$

37.
$$\begin{aligned} \frac{3}{3+i} &= \frac{3}{3+i} \cdot \frac{3-i}{3-i} = \frac{9-3i}{9-3i+3i-i^2} \\ &= \frac{9-3i}{10} = \frac{9}{10} - \frac{3}{10}i \end{aligned}$$

38. $i^{50} = i^{48} \cdot i^2 = (i^4)^{12} \cdot i^2 = 1^{12}(-1) = -1$

39.
$$\begin{aligned} (2+3i)^3 &= (2+3i)^2(2+3i) \\ &= (4+12i+9i^2)(2+3i) \\ &= (-5+12i)(2+3i) \\ &= -10-15i+24i+36i^2 \\ &= -46+9i \end{aligned}$$

40. $x^2 + x + 1 = 0$

$$a = 1, b = 1, c = 1,$$

$$b^2 - 4ac = 1^2 - 4(1)(1) = 1 - 4 = -3$$

$$x = \frac{-1 \pm \sqrt{-3}}{2(1)} = \frac{-1 \pm \sqrt{3}i}{2} = -\frac{1}{2} \pm \frac{\sqrt{3}}{2}i$$

The solution set is $\left\{-\frac{1}{2} - \frac{\sqrt{3}}{2}i, -\frac{1}{2} + \frac{\sqrt{3}}{2}i\right\}$.

41. $2x^2 + x - 2 = 0$

$$a = 2, b = 1, c = -2,$$

$$b^2 - 4ac = 1^2 - 4(2)(-2) = 1 + 16 = 17$$

$$x = \frac{-1 \pm \sqrt{17}}{2(2)} = \frac{-1 \pm \sqrt{17}}{4}$$

The solution set is $\left\{\frac{-1 - \sqrt{17}}{4}, \frac{-1 + \sqrt{17}}{4}\right\}$.

42. $x^2 + 3 = x$

$$x^2 - x + 3 = 0$$

$$a = 1, b = -1, c = 3,$$

$$b^2 - 4ac = (-1)^2 - 4(1)(3) = 1 - 12 = -11$$

$$x = \frac{-(-1) \pm \sqrt{-11}}{2(1)} = \frac{1 \pm \sqrt{11}i}{2} = \frac{1}{2} \pm \frac{\sqrt{11}}{2}i$$

The solution set is $\left\{\frac{1}{2} - \frac{\sqrt{11}}{2}i, \frac{1}{2} + \frac{\sqrt{11}}{2}i\right\}$.

43. $x(1-x) = 6$

$$-x^2 + x - 6 = 0$$

$$a = -1, b = 1, c = -6,$$

$$b^2 - 4ac = 1^2 - 4(-1)(-6) = 1 - 24 = -23$$

$$x = \frac{-1 \pm \sqrt{-23}}{2(-1)} = \frac{-1 \pm \sqrt{23}i}{-2} = \frac{1}{2} \pm \frac{\sqrt{23}}{2}i$$

The solution set is $\left\{\frac{1}{2} - \frac{\sqrt{23}}{2}i, \frac{1}{2} + \frac{\sqrt{23}}{2}i\right\}$.

44. $c = 50,000 + 95x$

45. Let x represent the amount of money invested in bonds. Then $70,000 - x$ represents the amount of money invested in CD's.

Since the total interest is to be \$5000, we have:

$$0.08x + 0.05(70,000 - x) = 5000$$

$$(100)(0.08x + 0.05(70,000 - x)) = (5000)(100)$$

$$8x + 350,000 - 5x = 500,000$$

$$3x + 350,000 = 500,000$$

$$3x = 150,000$$

$$x = 50,000$$

\$50,000 should be invested in bonds at 8% and \$20,000 should be invested in CD's at 5%.

46. Using $s = vt$, we have $t = 3$ and $v = 1100$.

Finding the distance s in feet:

$$s = 1100(3) = 3300$$

The storm is 3300 feet away.

47. $1600 \leq I \leq 3600$

$$1600 \leq \frac{900}{x^2} \leq 3600$$

$$\frac{1}{1600} \geq \frac{x^2}{900} \geq \frac{1}{3600}$$

$$\frac{9}{16} \geq x^2 \geq \frac{1}{4}$$

$$\frac{3}{4} \geq x \geq \frac{1}{2}$$

The range of distances is from 0.5 meters to 0.75 meters, inclusive.

48. Let s represent the distance the plane can travel.

	With wind	Against wind
Rate	$250 + 30 = 280$	$250 - 30 = 220$
Time	$\frac{(s/2)}{280}$	$\frac{(s/2)}{220}$
Dist.	$\frac{s}{2}$	$\frac{s}{2}$

Since the total time is at most 5 hours, we have:

$$\frac{(s/2)}{280} + \frac{(s/2)}{220} \leq 5$$

$$\frac{s}{560} + \frac{s}{440} \leq 5$$

$$11s + 14s \leq 5(6160)$$

$$25s \leq 30,800$$

$$s \leq 1232$$

The plane can travel at most 1232 miles or 616 miles one way and return 616 miles.

Chapter 1: Equations and Inequalities

49. Let t represent the time it takes the helicopter to reach the raft.

	Raft	Helicopter
Rate	5	90
Time	t	t
Dist.	$5t$	$90t$

Since the total distance is 150 miles, we have:

$$5t + 90t = 150$$

$$95t = 150$$

$$t \approx 1.58 \text{ hours} \approx 1 \text{ hour and } 35 \text{ minutes}$$

The helicopter will reach the raft in about 1 hour and 35 minutes.

50. Given that $s = 1280 - 32t - 16t^2$,

- a. The object hits the ground when $s = 0$.

$$0 = 1280 - 32t - 16t^2$$

$$t^2 + 2t - 80 = 0$$

$$(t+10)(t-8) = 0$$

$$t = -10, t = 8$$

The object hits the ground after 8 seconds.

- b. After 4 seconds, the object's height is

$$s = 1280 - 32(4) - 16(4)^2 = 896 \text{ feet.}$$

51. Let t represent the time it takes Clarissa to complete the job by herself.

	Clarissa	Shawna
Time to do job alone	t	$t+5$
Part of job done in 1 day	$\frac{1}{t}$	$\frac{1}{t+5}$
Time on job (days)	6	6
Part of job done by each person	$\frac{6}{t}$	$\frac{6}{t+5}$

Since the two people paint one house, we have:

$$\frac{6}{t} + \frac{6}{t+5} = 1$$

$$6(t+5) + 6t = t(t+5)$$

$$6t + 30 + 6t = t^2 + 5t$$

$$t^2 - 7t - 30 = 0$$

$$(t-10)(t+3) = 0$$

$$t = 10 \text{ or } t = -3$$

It takes Clarissa 10 days to paint the house when working by herself.

52. Let t represent the time it takes the smaller pump to empty the tank.

	Small Pump	Large Pump
Time to do job alone	t	$t-4$
Part of job done in 1 hr	$\frac{1}{t}$	$\frac{1}{t-4}$
Time on job (hrs)	5	5
Part of job done by each pump	$\frac{5}{t}$	$\frac{5}{t-4}$

Since the two pumps empty one tank, we have:

$$\frac{5}{t} + \frac{5}{t-4} = 1$$

$$5(t-4) + 5t = t(t-4)$$

$$5t - 20 + 5t = t^2 - 4t$$

$$t^2 - 14t + 20 = 0$$

We can solve this equation for t by using the quadratic formula:

$$t = \frac{-(-14) \pm \sqrt{(-14)^2 - 4(1)(20)}}{2(1)}$$

$$= \frac{14 \pm \sqrt{116}}{2} = \frac{14 \pm 2\sqrt{29}}{2}$$

$$= 7 \pm \sqrt{29} \approx 7 + 5.385$$

$$t = 12.385 \text{ or } t = 1.615 \text{ (not feasible)}$$

It takes the small pump approximately 12.385 hours (12 hr 23 min) to empty the tank.

53. Let x represent the amount of water added.

% salt	Tot. amt.	amt. of salt
10%	64	(0.10)(64)
0%	x	(0.00)(x)
2%	$64+x$	(0.02)($64+x$)

$$(0.10)(64) + (0.00)(x) = (0.02)(64+x)$$


$$6.4 = 1.28 + 0.02x$$

$$5.12 = 0.02x$$

$$x = 256$$

256 ounces of water must be added.

54. Consider the diagram



By the Pythagorean Theorem we have

$$w^2 + (w+2)^2 = (10)^2$$

$$w^2 + w^2 + 4w + 4 = 100$$

$$2w^2 + 4w - 96 = 0$$

$$w^2 + 2w - 48 = 0$$

$$(w+8)(w-6) = 0$$

$$w = -8 \text{ or } w = 6$$

The width is 6 inches and the length is $6 + 2 = 8$ inches.

55. Let x represent the amount of the 15% solution added.

% acid	tot. amt.	amt. of acid
40%	60	(0.40)(60)
15%	x	(0.15)(x)
25%	$60+x$	(0.25)(60+ x)

$$(0.40)(60) + (0.15)(x) = (0.25)(60+x)$$

$$24 + 0.15x = 15 + 0.25x$$

$$9 = 0.1x$$

$$x = 90$$

90 cubic centimeters of the 15% solution must be added, producing 150 cubic centimeters of the 25% solution.


56. a. Consider the following diagram:

$$4(s+6) = 50$$

$$4s + 24 = 50$$

$$4s = 26$$


$$s = 6.5$$



The painting is 6.5 inches by 6.5 inches.

$s+6 = 12.5$, so the frame is 12.5 inches by 12.5 inches.

- b. Consider the following diagram:



$$2(2w+6) + 2(w+6) = 50$$

$$4w+12 + 2w+12 = 50$$

$$6w = 26$$

$$w = \frac{26}{6} = 4\frac{1}{3}$$

$$l = 2w = 8\frac{2}{3}$$

The painting is $8\frac{2}{3}$ inches by $4\frac{1}{3}$ inches.

The frame is $14\frac{2}{3}$ inches by $10\frac{1}{3}$ inches.

57. Let x represent the amount Scott receives. Then $\frac{3}{4}x$ represents the amount Alice receives and $\frac{1}{2}x$ represents the amount Tricia receives. The total amount is \$900,000, so we have:

$$x + \frac{3}{4}x + \frac{1}{2}x = 900,000$$

$$4\left(x + \frac{3}{4}x + \frac{1}{2}x\right) = 4(900,000)$$

$$4x + 3x + 2x = 3,600,000$$

$$9x = 3,600,000$$

$$x = 400,000$$

$$\text{So, } \frac{3}{4}x = \frac{3}{4}(400,000) = 300,000 \text{ and}$$

$$\frac{1}{2}x = \frac{1}{2}(400,000) = 200,000.$$

Scott receives \$400,000, Alice receives \$300,000, and Tricia receives \$200,000.

58. Let t represent the time it takes the older machine to complete the job by itself.

	Old copier	New copier
Time to do job alone	t	$t-1$
Part of job done in 1 hr	$\frac{1}{t}$	$\frac{1}{t-1}$
Time on job (hrs)	1.2	1.2
Part of job done by each copier	$\frac{1.2}{t}$	$\frac{1.2}{t-1}$

Since the two copiers complete one job, we have:

$$\frac{1.2}{t} + \frac{1.2}{t-1} = 1$$

$$1.2(t-1) + 1.2t = t(t-1)$$

$$1.2t - 1.2 + 1.2t = t^2 - t$$

$$t^2 - 3.4t + 1.2 = 0$$

$$5t^2 - 17t + 6 = 0$$

$$(5t-2)(t-3) = 0$$

$$t = 0.4 \text{ or } t = 3$$

It takes the old copier 3 hours to do the job by itself. (0.4 hour is impossible since together it takes 1.2 hours.)

59. Let r_S represent Scott's rate and let r_T represent Todd's rate. The time for Scott to run 95 meters is the same as for Todd to run 100 meters.

$$\frac{95}{r_S} = \frac{100}{r_T}$$

$$r_S = 0.95r_T$$

$$d_S = t \cdot r_S = t(0.95r_T) = 0.95d_T$$

If Todd starts from 5 meters behind the start:

$$d_T = 105$$

$$d_S = 0.95d_T = 0.95(105) = 99.75$$

- a. The race does not end in a tie.
- b. Todd wins the race.
- c. Todd wins by 0.25 meters.
- d. To end in a tie:

$$100 = 0.95(100 + x)$$

$$100 = 95 + 0.95x$$

$$5 = 0.95x$$

$$x \approx 5.26 \text{ meters}$$

- e. $95 = 0.95(100)$ Therefore, the race ends in a tie.

60. The effective speed of the train (i.e., relative to the man) is $30 - 4 = 26$ miles per hour. The time

$$\text{is } 5 \text{ sec} = \frac{5}{60} \text{ min} = \frac{5}{3600} \text{ hr} = \frac{1}{720} \text{ hr.}$$

$$s = vt$$

$$= 26 \left(\frac{1}{720} \right)$$

$$= \frac{26}{720} \text{ miles}$$

$$= \frac{26}{720} \cdot 5280 \approx 190.67 \text{ feet}$$

The freight train is about 190.67 feet long.

Chapter 1 Test

$$\begin{aligned} 1. \quad \frac{2x}{3} - \frac{x}{2} &= \frac{5}{12} \\ 12 \left(\frac{2x}{3} - \frac{x}{2} \right) &= 12 \left(\frac{5}{12} \right) \\ 8x - 6x &= 5 \\ 2x &= 5 \\ x &= \frac{5}{2} \end{aligned}$$

The solution set is $\left\{\frac{5}{2}\right\}$.

$$\begin{aligned} 2. \quad x(x-1) &= 6 \\ x^2 - x &= 6 \\ x^2 - x - 6 &= 0 \\ (x-3)(x+2) &= 0 \\ x-3 = 0 \text{ or } x+2 &= 0 \\ x = 3 \text{ or } x &= -2 \end{aligned}$$

The solution set is $\{-2, 3\}$.

$$\begin{aligned} 3. \quad x^4 - 3x^2 - 4 &= 0 \\ (x^2 - 4)(x^2 + 1) &= 0 \\ x^2 - 4 = 0 \text{ or } x^2 + 1 &= 0 \\ x^2 = 4 \text{ or } x^2 &= -1 \\ x = \pm 2 \text{ or } &\text{Not real} \end{aligned}$$

The solution set is $\{-2, 2\}$.

$$\begin{aligned} 4. \quad \sqrt{2x-5} + 2 &= 4 \\ \sqrt{2x-5} &= 2 \\ (\sqrt{2x-5})^2 &= (2)^2 \\ 2x-5 &= 4 \\ 2x &= 9 \\ x &= \frac{9}{2} \end{aligned}$$

$$\text{Check: } \sqrt{2\left(\frac{9}{2}\right) - 5} + 2 = 4$$

$$\sqrt{9-5} + 2 = 4$$

$$\sqrt{4} + 2 = 4$$

$$2+2=4$$

$$4=4$$

The solution set is $\left\{\frac{9}{2}\right\}$.

5. $|2x-3|+7=10$

$$|2x-3|=3$$

$$2x-3=3 \quad \text{or} \quad 2x-3=-3$$

$$2x=6 \quad \text{or} \quad 2x=0$$

$$x=3 \quad \text{or} \quad x=0$$

The solutions set is $\{0, 3\}$.

6. $3x^3 + 2x^2 - 12x - 8 = 0$

$$x^2(3x+2) - 4(3x+2) = 0$$

$$(x^2 - 4)(3x+2) = 0$$

$$(x+2)(x-2)(3x+2) = 0$$

$$x+2=0 \quad \text{or} \quad x-2=0 \quad \text{or} \quad 3x+2=0$$

$$x=-2 \quad \text{or} \quad x=2 \quad \text{or} \quad x=-\frac{2}{3}$$

The solution set is $\left\{-2, -\frac{2}{3}, 2\right\}$.

7. $3x^2 - x + 1 = 0$

$$x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(3)(1)}}{2(3)}$$

$$= \frac{1 \pm \sqrt{-11}}{6} \quad (\text{Not real})$$

This equation has no real solutions.

8. $-3 \leq \frac{3x-4}{2} \leq 6$

$$2(-3) \leq 2\left(\frac{3x-4}{2}\right) \leq 2(6)$$

$$-6 \leq 3x-4 \leq 12$$

$$-2 \leq 3x \leq 16$$

$$-\frac{2}{3} \leq x \leq \frac{16}{3}$$

$$\left\{x \mid -\frac{2}{3} \leq x \leq \frac{16}{3}\right\} \text{ or } \left[-\frac{2}{3}, \frac{16}{3}\right]$$

9. $|3x+4| < 8$

$$-8 < 3x+4 < 8$$

$$-12 < 3x < 4$$

$$-4 < x < \frac{4}{3}$$

$$\left\{x \mid -4 < x < \frac{4}{3}\right\} \text{ or } \left(-4, \frac{4}{3}\right)$$

10. $2 + |2x-5| \geq 9$

$$|2x-5| \geq 7$$

$$2x-5 \leq -7 \quad \text{or} \quad 2x-5 \geq 7$$

$$2x \leq -2 \quad \text{or} \quad 2x \geq 12$$

$$x \leq -1 \quad \text{or} \quad x \geq 6$$

$$\{x \mid x \leq -1 \text{ or } x \geq 6\} \text{ or } (-\infty, -1] \cup [6, \infty).$$

11. $\frac{-2}{3-i} = \frac{-2}{3-i} \cdot \frac{3+i}{3+i} = \frac{-6-2i}{9+3i-3i-i^2} = \frac{-6-2i}{9-(-1)}$

$$= \frac{-6-2i}{10} = \frac{-3-i}{5} = -\frac{3}{5} - \frac{1}{5}i$$

12. $4x^2 - 4x + 5 = 0$

$$x = \frac{-(-4) \pm \sqrt{(-4)^2 - 4(4)(5)}}{2(4)}$$

$$= \frac{4 \pm \sqrt{-64}}{8} = \frac{4 \pm 8i}{8} = \frac{1}{2} \pm i$$

This solution set is $\left\{\frac{1}{2} - i, \frac{1}{2} + i\right\}$.

Chapter 1: Equations and Inequalities

13. Let x represent the amount of the \$8-per-pound coffee.

Amt. of coffee (pounds)	Price (\$)	Total \$
20	4	(20)(4)
x	8	(8)(x)
$20+x$	5	(5)(20+x)

$$80 + 8x = (5)(20+x)$$

$$80 + 8x = 100 + 5x$$

$$3x = 20$$

$$x = \frac{20}{3} = 6\frac{2}{3}$$

Add $6\frac{2}{3}$ pounds of \$8/lb coffee to get $26\frac{2}{3}$ pounds of \$5/lb coffee.

3. $T = 0.15$ board per second

$$0.15 = \frac{1}{0.2p+2}$$

$$(0.2p+2)(0.15) = 1$$

$$0.03p + 0.3 = 1$$

$$0.03p = 0.7$$

$$p \approx 23.3 \text{ parts per board}$$

Thus, only 23 parts per board will work.

For problems 4 – 6, C is requested, so solve for C first:

$$T = \frac{n}{Cnp + L + M}$$

$$(Cnp + L + M)T = n$$

$$CnpT + LT + MT = n$$

$$CnpT = n - LT - MT$$

$$C = \frac{n - LT - MT}{npT}$$

4. $T = 0.06, n = 3, p = 100, M = 1, L = 5$

$$C = \frac{3 - 5(0.06) - 1(0.06)}{3(100)(0.06)} \approx 0.147 \text{ sec}$$

5. $T = 0.06, n = 3, p = 150, M = 1, L = 5$

$$C = \frac{3 - 5(0.06) - 1(0.06)}{3(150)(0.06)} \approx 0.098 \text{ sec}$$

6. $T = 0.06, n = 3, p = 200, M = 1, L = 5$

$$C = \frac{3 - 5(0.06) - 1(0.06)}{3(200)(0.06)} \approx 0.073 \text{ sec}$$

7. As the number of parts per board increases, the tact time decreases, if all the other factors remain constant.

Chapter 1 Projects

Project I

Internet Based Project

Project II

1. $T = \frac{n}{Cnp + L + M}, n = 3, L = 5, M = 1, C = 0.2$

$$T = \frac{3}{0.2(3)p + 5 + 1} = \frac{3}{0.6p + 6} = \frac{1}{0.2p + 2}$$

2. All of the times given in problem 1 were in seconds, so $T = 0.1$ board per second needs to be used as the value for T in the equation found in problem 1.

$$0.1 = \frac{1}{0.2p + 2}$$

$$(0.2p+2)(0.1) = 1$$

$$0.02p + 0.2 = 1$$

$$0.02p = 0.8$$

$$p = 40 \text{ parts per board}$$